

STS SCA Data Specifications v2.9
Data Collection Form fields:
Updated: October, 2017

Introduction

This manual is intended to clarify field definition and intent. This document contains the most up to date instructions for v. 2.9 data abstraction. Do not refer to old manuals or other data definitions. **Please review this document prior to submitting clinical questions.** FAQs will be added to the document in red to provide additional examples and clarification. Please do not print this document since it will change frequently. Using the web version will ensure that you have the most up to date information. Occasionally there may be changes or important information that will be highlighted here and will be also included in STS Database Newsletters. Use the Ctrl + F function to search for a number or term of interest. Bookmarks have been added for October, 2017 updates.

General Information:

The STS data collection forms should be held for two years.
If you only collect data directly to the software you are not required to create data collection forms to save.

For all questions where the choices include “no” and “unknown,” how should the question be coded when there is no specific documentation?

When a history and physical or a consultation exists in the medical record and the values are not specifically addressed in the documentation, code no. Unknown should be coded only in the circumstance where no clinical documentation exists and the patient cannot give history and supportive documentation.

Will the STS plan to extract data from the EMR?

Some but not all could be collected from the EMR. Some fields are too complex to extract from the EMR in real time. Caution needs to be used with auto data transfer from the EMR.

Is there a list of the procedures that should be included in the Adult Cardiac Surgery Database?
While there is no all-inclusive list of procedures to be included, all procedures must include a surgeon that is listed in the participation agreement with the STS.

Administrative

Long Name: Software Vendor Identifier

Short Name: VendorID

Definition: Name (assigned by STS) given to identify software vendor (up to 8 characters). Vendors should use standard name identification across sites. Changes to Vendor Name Identification must be approved by the STS.

Intent/Clarification:

Name must match what is listed as the Active vendor for your Participant ID in the database. Any mismatch will cause your data file submission not to process.

SEQ. #: 10

Long Name: Software Version

Short Name: SoftVrsn

Definition: Vendor's software product name and version number identifying the software which created this record. Vendor controls the value in this field. Version passing certification/harvest testing will be noted at warehouse.

Intent/Clarification: -

SEQ. #: 15

Long Name: STS Data Version

Short Name: DataVrsn

Definition: Version number of the STS Data Specifications/Dictionary, to which each record conforms. It will identify which fields should have data, and what are the valid data for each field. This must be entered into the record automatically by the software.

Intent/Clarification:

Data version must be appropriate for the procedure date listed in the record. Valid date ranges can be found in the current Software Specifications.

Any mismatch will cause your data file submission not to process.

SEQ. #: 20

Long Name: On-Demand Files Version Number

Short Name: OnDemandVrsn

Definition: The version number of the On-Demand lists in use at the time this data record was created or edited. The value is inserted into the record at the time the record is created or is modified by the user. The version numbers will be specified by the STS.

Intent/Clarification:

Inconsistencies here do not prevent your file from being processed. However, any mismatch will appear in your Data Quality Report (DQR) as a value that could not be

interpreted. You should contact your designated Data Submission Coordinator for assistance.

<http://www.sts.org/sts-national-database/database-managers/contact-information>

SEQ. #: 25

Long Name: Participant ID

Short Name: ParticID

Definition: Participant ID is a unique number assigned to each database participant by the STS. A database participant is defined as one entity that signs a Participation Agreement with the STS, submits one data file to the harvest, and gets back one report on their data. The participant ID must be entered into each record.

Each participant's data if submitted to harvest must be in one data file. If one participant keeps their data in more than one file (e.g. at two sites), then the participant must combine them back into one file for harvest submission.

If two or more participants share a single purchased software, and enter cases into one database, then the data must be extracted into two different files, one for each participant ID, with each record having the correct participant ID number.

Intent/Clarification:

Each participant's data, if submitted to harvest, must be in one data file. If one participant keeps the data in more than one file (e.g. at two sites), then the participant must combine them back into one file for harvest submission. If two or more participants share a single purchased software, and enter cases in to one database, then the data must be extracted into two different files, one for each participant ID, with each record having the correct participant ID number.

SEQ. #: 30

Long Name: Record ID

Short Name: RecordID

Definition: An arbitrary, unique value generated by the software that permanently identifies each record in the participant's database (note that unlike the PatID value, this does not identify the individual patient). The value of the identifier is a combination of a code assigned to the software developer by the STS, and a value generated by the software to create a unique value. Once assigned to a record, this value can never be changed or reused. The data warehouse will use this value to communicate issues about individual records with the participant. It may also be used by the data warehouse to link this record to other clinical data.

Intent/Clarification:

The data warehouse will use this value to communicate issues about individual records with the participant. It may also be used by the data warehouse to link this record to other clinical data.

SEQ. #: 35

Long Name: Cost Link

Short Name: CostLink

Definition: A participant specified alpha-numeric code that can be used to link this record's clinical data with the participant's cost information for this patient admission. This information may be used in the future to perform procedure cost analysis (for which the actual cost data would have to be harvested separately). The value in this field must not be the patient's Medical Record Number, Social Security Number or any other patient identifying value.

Intent/Clarification:

This information may be used in the future to perform procedure cost analysis, for which the actual cost data would have to be harvested separately. The value in this field must not be the patient's Medical Record Number, Social Security Number or any other patient identifying value.

SEQ. #: 40

Long Name: Patient ID

Short Name: PatID

Definition: An arbitrary value, (not a recognizable ID like Social Security Number or Medical Record Number) that uniquely and permanently identifies each patient. The value of the identifier is a combination of a code assigned to the software developer by the STS, and a value generated by the software to create a unique value. Once assigned to a patient, this can never be changed or reused. If a patient is admitted to the hospital more than once, each record for that patient will have the same value in this field.

Intent/Clarification:

The value of the identifier is a combination of a code assigned to the software developer by the STS, and a value generated by the software to create a unique value. Once assigned to a patient, this can never be changed or reused. If a patient is admitted to the hospital more than once, each record for that patient will have the same value in this field.

SEQ. #: 45

Long Name: Patient Participating In STS-Related Clinical Trial

Short Name: ClinTrial

Definition: Indicate which, if any, STS-related clinical trial in which the patient is participating. The STS will assign a code to each clinical trial as they begin collecting data.

Intent/Clarification:

A list of trials will be posted as they are started.

There are currently no STS trials underway in the Adult Cardiac Surgery Database.

SEQ. #: 46

Long Name: Patient Participating In STS-Related Clinical Trial - Patient ID

Short Name: ClinTrialPatID

Definition: Indicate the patient identifier used to identify the patient in the clinical trial.

Intent/Clarification: Instructions will be provided for each trial.

Demographics

SEQ. #: 50

Long Name: Patient Last Name

Short Name: PatLName

Definition: Indicate the patient's last name documented in the medical record. This field should be collected in compliance with state/local privacy laws.

Intent/Clarification: -

SEQ. #: 55

Long Name: Patient First Name

Short Name: PatFName

Definition: Indicate the patient's first name documented in the medical record. This field should be collected in compliance with state/local privacy laws.

Intent/Clarification: -

SEQ. #: 60

Long Name: Patient Middle Name

Short Name: PatMName

Definition: Indicate the patient's middle name as documented in the medical record. Leave "blank" if no middle name. This field should be collected in compliance with state/local privacy laws.

Intent/Clarification: -

SEQ. #: 65

Long Name: Date of Birth **Short Name:** DOB

Definition: Indicate the patient's date of birth using 4-digit format for year. This field should be collected in compliance with state/local privacy laws.

Intent/Clarification: Required date format: mm/dd/yyyy

SEQ. #: 70

Long Name: Patient Age

Short Name: Age

Definition: Indicate the patient's age in years, at time of surgery. This should be calculated from the date of birth and the date of surgery, according to the convention used in the USA (the number of birthdate anniversaries reached by the date of surgery). If age is less than 18, the data record will be accepted into the database, but will not be included in the national analysis and report.

Intent/Clarification: -

SEQ. #: 75

Long Name: Sex

Short Name: Gender

Definition: Indicate the patient's sex at birth as either male or female.

Intent/Clarification:

Patients who have undergone gender reassignment surgery maintain the risk associated with their chromosomal gender.

SEQ. #: 76

Long Name: National Identification (Social Security Number) Known

Short Name: SSNKnown

Definition: Indicate whether the patient's National Identification Number is known or if the patient refused to provide this information.

Intent/Clarification: - Refused means the patient did not wish to share the information. No means the information was not available or the participant site did not wish to provide.

SEQ. #: 80

Long Name: National Identification (Social Security Number)

Short Name: SSN

Definition: Indicate the patient's National Identification Number. Although this is the Social Security Number in the USA, other countries may have a different National Patient Identifier Number. For example in Canada, this would be the Social Insurance Number.

This field should be collected in compliance with state/local privacy laws.

Intent/Clarification: -

SEQ. #: 85

Long Name: Medical Record Number

Short Name: MedRecN

Definition: Indicate the patient's medical record number at the hospital where surgery occurred. This field should be collected in compliance with state/local privacy laws.

Intent/Clarification: -

SEQ. #: 90

Long Name: Patient's Street Address

Short Name: PatAddr

Definition: Indicate the street address at which the patient resides at time of admission. If patient is homeless, enter "Homeless".

This field should be collected in compliance with state/local privacy laws.

Intent/Clarification:

This may be a hotel or relative's home if the patient is not a local resident. This will track referrals and assist with follow-up. A post office box may be used if no other address is available.

SEQ. #: 95

Long Name: Patient's City

Short Name: PatCity

Definition: Indicate the city in which the patient resides at time of admission.

This field should be collected in compliance with state/local privacy laws.

Intent/Clarification: -

SEQ. #: 100

Long Name: Patient's Region

Short Name: PatRegion

Definition: Indicate the region of the country (i.e., state or province) in which the patient resides at time of admission.

Intent/Clarification:

Regional information is used to assess disparities in health care delivery.

SEQ. #: 105

Long Name: Patient's ZIP Code

Short Name: PatZIP

Definition: Indicate the ZIP Code of the patient's local residence. Outside the USA, this data may be known by other names such as Postal Code.

This field should be collected in compliance with state/local privacy laws.

Intent/Clarification:

Regional information is used to assess disparities in health care delivery.

SEQ. #: 115

Long Name: Patient's Country

Short Name: PatientCountry

Definition: Indicate the patient's country of residence at time of admission.

This field should be collected in compliance with state/local privacy laws.

Intent/Clarification:

List of country codes found in Data Specifications V2.9 (p.9)

SEQ. #: 120

Long Name: Permanent Address

Short Name: PermAddr

Definition: Indicate whether the patient considers the given address to be their permanent address.

Intent/Clarification:

The intent is to identify patients who travel outside their local area for treatment. CMS is tracking disparities in health care delivery and looking at underserved areas. This also assists with long term follow up locally.

SEQ. #: 150

Long Name: Race Documented
Short Name: RaceDocumented
Definition: Indicate whether race is documented

Intent/Clarification:

Race should be self-reported by the patient/family. Do not assign race or make assumptions if race is not documented.

SEQ. #: 155

Long Name: Race - White
Short Name: RaceCaucasian

Definition: Indicate whether the patient's race, as determined by the patient or family, includes White. "White" refers to a person having origins in any of the original peoples of Europe, the Middle East, or North Africa. It includes people who indicated their race(s) as "White" or reported entries such as Irish, German, Italian, Lebanese, Arab, Moroccan, or Caucasian. [The 2010 Census Redistricting Data (Public Law 94-171) Summary File]

Intent/Clarification:

The Census Bureau collects race data in accordance with guidelines provided by the U.S. Office of Management and Budget, these data are based on self-identification. The racial categories included in the census form generally reflect a social definition of race recognized in this country and are not an attempt to define race biologically, anthropologically or genetically. In addition, it is recognized that categories of the race item include racial and national origin or socio-cultural groups. People may choose to report more than one race to indicate their racial mixture, such as American Indian and White.

People who identify their origin (ETHNICITY) as Hispanic, Latino or Spanish may be of any race. In addition, it is recognized that the categories of the race item include both racial and national origin and socio-cultural groups. You may choose more than one race category.

SEQ. #: 160

Long Name: Race - Black / African American
Short Name: RaceBlack

Definition: Indicate whether the patient's race, as determined by the patient or family, includes Black / African American. "Black or African American" refers to a person having origins in any of the Black racial groups of Africa. It includes people who indicated their race(s) as "Black, African Am., or Negro" or reported entries such as African American, Kenyan, Nigerian, or Haitian. [The 2010 Census Redistricting Data (Public Law 94-171) Summary File]

Intent/Clarification:

This includes a person having origins in any of the black racial groups of Africa. Terms such as "Haitian" or "Negro" can be used in addition to "Black or African American."

Definition source: Standards for Maintaining, Collecting and Presenting Federal Data on Race and Ethnicity: The minimum categories for data on race and ethnicity for Federal statistics, program administrative reporting and civil rights compliance reporting.

SEQ. #: 165

Long Name: Race - Asian

Short Name: RaceAsian

Definition: Indicate whether the patient's race, as determined by the patient or family, includes Asian. "Asian" refers to a person having origins in any of the original peoples of the Far East, Southeast Asia, or the Indian subcontinent, including, for example, Cambodia, China, India, Japan, Korea, Malaysia, Pakistan, the Philippine Islands, Thailand, and Vietnam. It includes people who indicated their race(s) as "Asian" or reported entries such as "Asian Indian", "Chinese", "Filipino", "Korean", "Japanese", "Vietnamese", and "Other Asian" or provided other detailed Asian responses. [The 2010 Census Redistricting Data (Public Law 94-171) Summary File]

Intent/Clarification: -

Definition source: Standards for Maintaining, Collecting and Presenting Federal Data on Race and Ethnicity: The minimum categories for data on race and ethnicity for Federal statistics, program administrative reporting and civil rights compliance reporting.

SEQ. #: 170

Long Name: Race - American Indian / Alaskan Native

Short Name: RaceNativeAm

Definition: Indicate whether the patient's race, as determined by the patient or family, includes American Indian / Alaskan Native. "American Indian or Alaska Native" refers to a person having origins in any of the original peoples of North and South America (including Central America) and who maintains tribal affiliation or community attachment. This category includes people who indicated their race(s) as "American Indian or Alaska Native" or reported their enrolled or principle tribe, such as Navajo, Blackfeet, Inupiat, Yup'ik, or Central American Indian groups or South American Indian groups. [The 2010 Census Redistricting Data (Public Law 94-171) Summary File]

Intent/Clarification:

This includes all in North American native peoples such as American Indian/Alaskan Native, Inuit.

SEQ. #: 175

Long Name: Race - Native Hawaiian / Pacific Islander

Short Name: RacNativePacific

Definition: Indicate whether the patient's race, as determined by the patient or family, includes Native Hawaiian / Pacific Islander. "Native Hawaiian or Other Pacific Islander" refers to a person having origins in any of the original peoples of Hawaii, Guam, Samoa, or other Pacific Islands. It includes people who indicated their race(s) as "Pacific Islander" or reported entries such as "Native Hawaiian", "Guamanian or Chamorro", "Samoaan", and "Other Pacific Islander" or provided other detailed Pacific Islander responses. [The 2010 Census Redistricting Data (Public Law 94-171) Summary File]

Intent/Clarification: -

Definition source: Standards for Maintaining, Collecting and Presenting Federal Data on Race and Ethnicity. The minimum categories for data on race and ethnicity for Federal statistics, program administrative reporting and civil rights compliance reporting.

SEQ. #: 180

Long Name: Race - Other

Short Name: RaceOther

Definition: Indicate whether the patient's race, as determined by the patient or family, includes any other race. "Some Other Race" includes all other responses not included in the White, Black or African American, American Indian or Alaska Native, Asian, and Native Hawaiian or Other Pacific Islander race categories described above. [The 2010 Census Redistricting Data (Public Law 94-171) Summary File]

Intent/Clarification: -

SEQ. #: 185

Long Name: Hispanic or Latino or Spanish Ethnicity

Short Name: Ethnicity

Definition: Indicate if the patient is of Hispanic, Latino or Spanish ethnicity as reported by the patient / family. "Hispanic, Latino or Spanish" refers to a person of Cuban, Mexican, Puerto Rican, South or Central American, or other Spanish culture or origin regardless of race. [The 2010 Census Redistricting Data (Public Law 94-171) Summary File]

Intent/Clarification:

People who identify their origin as Hispanic, Latino or Spanish **may be of any race.** Do not make assumptions about ethnicity if it is not documented in the medical record.

Hospitalization

SEQ. #: 205

Long Name: Hospital Name

Short Name: HospName

Definition: Indicate the full name of the facility where the procedure was performed. Values should be full, official hospital name as it appears on the contract with the STS, with no abbreviations or variations in spelling for a single hospital. Values should also be in mixed-case.

Intent/Clarification:

User maintains list of valid values. New values are made available through a utility that is separate from entering a data record.

This must match what DCRI and STS have on your contract. If this name does not match your file will be rejected. Please update if any changes occur.

Update Hospital and Surgeon information here:

http://www.sts.org/sites/default/files/documents/Updated_STSContactUpdateForm_FIN_AL_03142017_0.pdf

SEQ. #: 210

Long Name: Hospital ZIP Code **Short Name:** HospZIP

Definition: Indicate the ZIP Code of the hospital. Outside the USA, these data may be known by other names such as Postal Code. This field should be collected in compliance with state/local privacy laws.

Intent/Clarification: -

SEQ. #: 215

Long Name: Hospital Region

Short Name: HospStat

Definition: Indicate the region of the country (i.e., state or province) in which the hospital is located.

Intent/Clarification: -

SEQ. #: 220

Long Name: Hospital National Provider Identifier

Short Name: HospNPI

Definition: Indicate the hospital's National Provider Identifier (NPI). This number, assigned by the Center for Medicare and Medicaid Services (CMS), is used to uniquely identify facilities for Medicare billing purposes.

Non-US participants will have a unique hospital ID number assigned by STS.

Intent/Clarification:

STS/DCRI maintains a list of Hospital NPIs associated with Participation Agreements. Data files that include other hospitals cannot be processed. **This is different from the Surgeon NPI.** <https://nppes.cms.hhs.gov/NPPES/NPIRegistryHome.do>. **If the field is missing or incorrect, the file will not be processed.**

If the hospital NPI is changed (e.g. thru mergers/acquisitions) it is crucial that STS and DCRI be notified as soon as possible. This will ensure records are handled appropriately at harvest.

Update Hospital and Surgeon information here:

http://www.sts.org/sites/default/files/documents/Updated_STSContactUpdateForm_FIN_AL_03142017_0.pdf

SEQ. #: 221

Long Name: Hospital CMS Certification Number

Short Name: HospCMSCert

Definition: Indicate the hospital's CMS certification number

Intent/Clarification:

In order to avoid confusion with the NPI, the Medicare/Medicaid Provider Number, also known as the OSCAR Provider Number, Medicare Identification Number or Provider Number) has been renamed the **CMS Certification Number (CCN)**.

FAQ August 2017: Where can I find the CMS certification number for my site?

Answer: Contact your medical records billing department for the number. You can also look for the number on the CMS website at <https://www.ahd.com/search.php>

SEQ. #: 291

Long Name: Primary Payor

Short Name: PayorPrim

Definition: Indicate the primary insurance payor for this admission.

Intent/Clarification:

When there is more than one payor, the primary payor pays first.

FAQ August 2017: How is a Medicare/Medicaid managed care product coded (ie. Humana Medicare, Star Molina Medicaid)?

Answer: The Medicare/Medicaid managed care products are captured in the primary payor category as Medicare or Medicaid only.

SEQ. #: 292

Long Name: Primary Payor Medicare Fee For Service

Short Name: PrimMCareFFS

Definition: Indicate whether the patient is covered by Medicare Fee For Service (Part B).

Intent/Clarification: -

SEQ. #: 293

Long Name: Secondary (Supplemental) Payor

Short Name: PayorSecond

Definition: Indicate which if any secondary insurance payor was used for this admission.

Intent/Clarification:

When there is more than one payor, the secondary payor pays after the primary payor.

FAQ August 2017: How is a Medicare/Medicaid managed care product coded (ie. Humana Medicare, Star Molina Medicaid)?

Answer: The Medicare/Medicaid managed care products are captured in the secondary payor category as Medicare or Medicaid only.

SEQ. #: 294

Long Name: Secondary Payor Medicare Fee For Service

Short Name: SecondMCareFFS

Definition: Indicate whether the patient is covered by Medicare Fee For Service (Part B).

Intent/Clarification: -

FAQ August 2017: When is secondary Medicare FFS coded?

Answer: Code a secondary Medicare FFS provider only when the FFS ID number is different from the primary Medicare FFS provider number.

SEQ. #: 305

Long Name: Date of Admission

Short Name: AdmitDt

Definition: Indicate the Date of Admission. For those patients who originally enter the hospital in an out-patient capacity (i.e., catheterization), the admit date is the date the patient's status changes to in-patient. In the event admission date comes after date of surgery, use date of surgery.

Intent/Clarification: Required date format: mm/dd/yyyy

SEQ. #: 310

Long Name: Date of Surgery

Short Name: SurgDt

Definition: Indicate the date of index cardiac surgical procedure. Index cardiac surgical procedure is defined as the initial major cardiac surgical procedure of the hospitalization.

Intent/Clarification:

The date the patient enters the operating room for surgery.

Required date format: mm/dd/yyyy

SEQ. #: 320

Long Name: Admit Source

Short Name: AdmitSrc

Definition: Indicate the source of admission for the patient to your facility.

Intent/Clarification:

Choose elective admission, through the ED, transferred in from another acute care facility or “other,” which includes transfers from non-acute care facilities such as nursing homes.

If a patient is admitted for an elective catheterization and is then held-over for surgery (elective or urgent), this should be coded as an elective admission; however, the surgery status should be coded as urgent based on the catheterization findings.

If the facility has a stand-alone “feeder” ER (with the same patient ID) then the source is ED.

The option “Other” includes **direct admits** from MD offices, providers, non-acute clinics, Rehab units. However, if patients is sent to the ED then ED should be selected as admit source.

SEQ. #: 325

Long Name: Other Hospital Performs Cardiac Surgery

Short Name: OthHosCS

Definition: The transferring hospital has the necessary personnel and facilities to have been able to perform cardiac surgery.

Intent/Clarification:

The intent is to capture patients whose acuity requires a higher level of care or more complex procedure than can be provided at the transferring facility, such as a transplant. The goal is to identify high acuity patients and does not reflect negatively on the referring hospital. Code “yes” if the transferring hospital performs heart surgery,

even if it is not the type of surgery the patient is being transferred for such as transplant or VAD.

Risk Factors

General Information:

If the patient is alone, intubated and unable to give history; use the information from the patient's family if they become available.

SEQ. #: 330

Long Name: Height (cm)

Short Name: HeightCm

Definition: Indicate the height of the patient in centimeters.

Intent/Clarification:

Used to calculate BSA (body surface area) and is a field for risk calculation. 1 inch = 2.54 cm.

For patients who have had lower extremity amputations, code the patient's original height.

SEQ. #: 335

Long Name: Weight (kg)

Short Name: WeightKg

Definition: Indicate the weight of the patient in kilograms closest to the date of procedure.

Intent/Clarification:

Used to calculate BSA (body surface area) and is a field for risk calculation. Record in kilograms. 1 Kg = 2.2 pounds.

SEQ. #: 355

Long Name: RF-Family History of Premature CAD

Short Name: FHCAD

Definition: Indicate if the patient has any direct blood relatives (parents, siblings, children) who have had any of the following at age <55 y for male relatives or <65 y for female relatives:

- Angina
- Acute MI
- Sudden cardiac death without obvious cause
- CABG surgery
- PCI

Intent/Clarification:

The disease, treatment (surgical, non-surgical or medical) and/or symptoms must have been present or reported to have occurred prior to age 55 in males and 65 in females. This is considered a strong predictor for development of CAD and may include, but is not limited to, angina, acute MI, CABG, PCI or sudden cardiac death with no known cause. Early onset of CAD in patient and or first generation family members predisposes patient to increased risk of mortality/morbidity.

Code family history as “No” if the patient is adopted and family history is unknown.

You must have the exact age (not age range or approximation) to document premature CAD.

SEQ. #: 360

Long Name: RF-Diabetes

Short Name: Diabetes

Definition: History of diabetes diagnosed and/or treated by a healthcare provider. The American Diabetes Association criteria include documentation of the following:

1. Hemoglobin A1c $\geq 6.5\%$; or
2. Fasting plasma glucose ≥ 126 mg/dL (7.0 mmol/L); or
3. 2-h Plasma glucose ≥ 200 mg/dL (11.1 mmol/L) during an oral glucose tolerance test; or
4. In a patient with classic symptoms of hyperglycemia or hyperglycemic crisis, a random plasma glucose ≥ 200 mg/dL (11.1 mmol/L)

This does not include gestational diabetes.

2013 ACCF/AHA Data Standards
Cannon et al. JACC Vol. 61, No. 9, 2013

Intent/Clarification:

Indicate if the patient has a history of diabetes mellitus regardless of duration of disease or need for anti-diabetic agents. Code no for patients with steroid induced hyperglycemia and gestational (transient) diabetes if there is no supportive documentation of diabetes such as a HbA1c and/or treatment.

Not all patients receiving diabetic medications are considered diabetic. It is important to remember that some medications used to treat diabetes may be used to treat other conditions.

A HbA1c value ≥ 6.5 , collected within 3 months prior to surgery, is acceptable for documentation of diabetes = “yes”.

SEQ. #: 365

Long Name: RF-Diabetes-Control

Short Name: DiabCtrl

Definition: Indicate the patient's diabetes control method as presented on admission. Patients placed on a preprocedure diabetic pathway of insulin drip at admission but whose diabetes was controlled by diet or oral methods are not coded as being treated with insulin.

Choose the most aggressive therapy from the order below

- Insulin: insulin treatment (includes any combination with insulin)
- Other subcutaneous medications (e.g., GLP-1 agonist)
- Oral: treatment with oral agent (includes oral agent with or without diet treatment)
- Diet only: Treatment with diet only
- None: no treatment for diabetes
- Other: other adjunctive treatment, non-oral/insulin/diet
- Unknown

2013 ACCF/AHA Data Standards

Cannon et al. JACC Vol. 61, No. 9, 2013

Intent/Clarification:

Code diet only for patients who have had a history of diabetes that is resolved and not taking medication.

For patients who have had pancreatic transplant, code other adjunctive treatment.

SEQ. #: 370

Long Name: RF-Dyslipidemia

Short Name: Dyslip

Definition: Indicate if the patient has a history of dyslipidemia that was diagnosed and/or treated by a physician. NCEP criteria include documentation of the following:

- Total cholesterol >200 mg/dL (5.18 mmol/L); or
- LDL \geq 130 mg/dL (3.37 mmol/L);
- HDL <40 mg/dL (1.04 mmol/L) in men and <50 mg/dL (1.30 mmol/L) in women;
- Currently receiving antilipidemic treatment

Intent/Clarification:

Code "Yes" if a patient is prescribed treatment for dyslipidemia resulting in normal lab values even if anti-lipids are prescribed prophylactically, even pre-operatively, prior diagnosis with current therapy, or new diagnosis with therapy, or new diagnosis that meets the lab value definition, or if the patient is on a statin preoperatively.

Studies indicate some of the cholesterol-independent or "pleiotropic" effects of statins involve improving endothelial function, enhancing stability of atherosclerotic plaques, decreasing oxidative stress and inflammation, and inhibiting thrombogenic response.

Code "Yes" when:

- A. Documented history of dyslipidemia
- B. Lab values prior to OR entry support the diagnosis of dyslipidemia, include historic lab values
- C. Patient is on a statin medication prior to admission

Code "No" when:

- A. The patient is put on a statin after admission without the diagnosis of dyslipidemia or laboratory documentation to support the diagnosis.

SEQ. #: 375

Long Name: RF-Renal Fail-Dialysis

Short Name: Dialysis

Definition: Indicate whether the patient is currently (prior to surgery) undergoing dialysis.

Intent/Clarification:

Includes any form of peritoneal or hemodialysis the patient is receiving prior to surgery. Also, may include Continuous Veno-Venous Hemofiltration (CVVH, CVVH-D), and Continuous Renal Replacement Therapy (CRRT) as dialysis.

Code "No" for renal dialysis if ultrafiltration is the only documentation found in the record since this is for volume management.

SEQ. #: 380

Long Name: RF-Hypertension

Short Name: Hypertn

Definition: Indicate if the patient has a current diagnosis of hypertension defined by any 1 of the following:

- History of hypertension diagnosed and treated with medication, diet, and/or exercise
- Prior documentation of blood pressure >140 mm Hg systolic and/or 90 mm Hg diastolic for patients without diabetes or chronic kidney disease, or prior documentation of blood pressure >130 mm Hg systolic or 80 mm Hg diastolic on at least 2 occasions for patients with diabetes or chronic kidney disease
- Currently undergoing pharmacological therapy for treatment of hypertension

Intent/Clarification: -

SEQ. #: 385

Long Name: RF- Endocarditis

Short Name: InfEndo

Definition: Indicate whether the patient has a history of endocarditis. Endocarditis must meet the current CDC definition:

Endocarditis must meet at least 1 of the following criteria:

1. Patient has organisms cultured from valve or vegetation.
2. Patient has 2 or more of the following signs or symptoms: fever (>38°C), new or changing murmur*, embolic phenomena*, skin manifestations* (i.e., petechiae, splinter hemorrhages, painful subcutaneous nodules), congestive heart failure*, or cardiac conduction abnormality*

* With no other recognized cause and at least 1 of the following:

- 1) Organisms cultured from 2 or more blood cultures
- 2) Organisms seen on Gram's stain of valve when culture is negative or not done
- 3) Valvular vegetation seen during an invasive procedure or autopsy
- 4) Positive laboratory test on blood or urine (e.g., antigen tests for H influenzae, S pneumoniae, N meningitidis, or Group B Streptococcus)
- 5) Evidence of new vegetation seen on echocardiogram and if diagnosis is made antemortem, physician institutes appropriate antimicrobial therapy.

Choose "Yes" for patients with pre-operative endocarditis who begin antibiotics post-op. Code "Yes" for patients who are diagnosed intraoperatively.

Intent/Clarification:

Marantic Endocarditis (Nonbacterial Thrombotic Endocarditis) (Lupus) should not be coded as infectious endocarditis.

SEQ. #: 390

Long Name: RF-Infect Endocard Type

Short Name: InfEndTy

Definition: Indicate the type of endocarditis the patient has. If the patient is currently being treated for endocarditis, the disease is considered active. If no antibiotic medication (other than prophylactic medication) is being given at the time of surgery and the cultures are negative, then the infection is considered treated.

Intent/Clarification:

- **Active** - currently being treated; also include patients who were diagnosed in the OR but began treatment postop.
 - **Treated** - no antibiotic medication at time of surgery (other than prophylactic medication).
- -----

SEQ. #: 395

Long Name: RF-Infect Endocard Culture

Short Name: InfEndCult

Definition: Indicate culture results (may use cultures obtained in the OR).

Intent/Clarification:

The most common causal agents are listed; choose "other" if none of these apply or "unknown" if no culture result is available. Culture Negative, Streptococcus species, Methicillian sensitive staphylococcus aureus (MRSA), Coagulase negative staphylococcus, Enterococcus species, Gram negative species, Polymicrobial, Mycobacterium (chimera),

Fungal, Other, or Unknown. You may use cultures obtained in the OR.

SEQ. #: 400

Long Name: RF-Tobacco Use

Short Name: TobaccoUse

Definition: Indicate current (within 30 days prior to admission) or previous use of any tobacco product, including Cigarettes, Pipe, Cigars, Smokeless Cans, Other tobacco products (orbs, strips, sticks, hookah, etc.). Meaningful Use Definition

[Http://www.healthit.gov/providers-professionals/achieve-meaningful-use/core-measures/record-smoking-status](http://www.healthit.gov/providers-professionals/achieve-meaningful-use/core-measures/record-smoking-status)

Intent/Clarification:

Electronic cigarettes (Ecig) = "No;" Electronic cigarettes are not considered tobacco products.

- Current – Every Day smoker (Tobacco use within the most recent 30 days – on a daily basis)
- Current – Some Days smoker (Tobacco use within the most recent 30 days – on a less than daily basis)
- Smoker, current status unknown (Tobacco use within the most recent 30 days– frequency of use is unknown)
- Former smoker (Tobacco use prior to the most recent 30 days, without use within the most recent 30 days.)
- Smoking Status unknown (No information is available on patient's smoking status)

Example: Patient who smoked prior to admission, has been in the hospital > 2 weeks prior to surgery, and did not smoke while in the hospital is captured as "Yes, Current Every Day Smoker". The patient smoked within the 30 day window.

SEQ. #: 405

Long Name: RF-Chronic Lung Disease

Short Name: ChrLungD

Definition: Indicate whether the patient has chronic lung disease, and the severity level according to the following classification:

No

Mild: FEV1 60% to 75% of predicted, and/or on chronic inhaled or oral bronchodilator therapy.

Moderate: FEV1 50% to 59% of predicted, and/or on chronic oral/systemic steroid therapy aimed at lung disease.

Severe: FEV1 < 50% and/or Room Air pO2 < 60 or pCO2 > 50.

CLD present, severity not documented.

Unknown

Time Frame: Do not use values obtained more than 12 months prior to the date of surgery.

A history of chronic inhalation reactive disease (asbestosis, mesothelioma, black lung disease or pneumoconiosis) may qualify as chronic lung disease. Radiation induced pneumonitis or radiation fibrosis also qualifies as chronic lung disease. (if above criteria is met) A history of atelectasis is a transient condition and does not qualify.

Chronic lung disease can include patients with chronic obstructive pulmonary disease, chronic bronchitis, or emphysema. It can also include a patient who is currently being chronically treated with inhaled or oral pharmacological therapy (e.g., beta-adrenergic agonist, anti-inflammatory agent, leukotriene receptor antagonist, or steroid). Patients with asthma or seasonal allergies are not considered to have chronic lung disease.

Intent/Clarification:

Bedside spirometry can be used to quantify chronic lung disease ONLY if the study is interpreted by a pulmonologist.

SEQ. #: 410

Long Name: RF-Chronic Lung Disease - Type

Short Name: ChrLungDType

Definition: Indicate the type of chronic lung disease.

Intent/Clarification:

- **Obstructive** - Obstructive chronic lung disease is characterized by chronically poor airflow. It typically worsens over time and the main symptoms include shortness of breath, cough, and sputum production (ex. COPD; Chronic Bronchitis; Emphysema);
 - **Reactive** - Reactive lung disease is a specific type of reactive airway disease, a term used to generally describe a condition where the individual experiences asthma-like symptoms after exposure to toxins. The condition is distinctly different from asthma which is not COPD, a chronic respiratory disease where allergic reactions induce wheezing, though sometimes the terms are used interchangeably. (Ex. asbestosis and mesothelioma);
 - **Interstitial Fibrosis** - Interstitial lung disease (ILD), also known as diffuse parenchymal lung disease (DPLD), refers to a group of lung diseases affecting the interstitium (the tissue and space around the air sacs of the lungs). It concerns alveolar epithelium, pulmonary capillary endothelium, basement membrane, perivascular and perilymphatic tissues. The term ILD is used to distinguish these diseases from obstructive airways diseases; (ex. ILD, DPLD, Cystic Fibrosis)
 - **Restrictive** - Restrictive lung diseases, or restrictive ventilatory defects, are a category of extrapulmonary, pleural, or parenchymal respiratory diseases that restrict lung expansion, resulting in a decreased lung volume, an increased work of breathing, and inadequate ventilation and/or oxygenation.
 - **Other** - chronic lung disease other than previously described (ex: Amiodarone toxicity)
 - **Multiple** - Multiple types of chronic lung disease conditions are present
 - **Not documented**
- -----

SEQ. #: 415

Long Name: RF-Pulmonary Function Test

Short Name: PFT

Definition: Indicate whether pulmonary function tests were performed.

Intent/Clarification:

Pulmonary function testing is a valuable tool for evaluating the respiratory system, representing an important adjunct to the patient history, various lung imaging studies, and invasive testing such as bronchoscopy and open-lung biopsy. Insight into underlying pathophysiology can often be gained by comparing the measured values for pulmonary function tests obtained on a patient at any particular point with normative values derived from population studies. The percentage of predicted normal is used to grade the severity of the abnormality. Pulmonary function testing is used in clinical medicine for evaluating respiratory symptoms such as dyspnea and cough, for stratifying preoperative risk, and for diagnosing common diseases such as asthma and chronic obstructive pulmonary disease.

Bedside spirometry can be used to quantify chronic lung disease ONLY if the study is interpreted by a pulmonologist.

Time Frame: Do not use values obtained more than 12 months prior to the date of surgery.

SEQ. #: 420

Long Name: RF-Forced Expiratory Volume Predicted

Short Name: FEV1

Definition: Indicate the FEV1 % predicted from the most recent pulmonary function test prior to procedure.

Choose the highest value reported for % predicted, whether or not a bronchodilator was used.

Intent/Clarification:

FEV₁ is the maximal amount of air forcefully exhaled in one second. It is then converted to a percentage of normal. For example, the FEV₁ may be 80% of predicted based on height, weight, and race. FEV₁ is a marker for the degree of obstruction. In normal persons, the FEV₁ accounts for the greatest part of the exhaled volume from a spirometric maneuver and reflects mechanical properties of the large and the medium-sized airways.

- FEV1 > 75% of predicted = Normal
- FEV1 60% to 75% of predicted = Mild obstruction
- FEV1 50% to 59% of predicted = Moderate obstruction
- FEV1 < 50% of predicted = Severe obstruction

SEQ. #: 425

Long Name: DLCO Test Performed

Short Name: DLCO

Definition: Indicate whether a lung diffusion test (DLCO) was performed.

Intent/Clarification:

The diffusing capacity (DLCO) is a test of the integrity of the alveolar-capillary surface area for gas transfer.

SEQ. #: 430

Long Name: DLCO Predicted **Short Name:** DLCOPred

Definition: Indicate the % predicted DLCO value obtained for the patient. Choose the value that represents the highest **lowest** % predicted whether or not it is the simple DLCO or the DLCO/VA.

Intent/Clarification:

The **lowest** value for DLCO uncorrected should be captured.

SEQ. #: 435

Long Name: RF-Arterial Blood Gas

Short Name: ABG

Definition: Indicate whether a room-air arterial blood gas was performed prior to surgery. Answer no if the only available arterial blood gasses were drawn while patient was receiving supplemental oxygen.

Intent/Clarification:

Arterial blood gasses may be drawn in patients with suspected lung disease or sometimes during cardiac catheterization. Do not use ABGs drawn after initiation of anesthetic management. They may not accurately reflect the patient's true baseline due to preop sedation, anxiety, pain and other factors.

Answer "No" if the only available arterial blood gasses were drawn while patient was receiving supplemental oxygen.

SEQ. #: 440

Long Name: RF-Carbon Dioxide Level

Short Name: PCO2

Definition: Indicate PCO2 on most recent room air blood gas prior to procedure.

Intent/Clarification:

Higher levels (CO₂ retention) may indicate hypoventilation and low levels are consistent with hyperventilation.

The normal range is 35-45 mmHg.

SEQ. #: 445

Long Name: RF-Oxygen Level

Short Name: PO2

Definition: Indicate PO2 result on most recent room air arterial blood gas prior to procedure.

Intent/Clarification:

The partial pressure of oxygen that is dissolved in arterial blood is known as PO₂. In persons over 60 years of age, the normal is lower.

Normal values 80-100mm Hg.

SEQ. #: 450

Long Name: RF-Home Oxygen

Short Name: HmO2

Definition: Indicate whether supplemental oxygen at home is prescribed and used.

Intent/Clarification:

Choices include the following:

- Yes, PRN
- Yes, Oxygen dependent
- No
- Unknown

Code "No" for patients who are using home O₂ on a prn basis but have not used > 1 month,

Code "Unknown" if there is no indication of when home O₂ was last used.

SEQ. #: 455

Long Name: RF-Inhaled Medication or Oral Bronchodilator Therapy

Short Name: BDTx

Definition: Indicate whether oral and/or inhaled bronchodilator or inhaled (not oral or IV) steroid medications were in use by the patient routinely prior to this procedure.

Intent/Clarification:

Capture patients with prescribed home bronchodilator therapy prior to admission. Capture only routine use. Do not capture for patients using bronchodilators to treat asthma.

A bronchodilator is a substance that dilates the bronchi and bronchioles, decreasing airway resistance and thereby facilitating airflow. They are most useful in obstructive lung diseases, of which asthma and chronic obstructive pulmonary disease are the most

common conditions. Bronchodilators are either short-acting or long-acting. Short-acting medications provide quick or "rescue" relief from acute bronchoconstriction. Long-acting bronchodilators help to control and prevent symptoms.

Code "Unknown" when there is conflicting information in the medical record and/or with the patient/family.

SEQ. #: 460

Long Name: RF-Sleep Apnea

Short Name: SlpApn

Definition: Indicate whether patient has a diagnosis of sleep apnea (may be described as obstructive sleep apnea or OSA).

Intent/Clarification:

Sleep apnea is a potentially serious sleep disorder in which breathing repeatedly stops and starts during sleep. Sleep apnea occurs in two main types: Obstructive Sleep Apnea, the more common form that occurs when throat muscles relax, and Central Sleep Apnea, which occurs when the brain doesn't send proper signals to the muscles that control breathing. Additionally, some people have complex sleep apnea, which is a combination of both. Sleep apnea has been associated with sudden death.

- Capture patients with prescribed home therapy despite frequency of use.
- Sleep apnea must be diagnosed by a physician/NP/PA. **Do not capture suspected sleep apnea or that reported by family members as sleep apnea.**
- CPAP or BiPAP therapy is no longer a requirement to code "yes" for sleep apnea.
- Code "No" to sleep apnea if sleep apnea has been surgically corrected.

Code "Yes" if sleep apnea is diagnosed using a diagnostic tool but is not treated.

SEQ. #: 465

Long Name: RF-Pneumonia

Short Name: Pneumonia

Definition: Indicate whether patient has a recent (within 30 days) or remote (more than 30 days) history of pneumonia.

Intent/Clarification:

Pneumonia is an infection of one or both lungs caused by bacteria, viruses, fungi, chemicals, or aspiration. It can be community acquired or acquired in a health care setting. Typical symptoms associated with pneumonia include cough, chest pain, fever, and difficulty in breathing. Diagnostic tools include x-rays and examination of the sputum. Treatment depends on the cause of pneumonia; bacterial pneumonia is treated with antibiotics.

Code as:

- Recent- pneumonia diagnosis within 30 days of procedure or
- Remote - pneumonia diagnosis more than 30 days prior to the procedure.

- No - meaning no history of pneumonia
- Unknown

There must be documentation of pneumonia to code "Yes". "**Possible** pneumonia" with antibiotic treatment should be coded "Unknown".

Pneumonitis, inflammation of the lung tissue, without infection is not considered pneumonia and should be coded as "no".

SEQ. #: 470

Long Name: RF-Illicit Drug Use **Short Name:** IVDrugAb

Definition: Indicate whether documented history of use of illicit drugs, such as heroin, marijuana, cocaine, or methamphetamine, or abuse of a controlled substance.

Do not include rare historical use. Do not include prescribed medicinal marijuana.

Treatment with methadone is not considered illicit drug use.

Intent/Clarification:

Capture patients with habitual use of illicit drugs. Include abuse of street and prescription medications. Illicit drug use is associated with numerous health and social problems, and age-related physiological, psychological, and social changes that could impact recovery from surgery.

- **Recent** - Within 30 days of procedure
 - **Remote** - More than 30 days prior to procedure
 - **No** – No illicit drug use
 - **Unknown** - patient/family unable to provide history
- -----

SEQ. #: 475

Long Name: RF-Depression

Short Name: Depression

Definition: Indicate whether there is a current or previous history of depression or documentation of a depressed mood or affect.

Intent/Clarification:

People with heart disease are more likely to suffer from depression than otherwise healthy people. Angina and heart attacks are closely linked with depression. Recovery following heart surgery may be negatively impacted by the presence of preoperative depression.

A clinical diagnosis of depression/treatment is not required to code "Yes". Patient/family stating depressed mood or affect is sufficient.

Do not code "Yes" if a patient taking antidepressant medications for something other than depression (examples: radiculopathy, smoking cessation, fibromyalgia, sleep disorders, hormonal imbalances).

Include patients who are reporting symptoms of depression; treated or not treated with medications or therapy.

Bipolar disorders are considered depression.

Code "Unknown" when there is conflicting information in the medical record and/or with the patient/family.

SEQ. #: 480

Long Name: RF-Alcohol Use

Short Name: Alcohol

Definition: Specify alcohol consumption history.

Intent/Clarification: 490

Code current alcohol use (within 30 days of surgery):

- ≤ 1 drink per week (rare or occasional drink) one beer, one glass of wine or one shot
- 2-7 drinks per week (Social)
- ≥ 8 drinks per week (Heavy drinker)
- None (Non-drinker)
- Unknown- patient/family unable to provide history

Alcohol abuse is not necessarily a quantity of alcohol but implies interference with home, work and life functioning. Documentation the patient is an alcoholic at the time of admission should be coded ≥ 8 drinks per week.

SEQ. #: 485

Long Name: RF-Liver Disease

Short Name: LiverDis

Definition: Indicate whether the patient has a history of hepatitis B, hepatitis C, cirrhosis, portal hypertension, esophageal varices, chronic alcohol abuse or congestive hepatopathy. Exclude NASH in the absence of cirrhosis.

Intent/Clarification:

LFTs or a MELD score alone **cannot** be used to code "Yes" to liver disease since other conditions impact these lab values. Liver fibrosis with recurrent ascites, supported by the MELD can be coded as liver disease.

The following are not coded as liver disease:

- ~~Patients who have had liver transplant without residual anatomic or systemic issues or associated MELD scores.~~
- Hepatitis A
- Gilberts syndrome
- Fatty liver
- Liver Cancer

FAQ September 2017: Patient had a liver transplant 11 years ago and is here for convergent A-Fib ablation. His MELD score this admission was 10.69, bilirubin was 1.8, and there was no GI/Hepatology consult.

How should I code liver disease? I wouldn't be able to show that he has history of liver transplant if I key no.

Answer: Capture yes to liver disease and yes to liver transplant. It is important to capture the history of liver disease.

SEQ. #: 486

Long Name: RF-Liver Disease - Child Pugh Class

Short Name: LiverChildPugh

Definition: Indicate the Child Pugh Class, if known.

Intent/Clarification:

Documentation includes the compilation of the MELD score, the clinical diagnosis and the controllability of ascites.

- Child-Pugh A
- Child-Pugh B
- Child-Pugh C
- Unknown

FAQ August 2017: Can I calculate the Child-Pugh classification?

Answer: No, it is the responsibility of the surgeon/physician to calculate the Child-Pugh classification and document the score in the medical record. If not documented, code unknown.

SEQ. #: 487

Long Name: RF-Liver Disease - Listed for Liver Transplant

Short Name: LiverTransList

Definition: Indicate whether the patient is listed for liver transplant.

Intent/Clarification: -

SEQ. #: 488

Long Name: RF-Liver Disease - Status Post Liver Transplant

Short Name: LiverStatusPost

Definition: Indicate whether the patient has received a liver transplant prior to this operation.

Intent/Clarification: -

FAQ September 2017: Patient had a liver transplant 11 years ago and is here for convergent A-Fib ablation. His MELD score this admission was 10.69, bilirubin was 1.8, and there was no GI/Hepatology consult.

How should I code liver disease? I wouldn't be able to show that he has history of liver transplant if I key no.

Answer: Capture yes to liver disease and yes to liver transplant. It is important to capture the history of liver disease.

SEQ. #: 490

Long Name: RF-Immunocompromise

Short Name: ImmSupp

Definition: Indicate whether immunocompromise is present due to immunosuppressive medication therapy within 30 days preceding the operative procedure or existing medical condition. This includes, but is not limited to systemic steroid therapy, anti-rejection medications and chemotherapy. This does not include topical steroid applications, one-time systemic therapy, inhaled steroid therapy or pre-procedure protocol.

Intent/Clarification:

Include patients being treated with IVIG. Patients who have had splenectomy are considered immunocompromised. Examples of conditions causing immunocompromise include Hypogammaglobulinemia and HIV infection.

SEQ. #: 495

Long Name: RF-Mediastinal Radiation

Short Name: MediastRad

Definition: Indicate whether patient has a history of radiation therapy to the mediastinum or chest.

Intent/Clarification:

Chest wall or mediastinal radiation can cause damage to blood vessels, heart valves and lung tissue. Scar tissue caused by radiation therapy can lead to increased bleeding, may make harvesting the internal mammary artery difficult and may interfere with sternal healing.

Include radiation to the "mantel/chest" area only – this includes breast cancer with radiation.

SEQ. #: 500

Long Name: RF-Cancer Within 5 Years

Short Name: Cancer

Definition: Indicate whether the patient has a history of cancer diagnosed within 5 years of procedure. Do not capture low grade skin cancers such as basal cell or squamous cell carcinoma.

Intent/Clarification:

Capture cancers that have or will require surgical intervention, chemotherapy and or radiation therapy. If the date of diagnosis is not known, then the date of the last treatment may be used to determine the 5 year interval.

SEQ. #: 505

Long Name: RF-Peripheral Arterial Disease

Short Name: PVD

Definition: Indicate whether the patient has a history of peripheral arterial disease (includes upper and lower extremity, renal, mesenteric, and abdominal aortic systems).

This can include:

1. Claudication, either with exertion or at rest,
2. Amputation for arterial vascular insufficiency,
3. Vascular reconstruction, bypass surgery, or percutaneous intervention to the extremities (excluding dialysis fistulas and vein stripping),
4. Documented abdominal aortic aneurysm with or without repair,
5. Positive noninvasive test (e.g., ankle brachial index \leq 0.9, ultrasound, magnetic resonance or computed tomography imaging of $>$ 50% diameter stenosis in any peripheral artery, i.e., renal, subclavian, femoral, iliac) or angiographic imaging

Peripheral arterial disease excludes disease in the carotid, cerebrovascular arteries or thoracic aorta.

PVD does not include DVT.

Intent/Clarification:

PAD is sometimes called PVD, code only arterial disease. PAD includes subclavian artery stenosis.

SEQ. #: 510

Long Name: RF-Thoracic Aorta Disease

Short Name: ThAoDisease

Definition: Indicate whether the patient has a history of disease of the thoracic or thoracoabdominal aorta.

Abdominal aortic disease without thoracic involvement is captured in peripheral artery disease.

Intent/Clarification:

Code "Yes" to aortic aneurysms, aortic dissection/rupture. Fusiform ascending thoracic aneurysm is more likely to dissect when the aortic cross clamp is applied and should be coded as thoracic aorta disease.

Code "No" to blunt trauma or infection.

This field is not intended to capture calcification of the aorta.

SEQ. #: 515

Long Name: RF-Syncope

Short Name: Syncope

Definition: Indicate whether the patient had a sudden loss of consciousness with loss of postural tone, not related to anesthesia, with spontaneous recovery and believed to be related to cardiac condition. Capture events occurring within the past one year as reported by patient or observer. Patient may experience syncope when supine.

Intent/Clarification:

Cardiac conditions including dysrhythmias, such as ventricular tachycardia or ventricular fibrillation, and aortic stenosis can cause syncope.

Near syncope should be coded as "no".

Cardiac arrest with resuscitation is **not** syncope.

SEQ. #: 520

Long Name: RF-Unresponsive Neurologic State

Short Name: UnrespStat

Definition: Indicate whether the patient has a history of non-medically induced, unresponsive state within 24 hours of the time of surgery. Patient experienced complete mental unresponsiveness and no evidence of psychological or physiologically appropriate responses to stimulation, includes patients who experience sudden cardiac death.

Intent/Clarification:

The intent is to identify those patients whose postoperative neurologic state may not be a result of the surgery but rather patient's unknown preoperative neurologic status.

Code "Yes" if the patient never regained consciousness prior to surgery.

Temporary loss of consciousness that resolved after cardiac arrest should not be coded as yes.

SEQ. #: 521

Long Name: RF-Chest Wall Deformity

Short Name: ChestWallDef

Definition: Indicate whether the patient has a chest wall deformity.

Intent/Clarification:

A deformity is thought to be caused by excessive growth of the costal cartilages (ribs), although the reason for this is unknown. This overgrowth causes the ribs and cartilages to buckle and pushes the sternum either inwards or outwards.

Pectus excavatum results in a sunken sternum sometimes called funnel chest and usually involves the lower half of the sternum.

In pectus carinatum the sternum protrudes in a convex shape and is sometimes called pigeon chest.

Chest wall deformity should be coded for the patient with an existing deformity or one that has been previously repaired.

SEQ. #: 525

Long Name: RF-Cerebrovascular Dis

Short Name: CVD

Definition: Indicate whether the patient has a current or previous history of any of the following:

A. Stroke: Stroke is an acute episode of focal or global neurological dysfunction caused by brain, spinal cord, or retinal vascular injury as a result of hemorrhage or infarction, where the neurological dysfunction lasts for greater than 24 hours.

B. TIA: is defined as a transient episode of focal neurological dysfunction caused by brain, spinal cord, or retinal ischemia, without acute infarction, where the neurological dysfunction resolves within 24 hours.

C. Noninvasive or invasive arterial imaging test demonstrating $\geq 50\%$ stenosis of any of the major extracranial or intracranial vessels of the brain

D. Previous cervical or cerebral artery revascularization surgery or percutaneous intervention

This does not include chronic (nonvascular) neurological diseases or other acute neurological insults such as metabolic and anoxic ischemic encephalopathy.

Intent/Clarification:

A positive CT scan, even in the patient with no symptoms, should be coded as cerebral vascular disease. A CT scan following surgery with evidence of old infarct should be coded no.

Subdural hematoma is not cerebral vascular disease.

SEQ. #: 530

Long Name: RF-Prior CVA

Short Name: CVA

Definition: Indicate whether the patient has a history of stroke. Stroke is an acute episode of focal or global neurological dysfunction caused by brain, spinal cord, or retinal vascular injury as a result of hemorrhage or infarction, where the neurological dysfunction lasts for greater than 24 hours.

Intent/Clarification:

Include any confirmed neurological deficit of abrupt onset caused by a disturbance in

cerebral blood supply that did not resolve within 24 hours of the event. The physical deficit can be in the form of extremity weakness, facial asymmetry, language (speech and/or cognitive thinking) impairment. The intent is to differentiate between neurological events that resolve within 24 hours and those that don't.

Code "yes" to prior CVA if the patient has no history of stroke and no symptoms but imaging study results show an infarct (old/chronic or new).

SEQ. #: 535

Long Name: RF-Prior CVA-When

Short Name: CVAWhen

Definition: Indicate when the CVA events occurred. Those events occurring within 30 days prior to the surgical procedure are considered recent, while all others are considered remote.

Intent/Clarification:

≤ 30 days is recent

> 30 days is remote

SEQ. #: 540

Long Name: RF-CVD TIA

Short Name: CVDTIA

Definition: Indicate whether the patient has a history of a Transient Ischemic Attack (TIA). Transient ischemic attack (TIA) is defined as a transient episode of focal neurological dysfunction caused by brain, spinal cord, or retinal ischemia, without acute infarction, where the neurological dysfunction resolves within 24 hours.

Intent/Clarification:

Choices are:

- Yes
- No
- Unknown

"Unknown" should be selected if any neurologic dysfunction occurred or was suspected, was resolved in 24 hours, and could not be confirmed or if patient/family unable to provide history.

SEQ. #: 545

Long Name: RF-CVD Carotid Stenosis

Short Name: CVDCarSten

Definition: Indicate which carotid artery was determined from any diagnostic test to be ≥ 50% stenotic.

Intent/Clarification:

Code what is found at the time of surgery,even if a prior stent is in place.

Choices are:

- None
- Right
- Left
- Both

If the results are reported in a range, such as “40-50%”, choose the highest level in the range.

SEQ. #: 550

Long Name: RF-CVD Carotid Stenosis - Right

Short Name: CVDStenRt

Definition: Indicate the severity of stenosis reported on the right carotid artery.

Intent/Clarification:

Indicate % stenosis:

50 - 79% or “moderate”

80 - 99% or “critical”, “severe”, or “subtotal”.

100% or “total”

Not documented

If the results are reported in a range, such as “40-50%”, choose the highest level in the range.

SEQ. #: 555

Long Name: RF-CVD Carotid Stenosis - Left **Short Name:** CVDStenLft

Definition: Indicate the severity of stenosis reported on the left carotid artery.

Intent/Clarification:

Indicate % stenosis:

50 - 79% = “moderate”

80 - 99% = “critical”, “severe”, or “subtotal”.

100% = “total”

Not documented

If the results are reported in a range, such as “40-50%”, choose the highest level in the range.

SEQ. #: 560

Long Name: RF-CVD Prior Carotid Surgery

Short Name: CVDPCarSurg

Definition: Indicate whether the patient has a history of previous carotid artery surgery and/or stenting.

Intent/Clarification:

Carotid endarterectomy is a surgical procedure during which a surgeon removes atherosclerotic plaque or other material obstructing the flow of blood from the artery. This procedure eliminates a substance called plaque from the artery and can restore blood flow.

Carotid artery stenting is a procedure in which a slender, metal-mesh tube, called a stent, is inserted and expands inside the carotid artery to increase blood flow in areas blocked by plaque.

Also includes internal carotid artery aneurysm coils.

General Information for Labs:

Use results closest to surgery, prior to anesthesia provider initiating care. STS recommends values within 30 days, unless otherwise stated.

Capture lab values if available. Not all patients will have, or need to have, all of the following labs drawn.

Do not use labs drawn after IV fluids are hung in holding area or OR. Include POC (point of care) results.

SEQ. #: 565

Long Name: RF-Last WBC Count

Short Name: WBC

Definition: Indicate the pre-operative White Blood Cell (WBC) count closest to the date and time prior to surgery but prior to anesthetic management (induction area or operating room).

Intent/Clarification:

White Blood Cells (leukocytes) are part of the body's immune defense and are often elevated in the presence of infection. The hospital laboratory report should be accessed first when coding this variable. If this is unavailable, then additional source documents may be referenced for lab results.

SEQ. #: 570

Long Name: RF-Hemoglobin

Short Name: RFHemoglobin

Definition: Indicate the pre-operative Hemoglobin level at the date and time closest to surgery but prior to anesthetic management (induction area or operating room). Capture only measured hemoglobin levels, not calculated values.

Intent/Clarification:

The hemoglobin (Hgb) test may be used to screen for, diagnose, or monitor a number of conditions and diseases that affect red blood cells (RBCs) and/or the amount of hemoglobin in blood. The hospital laboratory report should be accessed first when coding this variable. If this is unavailable, then additional source documents may be referenced for lab results.

SEQ. #: 575

Long Name: RF-Last Hematocrit

Short Name: Hct

Definition: Indicate the pre-operative Hematocrit level at the date and time closest to surgery but prior to anesthetic management (induction area or operating room). Capture only measured hematocrit levels, not calculated values.

Intent/Clarification:

Hematocrit (Hct) is the proportion of red cells in the blood. The hospital laboratory report should be accessed first when coding this variable. If this is unavailable, then additional source documents may be referenced for lab results.

SEQ. #: 580

Long Name: RF-Platelets

Short Name: Platelets

Definition: Indicate the platelet count closest to the date and time prior to surgery but prior to anesthetic management (induction area or operating room).

Intent/Clarification:

Platelets (plt) are a blood component instrumental in clot formation. The hospital laboratory report should be accessed first when coding this variable. If this is unavailable, then additional source documents may be referenced for lab results.

SEQ. #: 585

Long Name: RF-Last Creat Level

Short Name: CreatLst

Definition: Indicate the creatinine level closest to the date and time prior surgery but prior to anesthetic management (induction area or operating room).

A creatinine level should be collected on all patients, even if they have no prior history of renal disease. A creatinine value is a high predictor of a patient's outcome and is used in the predicted risk models.

Intent/Clarification:

Creatinine (Cr) is a chemical waste molecule excreted by the kidneys that is generated

from muscle metabolism. If the kidneys become impaired for any reason, the creatinine level in the blood will rise due to poor clearance by the kidneys. Abnormally high levels of creatinine thus warn of possible malfunction or failure of the kidneys.

SEQ. #: 590

Long Name: RF-Total Albumin

Short Name: TotAlbumin

Definition: Indicate the total albumin closest to the date and time prior to surgery but prior to anesthetic management (induction area or operating room).

Intent/Clarification:

Albumin (alb), produced only in the liver, is the major plasma protein that circulates in the bloodstream. Albumin is essential for maintaining the oncotic pressure in the vascular system. A decrease in oncotic pressure due to a low albumin level allows fluid to leak out from the interstitial spaces into the peritoneal cavity, producing ascites. Albumin is also very important in the transportation of many substances such as drugs, lipids, hormones, and toxins that are bound to albumin in the bloodstream. A low serum albumin indicates poor liver function. Decreased serum albumin levels are not seen in acute liver failure because it takes several weeks of impaired albumin production before the serum albumin level drops. The most common reason for a low albumin is chronic liver failure caused by cirrhosis. The serum albumin concentration is usually normal in chronic liver disease until cirrhosis and significant liver damage has occurred.

You can capture results up to 6 weeks prior to surgery provided there is no known acute liver disease process.

SEQ. #: 595

Long Name: RF-Total Bilirubin

Short Name: TotBlrbn

Definition: Indicate the total Bilirubin closest to the date and time prior to surgery but prior to anesthetic management (induction area or operating room).

Intent/Clarification:

Bilirubin (Tbili) testing checks for levels of bilirubin, an orange-yellow pigment, in blood. Bilirubin is a natural byproduct that results from the normal breakdown of red blood cells. As a normal process, bilirubin is carried in the blood and passes through the liver. Too much bilirubin may indicate liver damage or disease.

You can capture results up to 6 weeks prior to surgery provided there is no known acute liver disease process.

SEQ. #: 600

Long Name: RF-Last A1c Level

Short Name: A1cLvl

Definition: Indicate the pre-operative HbA1c level closest to the date and time prior surgery but prior to anesthetic management (induction area or operating room).

Intent/Clarification:

Glycosylated hemoglobin, HbA1c, is a form of hemoglobin used primarily to identify the average plasma glucose concentration over prolonged periods of time. It is formed in a non-enzymatic glycation pathway by hemoglobin's exposure to plasma glucose. Normal levels of glucose produce a normal amount of glycosylated hemoglobin. As the average amount of plasma glucose increases, the fraction of glycosylated hemoglobin increases in a predictable way. This serves as a marker for average blood glucose levels over the previous months prior to the measurement. The HbA1c level is proportional to average blood glucose concentration over the previous four weeks to three months.

The 2010 American Diabetes Association Standards of Medical Care in Diabetes added the A1c \geq 6.5% as a criterion for the diagnosis of diabetes.

This lab must be drawn preoperatively since cardiopulmonary bypass, fluid shifts and transfusions can alter results and not accurately reflect the prior glucose levels.

You can capture results up to 3 months prior to surgery.

SEQ. #: 605

Long Name: RF-HIT Antibodies

Short Name: HITAnti

Definition: Indicate whether Heparin Induced Thrombocytopenia (HIT) is confirmed by antibody testing.

Intent/Clarification:

Heparin induced thrombocytopenia (HIT) can be defined as any clinical event best explained by platelet factor 4 (PF4)/ heparin-reactive antibodies ('HIT antibodies') in a patient who is, or has recently received heparin. Thrombocytopenia is the most common 'event' in HIT and occurs in at least 90% of patients, depending upon the definition of thrombocytopenia. A high proportion of patients with HIT develop thrombosis.

Alternative, non-heparin, anticoagulant therapy reduces the risk of subsequent thrombosis. The SRA (serotonin release assay) test is the most definitive HIT test. The timeframe is any time prior to surgery. <http://emedicine.medscape.com/article/1357846-overview>

Choices are:

- Yes - Positive antibody testing (test was performed, HIT positive)
 - No - Negative antibody testing (test was performed, HIT negative)
 - NA - antibody testing not performed
- -----

SEQ. #: 610

Long Name: RF-INR

Short Name: INR

Definition: Indicate the International Normalized Ratio (INR) closest to the date and time prior to surgery but prior to anesthetic management (induction area or operating room).

Intent/Clarification:

INR is the standard unit used to report the result of a prothrombin (PT) test. An individual whose blood clots normally and who is not on anticoagulation should have an INR of approximately 1. The higher the INR, the longer it takes blood to clot. As the INR increases above a given level, the risk of bleeding and bleeding-related events increases. As the INR decreases below a given level, the risk of clotting events increases.

SEQ. #: 615

Long Name: RF-MELD Score

Short Name: MELDScr

Definition: MELD score value calculated by software to indicate severity of liver disease.

Intent/Clarification:

MELD is a validated liver disease severity scoring system that uses laboratory values for serum bilirubin, serum creatinine and the INR to predict survival. In patients with chronic liver disease, an increasing MELD score is associated with increasing risk of death.

MELD is not used to confirm liver disease, rather as a severity measure for patients with known liver disease.

≤ 15 predictive of 95% survival at 3 months

~ 30 predictive of 65% survival at 3 months

≥ 40 predictive of 10-15% survival at 3 months

MELD = 3.8[Ln serum bilirubin (mg/dL)] + 11.2[Ln INR] + 9.6[Ln serum creatinine (mg/dL)] + 6.4. Laboratory values of INR, total bilirubin and serum creatinine that are <1.0 are set to 1.0. In addition, serum creatinine levels >4.0 mg/dL are capped at 4.0 mg/dL, and patients on dialysis receive an assigned serum creatinine value of 4.0 mg/dL.

Reference: <http://www.mayoclinic.org/medical-professionals/model-end-stage-liver-disease>

SEQ. #: 620

Long Name: RF-BNP

Short Name: BNP

Definition: Indicate the BNP value.

Intent/Clarification:

Brain natriuretic peptide (BNP), now known as B-type natriuretic peptide or Ventricular Natriuretic Peptide (still BNP), is a 32-amino acid polypeptide secreted by the ventricles

of the heart in response to excessive stretching of heart muscle cells (cardiomyocytes). The physiologic actions of BNP are similar to those of ANP and include decrease in systemic vascular resistance and central venous pressure as well as an increase in natriuresis. Thus, the net effect of BNP and ANP is a decrease in blood volume, which lowers systemic blood pressure and afterload, yielding an increase in cardiac output, partly due to a higher ejection fraction.

SEQ. #: 645

Long Name: RF-Five Meter Walk Test Done

Short Name: FiveMWalkTest

Definition: Indicate whether the five meter walk test was done.

Intent/Clarification:

Frailty is a risk factor for surgery that has been difficult to quantify. This test quantifies frailty prior to surgery in ambulatory patients. **Prolonged times strongly correlate with increased risk and this risk factor will be assessed for possible inclusion in risk model updates being done in 2015.**

Instructions:

1. Accompany the patient to the designated area, which should be well-lit, unobstructed, and contain clearly indicated markings at 0 and 5 meters
2. Position the patient with his/her feet behind and just touching the 0-meter start line
3. Instruct the patient to "walk at your comfortable pace" until a few steps past the 5-meter mark (the patient should not start to slow down before the 5-meter mark)
4. Begin each trial on the word "Go"
5. Start the timer with the first footfall after the 0-meter line
6. Stop the timer with the first footfall after the 5-meter line
7. Repeat 3 times, allowing sufficient time for recuperation between trials. (If patient is unable to repeat x3, enter 1 or 2 times)

Note: Patient may use a walking aid (cane, walker). If the patient is receiving an IV drip, he/she should perform the test without the IV only if it can be interrupted temporarily without any potential risk to the patient, if not, then the patient may perform the test pushing the IV pole. If the time taken to walk 5 meters averages > 6 seconds, the patient is considered frail.

Choices are:

- Yes
- No
- Non-ambulatory patient (physically or medically unable to perform the test)

Reference: *Gait Speed as an Incremental Predictor of Mortality and Major Morbidity in Elderly...* Afilalo et al. J Am Coll Cardiol.2010; 56: 1668-1676

SEQ. #: 650

Long Name: RF-Five Meter Walk Time 1

Short Name: FiveMWalk1

Definition: Indicate the time in seconds it takes the patient to walk 5 meters for the first of three tests.

Intent/Clarification: -

SEQ. #: 655

Long Name: RF-Five Meter Walk Time 2

Short Name: FiveMWalk2

Definition: Indicate the time in seconds it takes the patient to walk 5 meters for the second of three tests.

Intent/Clarification: -

SEQ. #: 660

Long Name: RF-Five Meter Walk Time 3

Short Name: FiveMWalk3

Definition: Indicate the time in seconds it takes the patient to walk 5 meters for the third of three tests.

Intent/Clarification: -

SEQ. #: 661

Long Name: RF - Six Minute Walk Test Done

Short Name: SixMWalkDone

Definition: Indicate whether a six-minute walk test was done.

Intent/Clarification:

The 6MWT is a practical simple test that requires a 100-ft hallway but no exercise equipment or advanced training for technicians. Walking is an activity performed daily by all but the most severely impaired patients. This test measures the distance that a patient can quickly walk on a flat, hard surface in a period of 6 minutes (the 6MWD). It evaluates the global and integrated responses of all the systems involved during exercise, including the pulmonary and cardiovascular systems, systemic circulation, peripheral circulation, blood, neuromuscular units, and muscle metabolism.

- Yes
 - No
-
-

SEQ. #: 662

Long Name: RF - Six Minute Walk Test Distance

Short Name: SixMWalkDist

Definition: Indicate the distance in feet the patient walked during the six-minute walk test.

Intent/Clarification: -

Previous Cardiac Interventions

SEQ. #: 665

Long Name: Prev Cardiac Intervent

Short Name: PrCVInt

Definition: Indicate whether the patient has undergone any previous cardiovascular intervention, either surgical or non-surgical, which may include those done during the current admission.

Intent/Clarification:

A patient who had previous invasive cardiac procedures (PCI or surgery) will have increased risk due to a variety of factors; such as repeated exposure to heparin potentiating incidence of heparin antibodies, heparin resistance or surgical adhesions. This is intended to capture surgical and/or interventional procedures, not diagnostic ones like TEE or cath.

SEQ. #: 670

Long Name: Prev CAB

Short Name: PrCAB

Definition: Indicate whether the patient had a previous Coronary Bypass Graft prior to the current admission.

Intent/Clarification:

This applies only to surgical approach to revascularization. Angioplasty or other catheter based coronary artery occlusion treatment does not apply.

SEQ. #: 675

Long Name: Prev Valve

Short Name: PrValve

Definition: Indicate whether the patient had a previous surgical replacement and/or surgical repair of a cardiac valve. This may also include percutaneous valve procedures.

Intent/Clarification:

This may include percutaneous valve procedures such as percutaneous valvotomy or valvuloplasty, as well as surgical or transcatheter valve repair or replacement. Capture

all procedures that apply.
These do not have to be in chronological order.

SEQ. #: 695

Long Name: Prev Valve Procedure 1

Short Name: PrValveProc1

Definition: Indicate the first previous valve procedure.

Intent/Clarification: Indicate which specific valve procedure was performed:

No additional valve procedure(s)

Aortic valve balloon valvotomy/valvuloplasty

Aortic valve repair, surgical

Aortic valve replacement, surgical

Aortic valve replacement, transcatheter

Mitral valve balloon valvotomy/valvuloplasty

Mitral valve commissurotomy, surgical

Mitral valve repair, percutaneous

Mitral valve repair, surgical

Mitral valve replacement, surgical

Mitral valve replacement, transcatheter

Tricuspid valve balloon valvotomy/valvuloplasty

Tricuspid valve repair, percutaneous

Tricuspid valve repair, surgical

Tricuspid valve replacement, surgical

Tricuspid valve replacement, transcatheter

Tricuspid valvectomy

Pulmonary valve balloon valvotomy/valvuloplasty

Pulmonary valve repair, surgical

Pulmonary valve replacement, surgical

Pulmonary valve replacement, transcatheter

Pulmonary valvectomy

Other valve procedure

SEQ. #: 700

Long Name: Prev Valve Procedure 2

Short Name: PrValveProc2

Definition: Indicate the second previous valve procedure or select "No additional valve procedures"

Intent/Clarification:

If a second procedure was done, please select from the list above or select:

No Additional Valve Procedure(s) - Software will grey out any additional selections.

SEQ. #: 705

Long Name: Prev Valve Procedure 3

Short Name: PrValveProc3

Definition: Indicate the third previous valve procedure or select "No additional valve procedures"

Intent/Clarification:

If a third procedure was done, please select from the list above or select:

No Additional Valve Procedure(s) - Software will grey out any additional selections.

SEQ. #: 710

Long Name: Prev Valve Procedure 4

Short Name: PrValveProc4

Definition: Indicate the fourth previous valve procedure or select "No additional valve procedures"

Intent/Clarification:

If a fourth procedure was done, please select from the list above or select:

No Additional Valve Procedure(s) - Software will grey out any additional selections.

SEQ. #: 715

Long Name: Prev Valve Procedure 5

Short Name: PrValveProc5

Definition: Indicate the fifth previous valve procedure or select "No additional valve procedures"

Intent/Clarification:

If a fifth procedure was done, please select from the list above or select:

No Additional Valve Procedure(s) - Software will grey out any additional selections.

SEQ. #: 775

Long Name: Previous PCI

Short Name: POCPCI

Definition: Indicate whether a previous Percutaneous Coronary Intervention (PCI) was performed any time prior to this surgical procedure.

Percutaneous coronary intervention (PCI) is the placement of an angioplasty guide wire, balloon, or other device (e.g. stent, atherectomy, brachytherapy, or thrombectomy catheter) into a native coronary artery or coronary artery bypass graft for the purpose of mechanical coronary revascularization.

Intent/Clarification:

An **attempted**, even if unsuccessful, PCI should be coded as a Previous CV intervention-PCI. This is in an effort to harmonize with ACC-NCDR.

SEQ. #: 780

Long Name: Previous PCI-Within This Episode of Care

Short Name: POCPCIWhen

Definition: Indicate whether the previous Percutaneous Cardiac Intervention (PCI) was performed within this episode of care. Episode of care is defined as continuous inpatient hospitalization which includes transfer from one acute care hospital to another.

Intent/Clarification:

This field is intended to capture PCIs done during the same episode of care prior to the surgical procedure. Include patients who were transferred for surgery from another facility following PCI.

Do not code PCIs done after the surgical procedure here.

Do not code as the same episode of care if the patient is discharged home between interventions. Choices are:

- Yes, at this facility
 - Yes, at some other acute care facility
 - No
-
-

NOTE THAT SEQUENCE NUMBER 785 IS A CHILD TO SEQUENCE NUMBER 780.

SEQ. #: 785

Long Name: Previous PCI-Indication For Surgery

Short Name: POCPCIndSurg

Definition: Select the indication for surgery following the Percutaneous Cardiac Intervention (PCI).

Intent/Clarification: Indicate whether surgery was required due to:

- **PCI complication** - complication during PCI necessitating surgical intervention such as dissection or acute occlusion.
- **PCI failure with clinical deterioration** - PCI failed to yield expected and/or desired results, patient condition deteriorated, includes attempts to cross with the wire but unsuccessful.
- **PCI for STEMI, multi-vessel disease** - STEMI with primary PCI of culprit lesion and multi-vessel disease requiring CABG.
- **PCI failure without clinical deterioration** - PCI failed to yield expected and/or desired results, patient condition did not deteriorate, includes attempts to cross with the wire but unsuccessful.
- **PCI/Surgery staged procedure (not STEMI)** - PCI and surgical procedures performed in a staged fashion in a patient not experiencing STEMI.
- **Other** - other indication for surgery not described above.

SEQ. #: 790

Long Name: Previous PCI-Stent

Short Name: POCPCISt

Definition: Indicate whether an intracoronary stent was used during the previous Percutaneous Cardiac Intervention (PCI).

Intent/Clarification:

A stent is a small mesh tube that's used to treat narrow or weak arteries. It is placed in an artery as part of a procedure called percutaneous coronary intervention (PCI). PCI restores blood flow through narrow or blocked arteries and helps support the inner wall of the artery in the months or years after PCI. Doctors may also place stents in weak arteries to improve blood flow and help prevent the arteries from bursting. Stents usually are made of metal mesh, but sometimes they're made of fabric. Fabric stents, also called stent grafts, are used in larger arteries.

SEQ. #: 795

Long Name: Previous PCI-Stent Type

Short Name: POCPCIStTy

Definition: Indicate type of intracoronary stent placed.

Intent/Clarification:

Choices are:

- **Bare metal**
- **Drug-eluting** - coated with medicine that is slowly and continuously released into the artery. The medicine is intended to prevent the artery from becoming blocked again.
 - **Bio-resorbable** - A bio-resorbable, biodegradable, or bio-absorbable stent serves the same purpose, but is manufactured from a material intended to dissolve or be absorbed in the body.
- **Multiple types**
- **Unknown**

SEQ. #: 800

Long Name: Previous PCI-Interval

Short Name: POCPCIn

Definition: Indicate the interval of time between the previous PCI and the current surgical procedure.

Intent/Clarification:

The choices are ≤ 6 hours or > 6 hours prior to OR entry. The timing of surgery after PCI may influence outcomes such as renal failure due to contrast given during PCI.

SEQ. #: 805

Long Name: Previous Other Cardiac **Short Name:** POC

Definition: Indicate whether the patient had any other previous cardiac intervention.

Intent/Clarification:

If the patient had any other procedure involving the heart and/or great vessels not mentioned above, choose this field. These do not have to be in chronological order.

SEQ. #: 810

Long Name: Previous Other Cardiac Intervention 1 **Short Name:** POCInt1

Definition: Indicate the first other cardiac intervention that was performed.

Intent/Clarification:

No additional interventions

Ablation, catheter, atrial fibrillation

Ablation, catheter, other or unknown

Ablation, catheter, ventricular

Ablation, surgical, atrial fibrillation

Ablation, surgical, other or unknown

Aneurysmectomy, LV

Aortic procedure, arch

Aortic procedure, ascending

Aortic procedure, descending

Aortic procedure, root

Aortic procedure, thoracoabdominal

Aortic Procedure, TEVAR

Aortic root procedure, valve sparing

Atrial appendage obliteration, Left, surgical

Atrial appendage obliteration, Left, transcatheter

Cardiac Tumor

Cardioversion(s)

Closure device, atrial septal defect

Closure device, ventricular septal defect

Congenital cardiac repair, surgical

ECMO

Implantable Cardioverter Defibrillator (ICD) with or without pacemaker

Pacemaker

Pericardial Window/Pericardiocentesis

Pericardiectomy

Pulmonary thromboembolectomy

Total Artificial Heart (TAH)

Transmyocardial Laser Revascularization (TMR)

Transplant heart & lung

Transplant, heart

Transplant, lung(s)

Ventricular Assist Device (VAD), BiVAD

Ventricular Assist Device (VAD), left
Ventricular Assist Device (VAD), right
Other Cardiac Intervention (not listed)

SEQ. #: 815

Long Name: Previous Other Cardiac Intervention 2

Short Name: POCInt2

Definition: Indicate the second other cardiac intervention that was performed.

Intent/Clarification:

If a second procedure was done, please select from the list above or select “No Additional Interventions,” software will grey out any additional selections.

SEQ. #: 820

Long Name: Previous Other Cardiac Intervention 3

Short Name: POCInt3

Definition: Indicate the third other cardiac intervention that was performed.

Intent/Clarification:

If a third procedure was done, please select from the list above or select: No Additional Interventions - Software will grey out any additional selections.

SEQ. #: 825

Long Name: Previous Other Cardiac Intervention 4

Short Name: POCInt4

Definition: Indicate the fourth other cardiac intervention that was performed.

Intent/Clarification:

If a fourth procedure was done, please select from the list above or select “No Additional Interventions,” software will grey out any additional selections.

SEQ. #: 830

Long Name: Previous Other Cardiac Intervention 5

Short Name: POCInt5

Definition: Indicate the fifth other cardiac intervention that was performed.

Intent/Clarification:

If a fifth procedure was done, please select from the list above or select “No Additional Interventions,” software will grey out any additional selections.

SEQ. #: 835

Long Name: Previous Other Cardiac Intervention 6

Short Name: POCInt6

Definition: Indicate the sixth other cardiac intervention that was performed.

Intent/Clarification:

If a sixth procedure was done, please select from the list above or select “No Additional Interventions,” software will grey out any additional selections.

SEQ. #: 840

Long Name: Previous Other Cardiac Intervention 7

Short Name: POCInt7

Definition: Indicate the seventh other cardiac intervention that was performed.

Intent/Clarification:

If a seventh procedure was done, please select from the list above or select “No Additional Interventions,” software will grey out any additional selections.

Preoperative Cardiac Status

SEQ. #: 885

Long Name: Prior MI

Short Name: PrevMI

Definition: Indicate if the patient has had at least one documented previous myocardial infarction at any time prior to this surgery. (Refer to training manual for MI definition.)

Intent/Clarification:

Indicate if the patient has a history of MI. A myocardial infarction is evidenced by **any of the following** in addition to a rise and fall of cardiac biomarkers (preferably troponin) with at least one of the values in the abnormal range for that laboratory [typically above the 99th percentile of the upper reference limit (URL) for normal subjects] **together with at least one of the following** manifestations of myocardial ischemia:

- 1) Ischemic symptoms;
 - a) ECG changes indicative of new ischemia (new ST-T changes, new LBBB, or loss of R- wave voltage)
 - b) Development of pathological Q waves in ≥ 2 contiguous leads on ECG (or equivalent findings for posterior MI)
 - c) Imaging evidence of new loss of viable myocardium or new regional wall motion abnormality
 - d) Documentation in the medical record of the diagnosis of acute myocardial infarction based on the cardiac biomarker pattern in the absence of any items

- enumerated in and due to conditions that may mask their appearance (e.g., peri-operative infarct when the patient cannot report ischemic symptoms; baseline left bundle branch block or ventricular pacing)
- 2) ECG changes associated with prior myocardial infarction can include the following (with or without prior symptoms):
 - a) Any Q wave in leads V2-V3 ≥ 0.02 seconds or QS complex in leads V2 and V3.
 - b) Q wave ≥ 0.03 seconds and ≥ 0.1 mV deep or QS complex in leads I, II, aVL, aVF, or V4-V6 in any two leads of a contiguous lead grouping (I, aVL, V6; V4-V6; II, III, and aVF).
 - c) R-wave ≥ 0.04 seconds in V1-V2 and R/S ≥ 1 with a concordant positive T-wave in the absence of a conduction defect.
 - 3) Imaging evidence of a region with new loss of viable myocardium at rest in the absence of a non-ischemic cause. This can be manifest as:
 - a) Echocardiographic, CT, MR, ventriculographic or nuclear imaging evidence of left ventricular thinning or scarring and failure to contract appropriately (i.e., hypokinesis, akinesis, or dyskinesis)
 - b) Fixed (non-reversible) perfusion defects on nuclear radioisotope imaging (e.g., MIBI, thallium)
 - 4) Medical records documentation of prior myocardial infarction.

Do not use phrases such as “cannot rule out”, “suggestive”, “probable”, “cannot exclude”, etc. to code MI.

SEQ. #: 890

Long Name: MI-When

Short Name: MIWhen

Definition: Indicate the time period between the last documented myocardial infarction and surgery.

Intent/Clarification:

Time of surgery is documented as the hour the patient entered the operating room. Select the time-interval category based on information available on when the MI occurred. MI occurrence is the time of diagnosis and/or when confirmation of the last MI is documented prior to surgery. If the EKG indicates a prior MI of undetermined age Code as >21 days if the patient has no recently reported or documented symptoms. More recent infarctions would likely be described as “evolving” on the EKG.

SEQ. #: 895

Long Name: Cardiac Presentation/Symptoms - At Time Of This Admission

Short Name: CardSympTimeOfAdm

Definition: Indicate the patient's cardiac symptoms at the time of this admission.

Intent/Clarification:

Indicate the patient's cardiac presentation / symptoms. Choose the worst status.

Time Frame: The highest value at the time of admission. If this is a subsequent episode of care, within 7 days, do not code the CAD Presentation from the previous episode of care.

If the patient presents with atypical symptoms of myocardial ischemia (i.e. only shortness of breath, upper abdominal pain, left arm pain, etc.) that is known and documented to be myocardial ischemia, and is considered to be an angina equivalent, code the selection that fits their presentation. If these symptoms are not thought to be, or have not been proven to be the anginal equivalent, code "No Symptoms".

- **No symptoms** – No angina, no acute STEMI, non-STEMI, no anginal equivalent, and no other atypical chest pain.
- **Stable angina** without a change in frequency or pattern for the 6 weeks prior. Angina is controlled by rest and/or oral or transcutaneous medications.
- **Unstable angina:** There are three principal presentations of unstable angina.
 - Rest angina (occurring at rest and prolonged, usually >20 minutes)
 - New-onset angina (within the past 2 months, of at least Canadian Cardiovascular Society Class III severity)
 - Increasing angina (previously diagnosed angina that has become distinctly more frequent, longer in duration, or increased by 1 or more Canadian Cardiovascular Society class to at least CCS III severity)
- **Non-STEMI** The patient was hospitalized for a non-ST elevation myocardial infarction (STEMI) as documented in the medical record. Non-STEMIs are characterized by the presence of **both** criteria:
 - Cardiac biomarkers (creatinine kinase-myocardial band, Troponin T or I) exceed upper limit of normal according to the individual hospitals. Laboratory confirmation of myocardial necrosis; laboratory parameters with a clinical presentation consistent or suggestive of ischemia. ECG changes and/or ischemic symptoms may or may not be present.
 - Absence of ECG changes diagnostic of a STEMI (see STEMI).
- **ST-Elevation MI (STEMI)** or equivalent. The patient presented with a ST elevation myocardial infarction (STEMI) or its equivalent as documented in the medical record. STEMI is characterized by the presence of both criteria:
 - ECG evidence of STEMI: New/presumed new ST-segment elevation or new left bundle branch block not documented to be resolved within 20 minutes. ST-segment elevation is defined by new or presumed new sustained ST-segment elevation at the J-point in two contiguous ECG leads with the cut-off points: ≥ 0.2 mV in men or ≥ 0.15 mV in women in leads V2- V3 and/or ≥ 0.1 mV in other leads and lasting greater than or equal to 20 minutes. If no exact ST-elevation measurement is recorded in the medical chart, physician's written documentation of ST-elevation or Q waves is acceptable. If only one ECG is performed, then the assumption that the ST elevation persisted at least the required 20 minutes is acceptable. Left bundle branch block (LBBB) refers to new or presumed new LBBB on the initial ECG. Cardiac biomarkers (creatinine kinase-myocardial band, Troponin T or I) exceed the upper limit of normal according to the individual hospital's laboratory parameters and a clinical presentation which is consistent or suggestive of ischemia. Note: For purposes of the Registry, ST elevation in the posterior chest leads (V7 through V9), or ST depression that is maximal in V1-3, without ST-segment elevation in other leads, demonstrating posterobasal myocardial infarction, is considered a STEMI equivalent.
- **Anginal Equivalent** - An anginal equivalent is a symptom such as shortness of breath (dyspnea), diaphoresis, extreme fatigue, or belching, occurring in a patient at high cardiac risk. Anginal equivalents are considered to be symptoms of

myocardial ischemia. Anginal equivalents are considered to have the same importance as angina pectoris in patients presenting with elevation of cardiac enzymes or certain EKG changes which are diagnostic of myocardial ischemia. For the patient with diabetes who presents with “silent angina”, code anginal equivalent.

- **Other** – Aortic dissections, sudden death, heart block, arrhythmia, syncope or heart failure.

SEQ. #: 900

Long Name: Cardiac Symptoms - At Time Of Surgery

Short Name: CardSympTimeOfSurg

Definition: Indicate the patient's cardiac symptoms at the time of this procedure.

Intent/Clarification:

The intent is to capture changes between admission and surgery; whether a patient improves or deteriorates. The definition is the same as Seq. #895, although timeframes may overlap.

- For elective admissions, patient symptoms, same value/answer, will be entered twice for seq. #895 and 900.
- If the patient did not improve or deteriorate between admission and surgery, the code will be the same.
- If the patient presents with STEMI or Non-STEMI, they should be coded as such in both sequence numbers 895 and 900.
- If the patient remains longer than 7 days and in that case presentation at the time of admission would be STEMI or Non-STEMI and at the time of surgery would be coded as unstable angina.
- Unstable angina at the time of admission would be coded unstable angina at the time of surgery.

SEQ. #: 911

Long Name: Heart Failure

Short Name: HeartFail

Definition: Indicate whether there is physician documentation or report that the patient has been in a state of heart failure.

Intent/Clarification:

Heart failure is described as unusual dyspnea on light exertion, recurrent dyspnea occurring in the supine position, fluid retention; or the description of rales, jugular venous distension, pulmonary edema on physical exam, or pulmonary edema on chest x-ray presumed to be cardiac dysfunction. A low ejection fraction alone, without clinical evidence of heart failure does not qualify as heart failure. An elevated BNP without other supporting documentation should not be coded as CHF.

Heart failure is a complex clinical syndrome that results from any structural or functional impairment of ventricular filling or ejection of blood. The cardinal manifestations of HF are dyspnea and fatigue, which may limit exercise tolerance, and fluid retention, which

may lead to pulmonary and/or splanchnic congestion and/or peripheral edema. Some patients have exercise intolerance but little evidence of fluid retention, whereas others complain primarily of edema, dyspnea, or fatigue. Because some patients present without signs or symptoms of volume overload, the term “heart failure” is preferred over “congestive heart failure.” There is no single diagnostic test for HF because it is largely a clinical diagnosis based on a careful history and physical examination.

SEQ. #: 912

Long Name: Heart Failure Timing

Short Name: HeartFailTmg

Definition: Indicate whether heart failure is acute, chronic or both (acute on chronic)

Intent/Clarification:

- Acute heart failure is the rapid onset of symptoms and signs of heart failure and may occur with or without previous cardiac disease occurring within 2 weeks of surgery. Acute decompensated heart failure is a sudden worsening of the signs and symptoms of heart failure, which typically includes difficulty breathing (dyspnea), leg or feet swelling, and fatigue.
 - Chronic heart failure develops gradually over time with symptoms of shortness of breath, lower extremity swelling and fatigue without an acute exacerbation within the 2 weeks prior to admission.
 - Both involves patients with chronic heart failure who presents with acute symptoms presents with a worsening of symptoms within 2 weeks of surgery.
-
-

SEQ. #: 913

Long Name: Heart Failure Type

Short Name: HeartFailType

Definition: Indicate the type of heart failure.

Intent/Clarification:

- Systolic: The left ventricle lacks the force to push enough blood into the circulation.
 - Diastolic: The left ventricle is stiff and fails to relax sufficiently to allow adequate filling.
 - Both: Components of both systolic and diastolic failure exist.
 - Unavailable: The type of heart failure is not documented in the medical record.
-
-

SEQ. #: 915

Long Name: Classification-NYHA

Short Name: ClassNYH

Definition: Indicate the patient's worst dyspnea or functional class, coded as the New York Heart Association (NYHA) classification within the past 2 weeks. This is to be used for heart failure only, is not intended to classify angina.

Intent/Clarification:**NYHA is for congestive heart failure (CHF).**

Select the **highest level** of heart failure within the two weeks leading up to episode of hospitalization or at the time of the procedure. The intent is to capture the highest level of failure. Physician documentation should be in the medical record.

- **Class I:** Patient has cardiac disease but without resulting limitations of ordinary physical activity. Ordinary physical activity (e.g., walking several blocks or climbing stairs) does not cause undue fatigue, palpitation, dyspnea, or anginal pain. Limiting symptoms may occur with marked exertion.
- **Class II:** Patient has cardiac disease resulting in slight limitation of ordinary physical activity. Patient is comfortable at rest. Ordinary physical activity such as walking more than two blocks or climbing more than one flight of stairs results in limiting symptoms (e.g., fatigue, palpitation, dyspnea, or anginal pain).
- **Class III:** Patient has cardiac disease resulting in marked limitation of physical activity. Patient is comfortable at rest. Less than ordinary physical activity (e.g., walking one to two level blocks or climbing one flight of stairs) causes fatigue, palpitation, dyspnea, or anginal pain.
- **Class IV:** Patient has dyspnea at rest that increases with any physical activity. Patient has cardiac disease resulting in inability to perform any physical activity without discomfort. Symptoms may be present even at rest. If any physical activity is undertaken, discomfort is increased. The physician documents new onset CHF with an EF of 25% and SOB. There is no indication of what level of activity causes the SOB.

SEQ. #: 930

Long Name: Cardiogenic Shock

Short Name: CarShock

Definition: Indicate if the patient developed cardiogenic shock. Cardiogenic shock is defined as a sustained (>30 min) episode of hypoperfusion evidenced by systolic blood pressure <90 mm Hg and/or, if available, cardiac index <2.2 L/min per square meter determined to be secondary to cardiac dysfunction and/or the requirement for parenteral inotropic or vasopressor agents or mechanical support (e.g., IABP, extracorporeal circulation, VADs) to maintain blood pressure and cardiac index above those specified levels.

Note: Transient episodes of hypotension reversed with IV fluid or atropine do not constitute cardiogenic shock. The hemodynamic compromise (with or without extraordinary supportive therapy) must persist for at least 30 min.

ACCF/AHA 2013

Intent/Clarification:

- At the time of the procedure.
- This includes patients with cardiogenic shock who have been stabilized on IABP/inotropes at the time of surgery.
- Do not code yes to cardiogenic shock for patients with a low cardiac index who are asymptomatic and do not require mechanical or inotropic support.
- Hemodynamic issues that could be contributed to anesthesia induction problems

should not count in the preoperative status of the patient.

- Elective procedures should not be coded as cardiogenic shock.
- Do not code yes to cardiogenic shock just because the patient has a LVAD; the patient must meet the blood pressure and/or cardiac index parameters of the definition of cardiogenic shock.

SEQ. #: 935

Long Name: Resuscitation

Short Name: Resusc

Definition: Indicate whether the patient required cardiopulmonary resuscitation before the start of the operative procedure which includes the institution of anesthetic management. Capture resuscitation timeframe: within 1 hour or 1-24 hours pre-op.

Intent/Clarification:

Indicate whether the patient required cardiopulmonary resuscitation within 24 hours of the start of the operative procedure. The start of the procedure begins with the induction of anesthesia. Capture resuscitation timeframe: within 1 hour of surgery or 1-24 hours pre-operatively.

The additional time options were added to harmonize with NCDR, looking at 24 hours pre-procedure yet still mapping to previous STS versions and risk models.

- Resuscitation may include **complete** circulatory support such as ECMO/other mechanical assist devices (ex. Impella, LVAD) initiated emergently prior to surgery. Intra-aortic balloon counterpulsation (IABP) by itself does not qualify as complete circulatory support.
- Do not code yes for resuscitation started after induction of anesthesia. The goal is to identify patients who require CPR and/or mechanical circulatory support to maintain life in the 24 hour period preceding surgery.

SEQ. #: 945

Long Name: Cardiac Arrhythmia

Short Name: Arrhythmia

Definition: Indicate whether the patient has a history of a cardiac rhythm disturbance before the start of the operative procedure which includes the institution of anesthetic management.

Intent/Clarification:

- Yes
- No

SEQ. #: 947

Long Name: Cardiac Arrhythmia - Permanently Paced Rhythm

Short Name: ArrhythPPaced

Definition: Indicate whether the patient has a permanently paced rhythm, evidenced by pacemaker activity during heart rhythm evaluation.

Intent/Clarification:

- Yes
 - No
-
-

SEQ. #: 950

Long Name: Cardiac Arrhythmia - VTach / VFib

Short Name: ArrhythVV

Definition: Indicate whether arrhythmia was VTach or VFib.

Intent/Clarification:

V-tach rhythm must be sustained/persistent or paroxysmal and require some type of intervention (pharmacological and/or electrical) to interrupt and cease the arrhythmia. Do not include short runs of VT.

- None
 - Remote - more than 30 days prior to procedure
 - Recent - within 30 days of this procedure
-
-

SEQ. #: 955

Long Name: Cardiac Arrhythmia - Sick Sinus Syndrome

Short Name: ArrhythSSS

Definition: Indicate whether arrhythmia was sick sinus syndrome.

Intent/Clarification:

Sick sinus syndrome is a collection of heart rhythm disorders caused by dysfunction in the SA node, the heart's main pacemaker. SSS may present as: Sinus bradycardia -- slow heart rates from the natural pacemaker of the heart. Tachycardias - fast heart rates
Bradycardia-tachycardia -- alternating slow and fast heart rhythms

- None
 - Remote - more than 30 days prior to procedure
 - Recent - within 30 days of this procedure
-
-

SEQ. #: 960

Long Name: Cardiac Arrhythmia - AFlutter

Short Name: ArrhythAFlutter

Definition: Indicate whether arrhythmia was atrial flutter.

Intent/Clarification:

Atrial flutter (AFL) is an abnormal heart rhythm that occurs in the atria of the heart. When it first occurs, it is usually associated with a fast heart rate or tachycardia (beats over 100 per minute) which falls into the category of supra-ventricular tachycardias. While this rhythm occurs most often in individuals with cardiovascular disease (e.g. hypertension, coronary artery disease, and cardiomyopathy) and diabetes, it may occur spontaneously in people with otherwise normal hearts. It is typically not a stable rhythm, and frequently degenerates into atrial fibrillation (AF). However, it does rarely persist for months to years. If rhythm is described as fib/flutter, code fibrillation.

- None
 - Remote - more than 30 days prior to procedure
 - Recent - within 30 days of this procedure
-
-

SEQ. #: 961

Long Name: Cardiac Arrhythmia - Atrial Fibrillation

Short Name: ArrhythAtrFib

Definition: Indicate whether arrhythmia was atrial fibrillation.

Intent/Clarification:

- None
 - Remote - more than 30 days prior to procedure
 - Recent - within 30 days of this procedure
-
-

SEQ. #: 962

Long Name: Cardiac Arrhythmia - Atrial Fibrillation - Type

Short Name: ArrhythAFib

Definition: Indicate whether arrhythmia was atrial fibrillation and if so, which type.

Intent/Clarification:

If the diagnosis of atrial fibrillation is present code the type:

- Paroxysmal: Recurrent AF (> 2 episodes). Terminates spontaneously within 7 days.
- Persistent: Sustained episode > 7 days, or lasting < 7 days, but necessitating pharmacologic or electrical cardioversion.
- Long-Standing Persistent: Continuous episode of > 1 year duration.
- Permanent: Continuous episode of > 1 year duration.

FAQ August 2017: The definition of longstanding persistent and permanent are the same, can you clarify the difference?

Answer: Longstanding persistent atrial fibrillation lasts longer than 1 year but still responds to treatment ie. cardioversion or ablation therapy.

Permanent atrial fibrillation lasts longer than 1 year but no longer responds to any form of treatment. These patients are treated for rate control and prevention of stroke only.

SEQ. #: 965

Long Name: Cardiac Arrhythmia - Second Degree Heart Block

Short Name: ArrhythSecond

Definition: Indicate whether arrhythmia was second degree heart block.

Intent/Clarification:

In second degree heart block, some signals from the atria don't reach the ventricles. This causes "dropped beats." On an ECG, the P wave isn't followed by the QRS wave, because the ventricles weren't activated. There are two types: Type I second-degree heart block, or Mobitz Type I, or Wenckebach's AV block. Electrical impulses are delayed more and more with each heartbeat until a beat is skipped. This condition is not too serious but sometimes causes dizziness and/or other symptoms. Type II second-degree heart block, or Mobitz Type II. This is less common than Type I but generally more serious. Because electrical impulses can't reach the ventricles, an abnormally slow heartbeat may result. In some cases a pacemaker is needed.

- None
- Remote -more than 30 days prior to procedure
- Recent -within 30 days of this procedure

SEQ. #: 970

Long Name: Cardiac Arrhythmia - Third Degree Heart Block

Short Name: ArrhythThird

Definition: Indicate whether arrhythmia was third degree heart block.

Intent/Clarification:

Heart block is applicable only if the patient has or did have 3rd degree heart block (complete heart block). Complete heart block, also referred to as third-degree heart block, or third-degree atrioventricular (AV) block, is a disorder of the cardiac conduction system where there is no conduction through the AV node. Therefore, complete dissociation of the atrial and ventricular activity exists.

- None
- Remote- more than 30 days prior to procedure
- Recent - within 30 days of this procedure

Preoperative Medications

SEQ. #: 1020

Long Name: Meds-ACE Inhibitors or ARB Within 48 Hours

Short Name: MedACEI48

Definition: Indicate whether the patient received ACE Inhibitors or ARB within 48 hours preceding surgery (e.g., if indicated for LV dysfunction or acute MI).

Intent/Clarification:

ACE and ARBs are used in the treatment of hypertension, congestive heart failure (reduces the workload of the heart). The drug action is to inhibit the release of the

hormone angiotensin II that constricts blood vessels, causing an increase in blood pressure. Therefore, blood vessels dilate to increase systemic blood flow to the heart. Some ACE inhibitors have additional diuretic components to increase the elimination of excess fluid.

Studies have shown that preoperative use of ACEI/ARB is associated with a 27.6% higher risk for Acute Kidney Injury (AKI) postoperatively. Stopping ACEI or ARB before cardiac surgery may reduce the incidence of AKI. This includes renin inhibitors.

- **Yes** - Capture those who are prescribed to take medications on a regular schedule and are presumed to be at a therapeutic level, 48 hours preceding surgery, (entry into the OR)
- **No** - did not receive an ACE inhibitor or ARB within 48 hours preceding surgery
- **Contraindicated** - Documented evidence of contraindication: If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record (notation of a medication allergy prior to arrival) by physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist
- **Unknown** - conflicting information in the medical record and/or with the patient/family or no information available

SEQ. #: 1025

Long Name: Meds- Amiodarone Prior To Surgery

Short Name: MedAmiodarone

Definition: Indicate whether and when the patient received Amiodarone therapy prior to surgery. Dronedarone (Multaq) may be coded as Amiodarone.

Intent/Clarification:

Intended to capture **ongoing** medication administration prior to surgery.

Amiodarone may play a role in reducing the risk of post-operative arrhythmias, notably A-Fib.

- **Yes: on home therapy**
- **Yes: therapy started this admission**, can include patients where a preoperative protocol was initiated; this allows differentiation from those patients on long term home therapy.
- **No:** a single dose prior to surgery such as in ED does not count as “Yes,” , only capture those who are prescribed to take medications on a regular schedule and are presumed to be at a therapeutic level, preceding surgery (entry into the OR)
- **Unknown:** conflicting information in the medical record and/or with the patient/family or no information available

SEQ. #: 1030

Long Name: Meds-Beta Blockers Within 24 Hours

Short Name: MedBeta

Definition: Indicate whether or not the patient received beta blockers within 24 hours preceding surgery, or if beta blocker was contraindicated. The contraindication must be

documented in the medical record by a physician, nurse practitioner, or physician assistant. A "hold order" is not considered a contraindication.

Intent/Clarification:

NQF Endorsed Measure - Part of the medication bundle in the STS Composite Quality Rating (Star Rating).

Beta blockers have been proven to increase survival in cardiac patients. For the treatment of:

1. High blood pressure
2. Treating chest pain or angina
3. Controlling irregular heart rhythms, prevention of post op Afib
4. Slowing ventricular rate response
5. Treating congestive heart failure

Yes- include those who received within 24 hours prior to ***incision in the OR***. This can include onetime doses given prior to ***incision in OR***

No – Patient did not receive prior to ***incision in the OR***

Contraindicated - documented evidence of contraindication: If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record (examples might include allergy, bradycardia, hypotension, heart block, COPD, Other), check “Contraindication.” by physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist. Documents created by hospitals used to track Core Measure Information may be used but would still have to be countersigned by physician, Nurse Practitioner, Anesthesia, Physician Assistant.

SEQ. #: 1035

Long Name: Meds-Beta Blocker Therapy For More Than 2 Weeks Prior To Surgery

Short Name: MedBetaTher

Definition: Indicate whether the patient received beta blocker therapy for at least 2 weeks prior to surgery.

Intent/Clarification:

Studies have shown that the abrupt discontinuation of Beta-Blockers during the perioperative period in patients who were on chronic Beta-Blocker therapy prior to surgery can lead to increased mortality during the intraoperative and postoperative periods. The American College of Cardiology/American Heart Association has given the continuation of Beta-Blocker therapy throughout the perioperative period a Class I recommendation.

- **Yes** - Capture those who are prescribed to take a Beta-Blocker on a regular schedule (daily) and are presumed to be at a therapeutic level, for at least 2 weeks preceding surgery (entry into the OR). Do Not Include a one-time dose.
- **No** – Beta-Blocker was prescribed but patient is not taking a daily dose or not prescribed Beta-Blocker, within the two weeks preceding surgery
- **Contraindicated**- Documented evidence of contraindication: If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record, notation of a medication allergy prior to arrival, by Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist. If a “hold order” has parameters associated with it, this is acceptable as a

contraindication (i.e. hold if HR < 60 & there is documentation of the HR less than 60 in the medication administration record (MAR).

- **Unknown** – conflicting information in the medical record and/or with the patient/family or no information is available

SEQ. #: 1040

Long Name: Meds-Calcium Channel Blocker Therapy For More Than 2 Weeks Prior To Surgery

Short Name: MedCChanTher

Definition: Indicate whether the patient received calcium channel blocker therapy for at least 2 weeks prior to surgery.

Intent/Clarification:

Calcium channel blockers (CCB), calcium channel antagonists or calcium antagonists are a number of medications that disrupts the movement of calcium (Ca²⁺) through calcium channels. Calcium channel blockers are used as antihypertensive drugs, i.e. as medications to decrease blood pressure in patients with hypertension. CCBs are particularly effective against large vessel stiffness, one of the common causes of elevated systolic blood pressure in elderly patients. Calcium channel blockers are also frequently used to alter heart rate, to prevent cerebral vasospasm, and to reduce chest pain caused by angina pectoris

- **Yes** - Capture those who are prescribed to calcium channel blockers on a regular schedule and are presumed to be at a therapeutic level, for at least 2 weeks preceding surgery (entry into the OR)
- **No** – Patient did not receive a Calcium Channel Blocker for at least 2 weeks preceding surgery
- **Contraindicated** - Documented evidence of contraindication: If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record (notation of a medication allergy prior to arrival) by a Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist
- **Unknown** – conflicting information in the medical record and/or with the patient/family or no information is available

SEQ. #: 1045

Long Name: Meds-Long-Acting Nitrate Therapy For More Than 2 Weeks Prior To Surgery

Short Name: MedLongActNit

Definition: Indicate whether the patient received long-acting nitrate therapy for at least 2 weeks prior to surgery.

Intent/Clarification:

- **Yes** - Capture those prescribed to take medications on a regular schedule and are presumed to be at a therapeutic level, for at least 2 weeks preceding surgery (entry into the OR). Nitropaste or Nitropatch are long-acting nitrates. Do Not include a one-time dose

- **No** – Patient did not receive a Long-Acting Nitrate for at least 2 weeks preceding surgery
 - **Contraindicated** - Documented evidence of contraindication: If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record (notation of a medication allergy prior to arrival) by a Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist -BH
 - **Unknown** – conflicting information in the medical record and/or with the patient/family or no information is available
-
-

SEQ. #: 1050

Long Name: Meds-Nitrates-I.V. Within 24 Hours

Short Name: MedNitIV

Definition: Indicate whether the patient received IV Nitrates within 24 hours preceding surgery.

Intent/Clarification:

Nitrates act by increasing dilatation of the coronary arteries, thereby increasing blood flow to the myocardium and decreasing myocardial ischemic changes. Trade name is Nitroglycerin.

- **Yes** - Capture those who are prescribed to take IV Nitrates and are presumed to be at a therapeutic level, 24 hours preceding surgery (entry into the OR)
 - **No** – Patient did not receive IV Nitrates within 24 hours preceding surgery
-
-

SEQ. #: 1055

Long Name: Meds-Other Antianginal Medication Therapy For More Than 2 Weeks Prior To Surgery

Short Name: MedOthAntiang

Definition: Indicate whether the patient received any other antianginal medication therapy for at least 2 weeks prior to surgery.

Intent/Clarification:

- **Yes** - Capture those who are prescribed to take any other antianginal medication on a regular schedule and are presumed to be at a therapeutic level, for at least 2 weeks preceding surgery (entry into the OR) - Do Not Include a one-time dose
- **No** – Patient did not receive any other antianginal medication therapy for at least 2 weeks preceding surgery. Do not capture if patient was given a sublingual, IV, or short acting formula of one of these medications.
- **Contraindicated** - Documented evidence of contraindication: If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record (notation of a medication allergy prior to arrival) by a Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist -BH
- **Unknown** – conflicting information in the medical record and/or with the patient/family

SEQ. #: 1060

Long Name: Meds-ADP Inhibitors Within Five Days

Short Name: MedADP5Days

Definition: Indicate whether the patient has received ADP Inhibitors within 5 days preceding surgery.

Intent/Clarification:

ADP stands for Adenosine Diphosphate. The anticoagulant properties of these medications may increase the risk of bleeding by inhibiting platelet aggregation (clotting). This category includes P2Y12 inhibitors. They are often used to treat patients with a history of atherosclerotic cardiovascular disease and potentially reduce the incidence of major cardiovascular events (stroke, peripheral arterial disease events). Peak drug levels are reached within 3-7 days of initiated maintenance dosing, while termination of drug effects are not seen for 5 days after last dose.

- **Yes** - Capture those who are prescribed to ADP inhibitors on a regular schedule and are presumed to be at a therapeutic level within 5 day preceding surgery (entry into the OR) **and** those who received a one-time dose of Plavix, preceding surgery
- **No** - did not receive an ADP inhibitor within 5 days preceding surgery
- **Contraindicated** - Documented evidence of contraindication: If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record (notation of a medication allergy prior to arrival) by a Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist
- **Unknown** – conflicting information in the medical record and/or with the patient/family or no information available

FAQ October 2017: ADP is coded 'Yes' for patients who receive a single dose of Plavix, is that true for Brillinta and Effient?

Answer: Code 'Yes' for one-time doses of Brillinta and Effient.

SEQ. #: 1065

Long Name: Meds-ADP Inhibitors Discontinuation

Short Name: MedADPIDis

Definition: Indicate the number of days prior to surgery ADP Inhibitor use was discontinued. If less than 24 hours, enter "0".

Intent/Clarification:

Peak drug levels are reached within 3-7 days of initiated maintenance dosing, while termination of drug effects are not seen for 5 days after last dose, which may increase risk of bleeding.

SEQ. #: 1070

Long Name: Meds-Aspirin Within Five Days

Short Name: MedASA

Definition: Indicate whether or not the patient received Aspirin or Ecotrin within 5 days preceding surgery.

Intent/Clarification:

Anti-inflammatory, analgesic and antiplatelet action. Half-life of aspirin products is 5-7 days. Aspirin use may predispose patient to post op bleeding.

- **Yes** - Capture those who are prescribed to take Aspirin or Ecotrin on a regular schedule and are presumed to be at a therapeutic level, 5 days preceding surgery (entry into the OR) - The minimum dose should be at least 75 mg (i.e. Aggrenox, which is only 25mg, should not be included).
- **No** – Patient did not receive Aspirin within 5 days preceding surgery.
- **Contraindicated** - Documented evidence of contraindication: If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record (notation of a medication allergy prior to arrival) by a Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist
- **Unknown** – conflicting information in the medical record and/or with the patient/family or no information available.

SEQ. #: 1071

Long Name: Meds-Aspirin Discontinuation

Short Name: MedASADis

Definition: Indicate the number of days prior to surgery Aspirin use was discontinued. If less than 24 hours, enter "0".

Intent/Clarification: -

SEQ. #: 1072

Long Name: Meds-Aspirin One-Time Dose

Short Name: MedASAOnce

Definition: Indicate whether the patient received a one-time dose of Aspirin and is not on daily aspirin.

Intent/Clarification: -

SEQ. #: 1073

Long Name: Meds-Glycoprotein IIb/IIIa Inhibitor Within 24 Hours

Short Name: MedGP

Definition: Indicate whether the patient received Glycoprotein IIb/IIIa inhibitors within 24 hours preceding surgery.

Intent/Clarification:

- **Yes:** if the patient received a IIb/IIIa inhibitor within 24 hours of OR entry date and

time.

- **No**

SEQ. #: 1075

Long Name: Meds-Anticoagulants Within 48 Hours

Short Name: MedACoag

Definition: Indicate whether the patient received IV and/or sub-q anticoagulants within 48 hours preceding surgery.

Do NOT include Coumadin or one-time boluses of Heparin.

Intent/Clarification:

Anticoagulant therapy inhibits platelet aggregation and clot formation, is used to treat and prevent blood clots, decreasing the viscosity of the blood. These medications may increase the risk of bleeding

- **Yes:** Only capture those who are prescribed to take IV and/or Sub-Q anticoagulants on a regular schedule and are presumed to be at a therapeutic level, within 48 hours preceding surgery (entry into the OR).
- **No:** Patient did not receive IV and/or Sub-Q anticoagulants within 48 hour preceding surgery. Do not capture one-time heparin, Lovenox/Enoxaparin doses used during the cardiac cath or any time within 48 hours preceding surgery.

SEQ. #: 1080

Long Name: Meds-Anticoagulants-Medication Name

Short Name: MedACMN

Definition: Indicate the name of the anticoagulant the patient received within 48 hours preceding surgery.

Intent/Clarification:

- Heparin (Unfractionated)
- Heparin (Low Molecular)
- Both
- Other

SEQ. #: 1091

Long Name: Meds-Warfarin (Coumadin) Within 5 Days

Short Name: MedCoum5Days

Definition: Indicate whether the patient has received Warfarin (Coumadin) within 5 days preceding surgery.

Intent/Clarification:

This is collected to capture the risk of bleeding related to anticoagulation therapy.

- **Yes** - Capture those who took Coumadin within 5 days preceding surgery and are

- presumed to be at a therapeutic level within 24 hours prior to OR entry date and time.
- **No** – Patient did not receive a Coumadin within 5 days prior to OR entry date and time.
 - **Unknown** – Conflicting information in the medical record and/or with the patient/family or no information is available
- -----

SEQ. #: 1092

Long Name: Meds-Warfarin (Coumadin) Discontinuation

Short Name: MedCoun5Dis

Definition: Indicate the number of days prior to surgery Warfarin (Coumadin) use was discontinued. If less than 24 hours, enter "0".

Intent/Clarification:

SEQ. #: 1101

Long Name: Meds-Factor Xa Inhibitors Within 5 Days

Short Name: MedXa5Days

Definition: Indicate whether the patient has received Factor Xa Inhibitors within 5 days preceding surgery.

Intent/Clarification:

Direct factor Xa inhibitors ("xabans ") are a class of anticoagulant drugs which act directly upon Factor X in the coagulation cascade, without using anti-thrombin as a mediator. This is collected to capture risk of bleeding related to anticoagulation therapy

- **Yes** - Capture those who are prescribed to take Factor Xa inhibitors on a regular schedule within the 5 days preceding surgery and are presumed to be at a therapeutic level, within 24 hours prior to OR entry date and time.
 - **No** – Patient did not receive Factor Xa Inhibitors within 5 days prior to OR entry date and time.
 - **Unknown** – Conflicting information in the medical record and/or with the patient/family or no information is available.
- .

FAQ August 2017: Clarify which medications should be coded in this category.

Answer: Confusion lies when the medications could be coded in any or all of three categories, Factor Xa, Thrombin Inhibitors, or NOAC. Initially all three categories were included to differentiate medications that did not have antidotes. With more antidotes available, the three medications seem redundant. Capture the medication according to the manufacturer's category description. The only caveat to remember is that you should only code ONE of the three medications so as not to have it appear that the patient is receiving more than one.

SEQ. #: 1102

Long Name: Meds-Factor Xa Inhibitors Discontinuation

Short Name: MedXa5DDis

Definition: Indicate the number of days prior to surgery Factor Xa Inhibitor use was discontinued. If less than 24 hours, enter "0".

Intent/Clarification:

SEQ. #: 1111

Long Name: Meds-Novel Oral Anticoagulant Within 5 Days

Short Name: MedNOAC5Days

Definition: Indicate whether the patient has received Novel Oral Anticoagulant within 5 days preceding surgery.

Intent/Clarification:

New agents have been introduced that are collectively referred to as **novel oral anticoagulants (NOACs)** or **directly acting oral anticoagulants (DOACs)**. They have been shown to be as good as or possibly better than Coumadin with less serious side effects. The newer anticoagulants (NOACs/DOACs), are more expensive than the traditional ones and should be used with care in patients with kidney problems. Additionally, there is no antidote for the factor Xa inhibitors, so it is difficult to stop their effects in the body in cases of emergency (accidents, urgent surgery). [Idarucizumab](#) was FDA approved for the reversal of dabigatran in 2015.

- Yes - Capture those who are prescribed to take novel oral anticoagulants on a regular schedule within the 5 days preceding surgery and are presumed to be at a therapeutic level, within 24 hours prior to OR entry date and time.
- No - Patient did not receive a novel oral anticoagulant within 5 days prior to OR entry date and time.
- Unknown - Conflicting information in the medical record and/or with the patient/family or no information is available.

FAQ August 2017: Clarify which medications should be coded in this category.

Answer: Confusion lies when the medications could be coded in any or all of the three categories, Factor Xa, Thrombin Inhibitors, or NOAC. Initially all three categories were included to differentiate medications that did not have antidotes. With more antidotes available, the three medications seem redundant. Capture the medication according to the manufacturer's category description. The only caveat to remember is that you should only code ONE of the three medications so as not to have it appear that the patient is receiving more than one.

SEQ. #: 1112

Long Name: Meds-Novel Oral Anticoagulant Discontinuation

Short Name: MedNOACDisc

Definition: Indicate the number of days prior to surgery Novel Oral Anticoagulant use was discontinued. If less than 24 hours, enter "0".

Intent/Clarification:

SEQ. #: 1121

Long Name: Meds-Thrombin Inhibitors Within 5 Days

Short Name: MedThromIn5Days

Definition: Indicate whether the patient has received Thrombin Inhibitors within 5 days preceding surgery.

Intent/Clarification:

- **Yes** - Capture those who are prescribed to take thrombin inhibitors within the 5 days preceding surgery and are presumed to be at a therapeutic level, within 24 hours preceding surgery (entry into the OR) - Do Not Include a one-time dose
- **No** – Patient did not receive a Thrombin Inhibitor medication within 5 days prior to OR entry date and time.
- **Contraindicated** - Documented evidence of contraindication: If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record (notation of a medication allergy prior to arrival) by a Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist
- **Unknown** – conflicting information in the medical record and/or with the patient/family or no information is available.

FAQ August 2017: Clarify which medications should be coded in this category.
Answer: Confusion lies when the medications could be coded in any or all of three categories, Factor Xa, Thrombin Inhibitors, or NOAC. Initially all three categories were included to differentiate medications that did not have antidotes. With more antidotes available, the three medications seem redundant. Capture the medication according to the manufacturer's category description. The only caveat to remember is that you should only code ONE of the three medications so as not to have it appear that the patient is receiving more than one.

SEQ. #: 1122

Long Name: Meds-Thrombin Inhibitors Discontinuation

Short Name: MedThromInDisc

Definition: Indicate the number of days prior to surgery Thrombin Inhibitor use was discontinued. If less than 24 hours, enter "0".

Intent/Clarification:

SEQ. #: 1125

Long Name: Meds-Thrombolytics Within 24 Hours

Short Name: MedThrom

Definition: Indicate whether the patient received thrombolytics within 24 hours preoperatively.

Intent/Clarification:

Thrombolytic (fibrinolytic) therapy is the use of drugs to break up or dissolve blood clots, which are the main cause of both heart attacks and stroke. It can predispose a patient to bleeding if given within 24 hours prior to surgery. There are three major classes of thrombolytic drugs: tissue plasminogen activator (tPA), streptokinase (SK), and urokinase (UK). This includes one-time doses.

- **Yes** - Capture those who received thrombolytics within 24 hours preceding of OR entry date and time.
- **No** – Patient did not receive thrombolytics within 24 hours preceding surgery.
- **ONLY CAPTURE DOSES GIVEN WITHIN 24 HOURS NOT 48 AS LISTED IN THE V2.9 DATA COLLECTION FORM.**

SEQ. #: 1130

Long Name: Meds-Inotropes Within 48 Hours

Short Name: MedInotr

Definition: Indicate whether the patient received IV inotropic agents within 48 hours preceding surgery.

Intent/Clarification:

Positive Inotropic agent actions act at the cellular level, increasing intracellular calcium. Cardiovascular effects range from increasing or decreasing the heart rate, increasing force of the heart muscle contraction, peripheral or extremity arterial or venous constriction. The degree to which these systems are affected are dose dependent. As well, these drugs may lose their cardiovascular effect causing a negative response at higher dosing levels. Initiation of these drugs typically is in response to some hemodynamic instability in the patient.

This field is in the risk models. Use of inotropic agents preoperatively is associated with increased risk of mortality and morbidity- including renal failure, prolonged vent, reoperation, and length of stay.

- **Yes** - Capture those who received IV inotropic agent(s), within 48 hours preceding OR entry date and time.
- **No** – Patient did not receive Inotropes within 48hours preceding surgery.

SEQ. #: 1135

Long Name: Meds-Lipid Lowering Within 24 Hours

Short Name: MedLipid

Definition: Indicate whether or not the patient received lipid lowering medication within 24 hours preceding surgery.

Intent/Clarification:

Capture medications administered to lower the total cholesterol, LDL, HDL or triglyceride

levels. Patient may be on prescribed medication and have normal cholesterol values, these patients should still be coded as “Yes,” for dyslipidemia.

Note that non-statis are listed here but are no longer considered effective lipid lowering agents according to AHA guidelines. **Only statins will count in the measure component of the composite.** A contraindication to statins takes the patient out of the denominator. This will be noted in the report.

- **Yes** - Capture those who are prescribed to take lipid-lowering medication on a regular schedule and are presumed to be at a therapeutic level 24 hours preceding surgery (entry into the OR) - Do Not Include a one-time dose
- **No** – Patient did not receive a lipid lowering medication within 24 hours preceding surgery
- **Contraindicated** - Documented evidence of contraindication: If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record (notation of a medication allergy prior to arrival) by physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist
- **Unknown** – Conflicting information in the medical record and/or with the patient/family or no information is available

SEQ. #: 1141

Long Name: Meds-Lipid Lowering-Medication Type

Short Name: MedLipType

Definition: Indicate the type of lipid lowering medication the patient received within 24 hours preceding surgery.

Intent/Clarification:

- Statin
- Statin + Other
- Non-Statin/Other

SEQ. #: 1143

Long Name: Meds-Steroids Within 24 Hours

Short Name: MedSter

Definition: Indicate whether the patient was taking steroids within 24 hours of surgery. This does not include a one-time dose related to prophylaxis therapy (i.e. for IV dye exposure during cath procedure or surgery pre-induction period). Non-systemic medications are not included in this category (i.e., nasal sprays, topical creams).

Intent/Clarification:

Systemic delivery only. Non-systemic delivery is not included in this data element. Non-systemic delivery includes topical creams, nasal sprays, inhalers or ophthalmic or otic drops.

Do not include one-time systemic dose as part of clinical pathway guideline or procedure/surgical preparatory order.

- **Yes** - Capture those who are prescribed to take **systemic steroids** within 24 hours preceding surgery and are presumed to be at a therapeutic level within 24

hours preceding OR entry date and time - Do Not Include a one-time dose.

- **No** – Patient did not receive a systemic steroids within 24 hours preceding surgery
- **Contraindicated** - Documented evidence of contraindication: If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record (notation of a medication allergy prior to arrival) by physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist -BH
- **Unknown** – Conflicting information in the medical record and/or with the patient/family or no information is available

Hemodynamics/Cath/Echo

****General Information: Hemodynamic values for ejection fraction, pulmonary artery pressure, and valve insufficiency and stenosis should be captured from studies done **closest to the time of surgery.****

SEQ. #: 1145

Long Name: Cardiac Catheterization Performed

Short Name: CarCathPer

Definition: Indicate whether cardiac catheterization and/or CT angio was performed.

Intent/Clarification:

Diagnostic coronary angiography is defined as the passage of a catheter into the aortic root or other great vessels for the purpose of angiography of the native coronary arteries or bypass grafts supplying native coronary arteries.

Capture procedures done within 6 months prior to surgery. Do not include stand-alone right heart catheterization in this field; include coronary angiogram either done with or without right and/or left heart pressures.

SEQ. #: 1150

Long Name: Cardiac Catheterization Date

Short Name: CarCathDt

Definition: Indicate the date cardiac catheterization was performed.

Intent/Clarification:

If more than one was performed, capture the date closest to surgery. Do not include stand-alone RHC (right heart cath) in this field. While it is preferred that the cath be done within 6 months, they can be used for up to one year.

Required date format: mm/dd/yyyy.

SEQ. #: 1155

Long Name: Coronary Anatomy/Disease Known

Short Name: CorAnatDisKnown

Definition: Indicate whether coronary artery anatomy and/or disease is documented and available prior to surgery.

Intent/Clarification:

Indicated if coronary artery anatomy and/or disease is documented **prior** to surgery. Sometimes the results are known and verbally communicated to the surgeon, but the Cath Lab Report is not documented in the medical record until after surgery has started; this is particularly true for emergent cases. This can be captured even if dictation was not completed until after the surgery. Results dictated following the procedure may be used.

SEQ. #: 1160

Long Name: Dominance

Short Name: Dominance

Definition: Indicate whether coronary artery dominance is documented prior to surgery.

Intent/Clarification:

- **Left** - The posterior descending artery (PDA) arises from the left circumflex artery.
- **Right** - The posterior descending artery (PDA) arises from the right coronary artery.
- **Co-dominant** - The right coronary artery supplies the posterior descending artery (PDA) and the circumflex supplies the posterolateral artery (PLA). Thus, there is approximately equal contribution to the inferior surface of the left ventricle from both the left circumflex and right coronary arteries.
- **Not documented**

SEQ. #: 1165

Long Name: Source(s) Used To Quantify Stenosis

Short Name: StenSource

Definition: Indicate source or sources used to quantify coronary artery stenosis.

Intent/Clarification:

- Angiogram
- CT
- IVUS
- Progress/OP Note
- Other
- Multiple

If multiple sources are available, select surgeon "s documentation of the degree of stenosis. This is the degree of stenosis the surgeon used to develop the operative plan.

FAQ August 2017: Should IVUS results be used for the LM only or can it be used to quantify disease in any artery?

Answer: IVUS can be used to quantify the stenosis in any artery.

SEQ. #: 1170

Long Name: Num Dis Vessels

Short Name: NumDisV

Definition: Indicate the number of diseased major native coronary vessel systems: LAD system, Circumflex system, and/or Right system with $\geq 50\%$ narrowing of any vessel preoperatively.

NOTE: Left main disease ($\geq 50\%$) is counted as TWO vessels (LAD and Circumflex, which may include a Ramus Intermedius). For example, left main and RCA would count as three total.

A vessel that has ever been considered diseased, should always be considered diseased.

Intent/Clarification:

There are three (3) major coronary systems; Left Anterior Descending (LAD), Circumflex and Right Coronary System (RCA). Each system has "branches" that are considered part of their corresponding system. Vessel stenosis or narrowing is measured in percentages (%), most often expressed as a range of "stenosis".

The Ramus Intermedius is a vessel that can function as part of the LAD system or as part of the Circumflex system depending on its course. If the Ramus is part of the LAD system and functions much like a diagonal, code 1 vessel disease. If the Ramus is part of the Circumflex system and functions much like an obtuse marginal AND the patient has LAD disease, code 2 vessel disease.

If there is any confusion about the distribution of the Ramus as it relates to the LAD or Circumflex coronary artery, consult with your surgeon.

The number of diseased vessels may not necessarily match the number of bypass grafts performed.

Left main disease ($\geq 50\%$) is counted as TWO vessels (LAD and Circumflex). For example, left main $\geq 50\%$ and RCA would count as a total of three.

If bypass is performed for an anomalous, kinked or damaged vessel, this vessel is counted as one diseased or abnormal vessel.

Code the number of vessels diseased only for those vessels that have a stenosis greater than or equal to 50%.

SEQ. #: 1175

Long Name: Percent Native Artery Stenosis Known

Short Name: PctStenKnown

Definition: Indicate whether the percent stenosis of native coronary stenosis is known.

Intent/Clarification:

A patient may never have more than three vessel disease. Once a coronary artery is found to be diseased, for the purposes of the STS, the vessel is considered diseased for the remainder of the patient “s life and all subsequent reoperations regardless of previous interventions.

The Coronary section is arranged in a grid format. Each column header has a ““Yes,”/no” field. If any column has a ““Yes,”“answer, at least one vessel below must have documentation. If the medical record has conflicting reports on the vessel name, for example a vessel is described as OM 1 by one provider and the same vessel is referred to as the Ramus by another provider, use the surgeon “s description of the lesion location.

Each Column with a ““Yes,”” response below must have documentation on at least one vessel

Coronary	Native Artery % Stenosis Known: PctStenKnown (1175) <input type="checkbox"/> Yes <input type="checkbox"/> No (If yes↓)	Graft(s) Graft(s) Present: GraftsPrsnt (1180) <input type="checkbox"/> Yes <input type="checkbox"/> No (If yes↓)	Stent(s) Stent(s) Present: StentPrsnt (1185) <input type="checkbox"/> Yes <input type="checkbox"/> No (If yes↓)	Fractional Flow Reserve (FFR) performed: FFRPerf (1190) <input type="checkbox"/> Yes <input type="checkbox"/> No (If yes↓)	Instantaneous wave-free ratio (iFR) performed: IFRPerf (1191) <input type="checkbox"/> Yes <input type="checkbox"/> No (If yes↓)
----------	---	---	--	--	--

SEQ. #: 1180

Long Name: Graft(s) Present

Short Name: GraftsPrsnt

Definition: Indicate whether one or more coronary artery bypass grafts are present prior to this surgery.

Intent/Clarification:

- **Yes** – A previous coronary bypass graft is documented in the medical record.
- **No** – No previous graft documented in the medical record.

SEQ. #: 1185

Long Name: Stent(s) Present

Short Name: StentPrsnt

Definition: Indicate whether one or more intracoronary stents are present prior to this surgery.

Intent/Clarification:

- **Yes** – a previously placed coronary artery stent is documented in the medical record.
- **No** – no previous coronary artery stent documented in the medical record.

SEQ. #: 1190

Long Name: Fractional Flow Reserve (FFR) Performed

Short Name: FFRPerf

Definition: Indicate whether Fractional Flow Reserve (FFR) was performed.

Intent/Clarification:

Fractional flow reserve (FFR) is a technique used in coronary catheterization to measure pressure differences across a coronary artery stenosis (narrowing, usually due to atherosclerosis) to determine the likelihood that the stenosis impedes oxygen delivery to the heart muscle (myocardial ischemia).

Fractional flow reserve is defined as the pressure behind (distal to) a stenosis relative to the pressure before the stenosis. The result is an absolute number; an FFR of 0.80 means that a given stenosis causes a 20% drop in blood pressure. In other words, FFR expresses the maximal flow down a vessel in the presence of a stenosis compared to the maximal flow in the hypothetical absence of the stenosis.

- **Yes** – a vessel that has a Fractional Flow Reserve documented in the medical record. If the value from the FFR is higher than the cardiac catheterization; code from the FFR.
- **No** – a vessel that has no Fractional Flow Reserve documented in the medical record.

SEQ. #: 1191

Long Name: Instantaneous Wave-Free Ration (iFR) Performed

Short Name: IFRPerf

Definition: Indicate whether Instantaneous wave-free ration (iFR) was performed.

Intent/Clarification:

Instantaneous wave-free ratio is performed using high fidelity pressure wires that are passed distal to the coronary stenosis. iFR isolates a specific period in diastole, called the wave-free period, and uses the ratio of distal coronary pressure (Pd) to the pressure observed in the aorta (Pa) over this period. During this wave-free period, the competing forces (waves) that affect coronary flow are quiescent meaning pressure and flow are linearly related as compared to the rest of the cardiac cycle.

When stenoses are flow limiting, Pd and Pa pressures over the wave-free period diverge; a normal ratio is 1.0 and iFR values of below 0.90 suggest flow limitation. iFR can be calculated using dedicated consoles available for medical use and typically uses an average over 5 heart beats but can be performed using a single heartbeat. iFR is measured at rest, without the need for pharmacological vasodilators or stressors and compares well to other invasive and non-invasive markers of ischemia or flow limitation.

- **Yes** – a vessel that has an Instantaneous Wave-Free Ration documented in the medical record.
 - **No** – a vessel that has no Instantaneous Wave-Free Ration documented in the medical record.
-

GENERAL STATEMENT (10/2017): The following descriptive terms and associated percentages can be used to quantify the % stenosis in any coronary artery.

Stenosis Percent / Descriptive Terms		Percentage Coded
Stenosis <50%		
▪ No stenosis/disease noted	▪ Normal	Stenosis <50% is <i>not</i> captured in v5
▪ No significant disease	▪ Patent vessel	
▪ Luminal irregularities	▪ Minimal disease	
▪ Diffuse disease	▪ Mild disease	
Stenosis ≥50%		
▪ Borderline obstructive/obstruction	▪ Borderline disease	50%
▪ Moderate disease	▪ Intermediate disease	
Stenosis ≥70%		
▪ Significant	▪ Obstructive disease	70%
▪ Flow-limiting		
Stenosis ≥ 90%		
▪ Critical	▪ Subtotaled	90%
▪ Severe	▪ Tight	
▪ Occlusive		
Stenosis 100%		
▪ Total occlusion	▪ Chronic Total Occlusion (CTO)	100%
▪ Occluded		

*Code the highest value when a range is document (Example: if 'mild to moderate' is documented, 50% will be coded)

SEQ. #: 1195

Long Name: Percent Stenosis - Left Main

Short Name: PctStenLMain

Definition: Indicate the highest percent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture % stenosis for vessels with documented stenosis ≥50%
 If “Native Artery % Stenosis Known” (field 1175) is marked “Yes,” at least one vessel must have a percent stenosis marked to avoid a missing data flag in the DQR.
 If there is no stenosis or no documentation or mention of a vessel, leave the selection blank.

In instances where multiple lesions are present, enter the single highest percent stenosis noted in that vessel. When ranges are reported, such as 45- 50% for stenosis, **report as the highest percent in range, in this case 50%.**

Stenosis at the ostia of the LAD and circumflex is not considered left main disease for the purpose of Society of Thoracic Surgeons (STS). **Stenosis needs to be in the left main artery.**

If the cath report states 40% disease, but the Intravascular Ultrasound (IVUS) shows 70%, code 70%.

If multiple sources are available, select surgeon “s documentation degree of stenosis. This is the degree of stenosis that the surgeon used to develop the surgical treatment plan.

FAQ September 2017:

Answer: When coding the % stenosis in a native coronary artery, code all the known percentages even if they are less than 50%. Understand that these fields will only be open to be completed when at least one vessel has a stenosis greater than or equal to 50%.

FAQ October 2017: How would bilateral spontaneous coronary dissections from the ostium of both the Left and Right main coronary arteries to the distal end of the coronary tree be coded.

Answer: Code 100% for both LM and RCA.

SEQ. #: 1200

Long Name: Graft Stenosis - Left Main

Short Name: GrftStenLMain

Definition: Indicate the highest percent stenosis in this graft at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of stenosis in a graft if one is present.

If "Graft Present" (field 1180) is marked "Yes," at least one vessel must have a graft stenosis marked to avoid a missing data flag in the DQR.

If there is no stenosis documented code patent. If no documentation of graft stenosis, code not documented.

- Patent
 - Stenosis \geq 50%
 - 100% occlusion
 - Not documented
-
-

SEQ. #: 1205

Long Name: Stent Stenosis - Left Main

Short Name: StntStenLMain

Definition: Indicate the highest percent of stent stenosis at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of in-stent stenosis if present.

If no documentation of in-stent stenosis, leave blank.

If "Stent Present" (field 1185) is marked "Yes," at least one vessel must have a stent present marked to avoid a missing data flag in the DQR.

If no documentation of in-stenosis, code not documented.

- Patent
 - Stenosis \geq 50%
 - Not documented
-
-

SEQ. #: 1210

Long Name: Fractional Flow Reserve (FFR) - Left Main

Short Name: FFRLMain

Definition: Indicate the FFR in this vessel.

Intent/Clarification:

The intent is to capture the pressure difference across a coronary artery when “FFR” (field 1190) is marked ““Yes,”” and has been performed and documented in at least one vessel.

If there is no FFR reported for this vessel, leave blank. Choose the **lowest value** documented in the medical record. A FFR of 0.70 indicates a higher level of stenosis than a FFR of 0.80.

SEQ. #: 1212

Long Name: Instantaneous Wave-Free Ration (iFR) - Left Main

Short Name: IFRLMain

Definition: Indicate the iFR in this vessel at the time of this surgery.

Intent/Clarification:

If there is no iFR reported for this vessel, leave blank. Choose the **lowest value** documented in the medical record. iFR values of below 0.90 suggest flow limitation.

SEQ. #: 1215

Long Name: Percent Stenosis - Proximal LAD

Short Name: PctStenProxLAD

Definition: Indicate the highest percent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture % stenosis for vessels with documented stenosis $\geq 50\%$ If “Native Artery % Stenosis Known” (field 1175) is marked ““Yes,”” at least one vessel must have a percent stenosis marked to avoid a missing data flag in the DQR.

If there is no stenosis or no documentation or mention of a vessel, leave the selection blank

In instances where multiple lesions are present, enter the single highest percent stenosis noted in that vessel. When ranges are reported, such as 45- 50% for stenosis, **report as the highest percent in range, in this case 50%.**

FAQ September 2017:

General Statement When coding the % stenosis in a native coronary artery, code all the known percentages even if they are less than 50%. Understand that these fields will only for completion when at least one vessel has a stenosis greater than or equal to 50%.

SEQ. #: 1220

Long Name: Graft Stenosis - Proximal LAD

Short Name: GrftStenProxLAD

Definition: Indicate the highest percent stenosis in this graft at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of stenosis in a graft if one is present.

If "Graft Present" (field 1180) is marked "Yes," at least one vessel must have a graft stenosis marked to avoid a missing data flag in the DQR.

If no stenosis, or no documentation of graft stenosis, code not documented.

- Patent
- Stenosis \geq 50%
- 100% Occlusion
- Not documented

SEQ. #: 1225

Long Name: Stent Stenosis - Proximal LAD

Short Name: StntStenProxLAD

Definition: Indicate the highest percent of stent stenosis at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of in-stent stenosis is present.

If no documentation of in-stent stenosis, leave blank.

If "Stent Present" (field 1185) is marked "Yes," at least one vessel must have a stent present marked to avoid a missing data flag in the DQR.

If no documentation of in- stenosis, leave blank.

- Patent
- Stenosis \geq 50%
- Not documented

SEQ. #: 1230

Long Name: Fractional Flow Reserve (FFR) - Proximal LAD

Short Name: FFRProxLAD

Definition: Indicate the FFR in this vessel.

Intent/Clarification:

The intent is to capture the pressure difference across a coronary artery when FFR (field 1190) is "Yes," and has been performed and documented in at least one vessel.

If there is no FFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. A FFR of 0.70 indicates a higher level of stenosis than a FFR of 0.80.

SEQ. #: 1232

Long Name: Instantaneous Wave-Free Ratio (iFR) - Proximal LAD

Short Name: IFRProxLAD

Definition: Indicate the iFR in this vessel at the time of this surgery.

Intent/Clarification:

If there is no iFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. iFR values of below 0.90 suggest flow limitation.

SEQ. #: 1235

Long Name: Percent Stenosis - Mid LAD

Short Name: PctStenMidLAD

Definition: Indicate the highest percent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture % stenosis for vessels with documented stenosis $\geq 50\%$

If "Native Artery % Stenosis Known" (1175) is marked "Yes," at least one vessel must have a percent stenosis marked to avoid a missing data flag in the DQR.

If there is no stenosis or no documentation or mention of a vessel, leave the selection blank

In instances where multiple lesions are present, enter the single highest percent stenosis noted in that vessel. When ranges are reported, such as 45- 50% for stenosis, **report as the highest percent in range, in this case 50%.**

FAQ General Statement 2017:

General Statement When coding the % stenosis in a native coronary artery, code all the known percentages even if they are less than 50%. Understand that these fields will only for completion when at least one vessel has a stenosis greater than or equal to 50%.

SEQ. #: 1240

Long Name: Graft Stenosis - Mid LAD

Short Name: GrftStenMidLAD

Definition: Indicate the highest percent stenosis in this graft at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of stenosis in a graft if one is present.

If "Graft Present" (field 1180) is marked "Yes," at least one vessel must have a graft

stenosis marked to avoid a missing data flag in the DQR.
If no stenosis, or no documentation of graft stenosis, leave blank.

- Patent
- Stenosis \geq 50%
- 100% Occlusion
- Not documented

SEQ. #: 1245

Long Name: Stent Stenosis - Mid LAD

Short Name: StntStenMidLAD

Definition: Indicate the highest percent of stent stenosis at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of in-stent stenosis is present.

If no documentation of in-stent stenosis, leave blank.

If “Stent Present” (field 1185) is marked “Yes,” at least one vessel must have a stent present marked to avoid a missing data flag in the DQR.

If no documentation of in- stenosis, leave blank.

- Patent
- Stenosis \geq 50%
- Not documented

SEQ. #: 1250

Long Name: Fractional Flow Reserve (FFR) - Mid LAD

Short Name: FFRMidLAD

Definition: Indicate the FFR in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the pressure difference across a coronary artery when FFR (field 1190) is “Yes,” and has been performed and documented in at least one vessel.

If there is no FFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. A FFR of 0.70 indicates a higher level of stenosis than a FFR of 0.80.

SEQ. #: 1252

Long Name: Instantaneous Wave-Free Ration (iFR) - Mid LAD

Short Name: IFRMidLAD

Definition: Indicate the iFR in this vessel at the time of this surgery.

Intent/Clarification:

If there is no iFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. iFR values of below 0.90 suggest flow limitation.

SEQ. #: 1255

Long Name: Percent Stenosis - Distal LAD

Short Name: PctStenDistLAD

Definition: Indicate the highest percent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture % stenosis for vessels with documented stenosis $\geq 50\%$

If "Native Artery % Stenosis Known" (field 1175) is marked "Yes," at least one vessel must have a percent stenosis marked to avoid a missing data flag in the DQR.

If there is no stenosis or no documentation or mention of a vessel, leave the selection blank

In instances where multiple lesions are present, enter the single highest percent stenosis noted in that vessel. When ranges are reported, such as 45- 50% for stenosis, **report as the highest percent in range, in this case 50%.**

FAQ September 2017:

General Statement When coding the % stenosis in a native coronary artery, code all the known percentages even if they are less than 50%. Understand that these fields will only for completion when at least one vessel has a stenosis greater than or equal to 50%.

SEQ. #: 1260

Long Name: Graft Stenosis - Distal LAD

Short Name: GrftStenDistLAD

Definition: Indicate the highest percent stenosis in this graft at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of stenosis in a graft if one is present.

If "Graft Present" (field 1180) is marked "Yes," at least one vessel must have a graft stenosis marked to avoid a missing data flag in the DQR.

If no stenosis, or no documentation of graft stenosis, leave blank.

- Patent
 - Stenosis $\geq 50\%$
 - 100% Occlusion
 - Not documented
-
-

SEQ. #: 1265

Long Name: Stent Stenosis - Distal LAD

Short Name: StntStenDistLAD

Definition: Indicate the highest percent of stent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of in-stent stenosis is present.

If no documentation of in-stent stenosis, leave blank.

If “Stent Present” (field 1185) is marked “Yes,” at least one vessel must have a stent present marked to avoid a missing data flag in the DQR.

If no documentation of in- stenosis, leave blank.

- Patent
- Stenosis \geq 50%
- Not documented

SEQ. #: 1270

Long Name: Fractional Flow Reserve (FFR) - Distal LAD

Short Name: FFRDistLAD

Definition: Indicate the FFR in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the pressure difference across a coronary artery when FFR (field 1190) is “Yes,” and has been performed and documented in at least one vessel.

If there is no FFR reported for this vessel, leave blank. Choose the **lowest** value

documented in the medical record. A FFR of 0.70 indicates a higher level of stenosis than a FFR of 0.80.

SEQ. #: 1272

Long Name: Instantaneous Wave-Free Ration (iFR) - Distal LAD

Short Name: IFRDistLAD

Definition: Indicate the iFR in this vessel at the time of this surgery.

Intent/Clarification:

If there is no iFR reported for this vessel, leave blank. Choose the **lowest** value

documented in the medical record. iFR values of below 0.90 suggest flow limitation.

SEQ. #: 1275

Long Name: Percent Stenosis - Diagonal 1

Short Name: PctStenDiag1

Definition: Indicate the highest percent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture % stenosis for vessels with documented stenosis \geq 50%

If “Native Artery % Stenosis Known” (field 1175) is marked “Yes,” at least one vessel must have a percent stenosis marked to avoid a missing data flag in the DQR.

If there is no stenosis or no documentation or mention of a vessel, leave the selection blank

In instances where multiple lesions are present, enter the single highest percent stenosis noted in that vessel. When ranges are reported, such as 45- 50% for stenosis, **report as the highest percent in range, in this case 50%.**

FAQ September 2017:

General Statement When coding the % stenosis in a native coronary artery, code all the known percentages even if they are less than 50%. Understand that these fields will only for completion when at least one vessel has a stenosis greater than or equal to 50%.

SEQ. #: 1280

Long Name: Graft Stenosis - Diagonal 1

Short Name: GrftStenDiag1

Definition: Indicate the highest percent stenosis in this graft at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of stenosis in a graft if one is present.

If “Graft Present” (field 1180) is marked “Yes,” at least one vessel must have a graft stenosis marked to avoid a missing data flag in the DQR.

If no stenosis, or no documentation of graft stenosis, leave blank.

- Patent
- Stenosis \geq 50%
- 100% Occlusion
- Not documented

SEQ. #: 1285

Long Name: Stent Stenosis - Diagonal 1

Short Name: StntStenDiag1

Definition: Indicate the highest percent of stent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

If no documentation of in-stent stenosis, leave blank.

The intent is to capture the degree of in-stent stenosis is present.

If “Stent Present” (field 1185) is marked “Yes,” at least one vessel must have a stent present marked to avoid a missing data flag in the DQR.

If no documentation of in- stenosis, leave blank.

- Patent
- Stenosis \geq 50%
- Not documented

SEQ. #: 1290

Long Name: Fractional Flow Reserve (FFR) - Diagonal 1

Short Name: FFRDiag1

Definition: Indicate the FFR in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the pressure difference across a coronary artery when FFR (field 1190) is “Yes,” and has been performed and documented in at least one vessel. If there is no FFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. A FFR of 0.70 indicates a higher level of stenosis than a FFR of 0.80.

SEQ. #: 1292

Long Name: Instantaneous Wave-Free Ratio (iFR) - Diagonal 1

Short Name: IFRDiag1

Definition: Indicate the iFR in this vessel at the time of this surgery.

Intent/Clarification:

If there is no iFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. iFR values of below 0.90 suggest flow limitation.

SEQ. #: 1295

Long Name: Percent Stenosis - Diagonal 2

Short Name: PctStenDiag2

Definition: Indicate the highest percent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture % stenosis for vessels with documented stenosis $\geq 50\%$. If “Native Artery % Stenosis Known” (field 1175) is marked “Yes,” at least one vessel must have a percent stenosis marked to avoid a missing data flag in the DQR. If there is no stenosis or no documentation or mention of a vessel, leave the selection blank. In instances where multiple lesions are present, enter the single highest percent stenosis noted in that vessel. When ranges are reported, such as 45- 50% for stenosis, **report as the highest percent in range, in this case 50%.**

FAQ September 2017:

General Statement When coding the % stenosis in a native coronary artery, code all the known percentages even if they are less than 50%. Understand that these fields will

only for completion when at least one vessel has a stenosis greater than or equal to 50%.

SEQ. #: 1300

Long Name: Graft Stenosis - Diagonal 2

Short Name: GrftStenDiag2

Definition: Indicate the highest percent stenosis in this graft at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of stenosis in a graft if one is present.

If “Graft Present” (field 1180) is marked “Yes,” at least one vessel must have a graft stenosis marked to avoid a missing data flag in the DQR.

If no stenosis, or no documentation of graft stenosis, leave blank.

- Patent
 - Stenosis \geq 50%
 - 100% Occlusion
 - Not documented
-
-

SEQ. #: 1305

Long Name: Stent Stenosis - Diagonal 2

Short Name: StntStenDiag2

Definition: Indicate the highest percent of stent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of in-stent stenosis is present.

If no documentation of in-stent stenosis, leave blank.

If “Stent Present” (field 1185) is marked “Yes,” at least one vessel must have a stent present marked to avoid a missing data flag in the DQR.

If no documentation of in- stenosis, leave blank.

- Patent
 - Stenosis \geq 50%
 - Not documented
-
-

SEQ. #: 1310

Long Name: Fractional Flow Reserve (FFR) - Diagonal 2

Short Name: FFRDiag2

Definition: Indicate the FFR in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the pressure difference across a coronary artery when FFR (field 1190) is “Yes,” and has been performed and documented in at least one vessel.

If there is no FFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. A FFR of 0.70 indicates a higher level of stenosis than a FFR of 0.80.

SEQ. #: 1312

Long Name: Instantaneous Wave-Free Ration (iFR) - Diagonal 2

Short Name: IFRDiag2

Definition: Indicate the iFR in this vessel at the time of this surgery.

Intent/Clarification:

If there is no iFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. iFR values of below 0.90 suggest flow limitation.

SEQ. #: 1315

Long Name: Percent Stenosis - Diagonal 3

Short Name: PctStenDiag3

Definition: Indicate the highest percent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture % stenosis for vessels with documented stenosis $\geq 50\%$

If "Native Artery % Stenosis Known" (field 1175) is marked "Yes," at least one vessel must have a percent stenosis marked to avoid a missing data flag in the DQR.

If there is no stenosis or no documentation or mention of a vessel, leave the selection blank

In instances where multiple lesions are present, enter the single highest percent stenosis noted in that vessel. When ranges are reported, such as 45- 50% for stenosis, **report as the highest percent in range, in this case 50%.**

FAQ September 2017:

General Statement When coding the % stenosis in a native coronary artery, code all the known percentages even if they are less than 50%. Understand that these fields will only for completion when at least one vessel has a stenosis greater than or equal to 50%.

SEQ. #: 1320

Long Name: Graft Stenosis - Diagonal 3

Short Name: GrftStenDiag3

Definition: Indicate the highest percent stenosis in this graft at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of stenosis in a graft if one is present.

If “Graft Present” (field 1180) is marked “Yes,” at least one vessel must have a graft stenosis marked to avoid a missing data flag in the DQR.

If no stenosis, or no documentation of graft stenosis, leave blank.

- Patent
 - Stenosis \geq 50%
 - 100% Occlusion
 - Not documented
-
-

SEQ. #: 1325

Long Name: Stent Stenosis - Diagonal 3

Short Name: StntStenDiag3

Definition: Indicate the highest percent of stent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of in-stent stenosis is present.

If no documentation of in-stent stenosis, leave blank.

If “Stent Present” (field 1185) is marked “Yes,” at least one vessel must have a stent present marked to avoid a missing data flag in the DQR.

If no documentation of in-stenosis, leave blank.

- Patent
 - Stenosis \geq 50%
 - Not documented
-
-

SEQ. #: 1330

Long Name: Fractional Flow Reserve (FFR) - Diagonal 3

Short Name: FFRDiag3

Definition: Indicate the FFR in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the pressure difference across a coronary artery when FFR (field 1190) is “Yes,” and has been performed and documented in at least one vessel.

If there is no FFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. A FFR of 0.70 indicates a higher level of stenosis than a FFR of 0.80.

SEQ. #: 1332

Long Name: Instantaneous Wave-Free Ration (iFR) - Diagonal 3

Short Name: IFRDiag3

Definition: Indicate the iFR in this vessel at the time of this surgery.

Intent/Clarification:

If there is no iFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. iFR values of below 0.90 suggest flow limitation.

SEQ. #: 1335

Long Name: Percent Stenosis - Circumflex

Short Name: PctStenCircflx

Definition: Indicate the highest percent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture % stenosis for vessels with documented stenosis $\geq 50\%$. If "Native Artery % Stenosis Known" (field 1175) is marked "Yes," at least one vessel must have a percent stenosis marked to avoid a missing data flag in the DQR.

If there is no stenosis or no documentation or mention of a vessel, leave the selection blank.

In instances where multiple lesions are present, enter the single highest percent stenosis noted in that vessel. When ranges are reported, such as 45- 50% for stenosis, **report as the highest percent in range, in this case 50%.**

FAQ September 2017:

General Statement When coding the % stenosis in a native coronary artery, code all the known percentages even if they are less than 50%. Understand that these fields will only for completion when at least one vessel has a stenosis greater than or equal to 50%.

SEQ. #: 1340

Long Name: Graft Stenosis - Circumflex

Short Name: GrftStenCircflx

Definition: Indicate the highest percent stenosis in this graft at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of stenosis in a graft if one is present.

If "Graft Present" (field 1180) is marked "Yes," at least one vessel must have a graft stenosis marked to avoid a missing data flag in the DQR.

If no stenosis, or no documentation of graft stenosis, leave blank.

- Patent
 - Stenosis $\geq 50\%$
 - 100% Occlusion
 - Not documented
- -----

SEQ. #: 1345

Long Name: Stent Stenosis - Circumflex

Short Name: StntStenCircflx

Definition: Indicate the highest percent of stent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of stenosis in a graft if one is present.

If "Graft Present" (field 1180) is marked "Yes," at least one vessel must have a graft stenosis marked to avoid a missing data flag in the DQR.

If no stenosis, or no documentation of graft stenosis, leave blank.

- Patent
- Stenosis \geq 50%
- 100% Occlusion
- Not documented

SEQ. #: 1350

Long Name: Fractional Flow Reserve (FFR) - Circumflex

Short Name: FFRCircflx

Definition: Indicate the FFR in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the pressure difference across a coronary artery when FFR (field 1190) is "Yes," and has been performed and documented in at least one vessel.

If there is no FFR reported for this vessel, leave blank. Choose the lowest value documented in the medical record. A FFR of 0.70 indicates a higher level of stenosis than a FFR of 0.80.

SEQ. #: 1352

Long Name: Instantaneous Wave-Free Ration (iFR) - Circumflex

Short Name: IFRCircflx

Definition: Indicate the iFR in this vessel at the time of this surgery.

Intent/Clarification:

If there is no iFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. iFR values of below 0.90 suggest flow limitation.

SEQ. #: 1355

Long Name: Percent Stenosis - Obtuse Marginal 1

Short Name: PctStenOM1

Definition: Indicate the highest percent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture % stenosis for vessels with documented stenosis $\geq 50\%$
If "Native Artery % Stenosis Known" (field 1175) is marked "Yes," at least one vessel must have a percent stenosis marked to avoid a missing data flag in the DQR.

If there is no stenosis or no documentation or mention of a vessel, leave the selection blank

In instances where multiple lesions are present, enter the single highest percent stenosis noted in that vessel. When ranges are reported, such as 45- 50% for stenosis, **report as the highest percent in range, in this case 50%.**

FAQ September 2017:

General Statement When coding the % stenosis in a native coronary artery, code all the known percentages even if they are less than 50%. Understand that these fields will only for completion when at least one vessel has a stenosis greater than or equal to 50%.

SEQ. #: 1360

Long Name: Graft Stenosis - Obtuse Marginal 1

Short Name: GrftStenOM1

Definition: Indicate the highest percent stenosis in this graft at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of stenosis in a graft if one is present.

If "Graft Present" (field 1180) is marked "Yes," at least one vessel must have a graft stenosis marked to avoid a missing data flag in the DQR.

If no stenosis, or no documentation of graft stenosis, leave blank.

- Patent
- Stenosis $\geq 50\%$
- 100% Occlusion
- Not documented

SEQ. #: 1365

Long Name: Stent Stenosis - Obtuse Marginal 1

Short Name: StntStenOM1

Definition: Indicate the highest percent of stent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of in-stent stenosis is present.

If no documentation of in-stent stenosis, leave blank.

If "Stent Present" (field 1185) is marked "Yes," at least one vessel must have a stent present marked to avoid a missing data flag in the DQR.

If no documentation of in- stenosis, leave blank.

- Patent
- Stenosis \geq 50%
- Not documented

SEQ. #: 1370

Long Name: Fractional Flow Reserve (FFR) - Obtuse Marginal 1

Short Name: FFROM1

Definition: Indicate the FFR in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the pressure difference across a coronary artery when FFR (field 1190) is “Yes,” and has been performed and documented in at least one vessel. If there is no FFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. A FFR of 0.70 indicates a higher level of stenosis than a FFR of 0.80.

SEQ. #: 1372

Long Name: Instantaneous Wave-Free Ration (iFR) - Obtuse Marginal 1

Short Name: IFROM1

Definition: Indicate the iFR in this vessel at the time of this surgery.

Intent/Clarification:

If there is no iFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. iFR values of below 0.90 suggest flow limitation.

SEQ. #: 1375

Long Name: Percent Stenosis - Obtuse Marginal 2

Short Name: PctStenOM2

Definition: Indicate the highest percent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture % stenosis for vessels with documented stenosis \geq 50% If “Native Artery % Stenosis Known” (field 1175) is marked “Yes,” at least one vessel must have a percent stenosis marked to avoid a missing data flag in the DQR.

If there is no stenosis or no documentation or mention of a vessel, leave the selection blank

In instances where multiple lesions are present, enter the single highest percent stenosis noted in that vessel. When ranges are reported, such as 45- 50% for stenosis, **report as the highest percent in range, in this case 50%.**

FAQ September/2017:

General Statement When coding the % stenosis in a native coronary artery, code all the known percentages even if they are less than 50%. Understand that these fields will only for completion when at least one vessel has a stenosis greater than or equal to 50%.

SEQ. #: 1380

Long Name: Graft Stenosis - Obtuse Marginal 2

Short Name: GrftStenOM2

Definition: Indicate the highest percent stenosis in this graft at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of stenosis in a graft if one is present.

If “Graft Present” (field 1180) is marked “Yes,” at least one vessel must have a graft stenosis marked to avoid a missing data flag in the DQR.

If no stenosis, or no documentation of graft stenosis, leave blank.

- Patent
 - Stenosis \geq 50%
 - 100% Occlusion
 - Not documented
-
-

SEQ. #: 1385

Long Name: Stent Stenosis - Obtuse Marginal 2

Short Name: StntStenOM2

Definition: Indicate the highest percent of stent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of in-stent stenosis is present.

If no documentation of in-stent stenosis, leave blank.

If “Stent Present” (field 1185) is marked “Yes,” at least one vessel must have a stent present marked to avoid a missing data flag in the DQR.

If no documentation of in- stenosis, leave blank.

- Patent
 - Stenosis \geq 50%
 - Not documented
-
-

SEQ. #: 1390

Long Name: Fractional Flow Reserve (FFR) - Obtuse Marginal 2

Short Name: FFROM2

Definition: Indicate the FFR in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the pressure difference across a coronary artery when FFR (field 1190) is “Yes,” and has been performed and documented in at least one vessel. If there is no FFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. A FFR of 0.70 indicates a higher level of stenosis than a FFR of 0.80.

SEQ. #: 1392

Long Name: Instantaneous Wave-Free Ration (iFR) - Obtuse Marginal 2

Short Name: IFRM2

Definition: Indicate the iFR in this vessel at the time of this surgery.

Intent/Clarification:

If there is no iFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. iFR values of below 0.90 suggest flow limitation.

SEQ. #: 1395

Long Name: Percent Stenosis - Obtuse Marginal 3

Short Name: PctStenOM3

Definition: Indicate the highest percent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture % stenosis for vessels with documented stenosis $\geq 50\%$. If “Native Artery % Stenosis Known” (field 1175) is marked “Yes,” at least one vessel must have a percent stenosis marked to avoid a missing data flag in the DQR. If there is no stenosis or no documentation or mention of a vessel, leave the selection blank. In instances where multiple lesions are present, enter the single highest percent stenosis noted in that vessel. When ranges are reported, such as 45- 50% for stenosis, **report as the highest percent in range, in this case 50%.**

FAQ September 2017:

General Statement When coding the % stenosis in a native coronary artery, code all the known percentages even if they are less than 50%. Understand that these fields will only for completion when at least one vessel has a stenosis greater than or equal to 50%.

SEQ. #: 1400

Long Name: Graft Stenosis - Obtuse Marginal 3

Short Name: GrftStenOM3

Definition: Indicate the highest percent stenosis in this graft at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of stenosis in a graft if one is present.

If “Graft Present” (field 1180) is marked “Yes,” at least one vessel must have a graft stenosis marked to avoid a missing data flag in the DQR.

If no stenosis, or no documentation of graft stenosis, leave blank.

- Patent
- Stenosis \geq 50%
- 100% Occlusion
- Not documented

SEQ. #: 1405

Long Name: Stent Stenosis - Obtuse Marginal 3

Short Name: StntStenOM3

Definition: Indicate the highest percent of stent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

If no documentation of in-stent stenosis, leave blank.

The intent is to capture the degree of in-stent stenosis is present.

If “Stent Present” (field 1185) is marked “Yes,” at least one vessel must have a stent present marked to avoid a missing data flag in the DQR.

If no documentation of in- stenosis, leave blank.

- Patent
- Stenosis \geq 50%
- Not documented

SEQ. #: 1410

Long Name: Fractional Flow Reserve (FFR) - Obtuse Marginal 3

Short Name: FFROM3

Definition: Indicate the FFR in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the pressure difference across a coronary artery when FFR (field 1190) is “Yes,” and has been performed and documented in at least one vessel.

If there is no FFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. A FFR of 0.70 indicates a higher level of stenosis than a FFR of 0.80.

SEQ. #: 1412

Long Name: Instantaneous Wave-Free Ration (iFR) - Obtuse Marginal 3

Short Name: IFROM3

Definition: Indicate the iFR in this vessel at the time of this surgery.

Intent/Clarification:

If there is no iFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. iFR values of below 0.90 suggest flow limitation.

SEQ. #: 1415

Long Name: Percent Stenosis - Ramus

Short Name: PctStenRamus

Definition: Indicate the highest percent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture % stenosis for vessels with documented stenosis $\geq 50\%$. If "Native Artery % Stenosis Known" (field 1175) is marked "Yes," at least one vessel must have a percent stenosis marked to avoid a missing data flag in the DQR.

If there is no stenosis or no documentation or mention of a vessel, leave the selection blank.

In instances where multiple lesions are present, enter the single highest percent stenosis noted in that vessel. When ranges are reported, such as 45- 50% for stenosis, **report as the highest percent in range, in this case 50%.**

FAQ September 2017:

General Statement When coding the % stenosis in a native coronary artery, code all the known percentages even if they are less than 50%. Understand that these fields will only for completion when at least one vessel has a stenosis greater than or equal to 50%.

SEQ. #: 1420

Long Name: Graft Stenosis - Ramus

Short Name: GrftStenRamus

Definition: Indicate the highest percent stenosis in this graft at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of stenosis in a graft if one is present.

If "Graft Present" (field 1180) is marked "Yes," at least one vessel must have a graft stenosis marked to avoid a missing data flag in the DQR.

If no stenosis, or no documentation of graft stenosis, leave blank.

- Patent
- Stenosis $\geq 50\%$
- 100% Occlusion
- Not documented

SEQ. #: 1425

Long Name: Stent Stenosis - Ramus

Short Name: StntStenRamus

Definition: Indicate the highest percent of stent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of in-stent stenosis is present.

If no documentation of in-stent stenosis, leave blank.

If "Stent Present" (field 1185) is marked "Yes," at least one vessel must have a stent present marked to avoid a missing data flag in the DQR.

If no documentation of in- stenosis, leave blank.

- Patent
- Stenosis \geq 50%
- Not documented

SEQ. #: 1430

Long Name: Fractional Flow Reserve (FFR) - Ramus

Short Name: FFRRamus

Definition: Indicate the FFR in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the pressure difference across a coronary artery when FFR (field 1190) is "Yes," and has been performed and documented in at least one vessel.

If there is no FFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. A FFR of 0.70 indicates a higher level of stenosis than a FFR of 0.80.

SEQ. #: 1432

Long Name: Instantaneous Wave-Free Ration (iFR) - Ramus

Short Name: IFRRamus

Definition: Indicate the iFR in this vessel at the time of this surgery.

Intent/Clarification:

If there is no iFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. iFR values of below 0.90 suggest flow limitation.

If there is no iFR reported for this vessel, leave blank. Choose the lowest value documented in the medical record. iFR values of below 0.90 suggest flow limitation.

SEQ. #: 1435

Long Name: Percent Stenosis - RCA

Short Name: PctStenRCA

Definition: Indicate the highest percent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture % stenosis for vessels with documented stenosis $\geq 50\%$

If “Native Artery % Stenosis Known” (field 1175) is marked “Yes,” at least one vessel must have a percent stenosis marked to avoid a missing data flag in the DQR.

If there is no stenosis or no documentation or mention of a vessel, leave the selection blank.

In instances where multiple lesions are present, enter the single highest percent stenosis noted in that vessel. When ranges are reported, such as 45- 50% for stenosis, **report as the highest percent in range, in this case 50%.**

FAQ September 2017:

General Statement When coding the % stenosis in a native coronary artery, code all the known percentages even if they are less than 50%. Understand that these fields will only for completion when at least one vessel has a stenosis greater than or equal to 50%.

FAQ October 2017: How would bilateral spontaneous coronary dissections from the ostium of both the Left and Right main coronary arteries to the distal end of the coronary tree be coded.

Answer: Code 100% for both LM and RCA.

SEQ. #: 1440

Long Name: Graft Stenosis - RCA

Short Name: GrftStenRCA

Definition: Indicate the highest percent stenosis in this graft at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of stenosis in a graft if one is present.

If “Graft Present” (field 1180) is marked “Yes,” at least one vessel must have a graft stenosis marked to avoid a missing data flag in the DQR.

If no stenosis, or no documentation of graft stenosis, leave blank.

- Patent
- Stenosis $\geq 50\%$
- 100% Occlusion
- Not documented

SEQ. #: 1445

Long Name: Stent Stenosis - RCA

Short Name: StntStenRCA

Definition: Indicate the highest percent of stent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

If no documentation of in-stent stenosis, leave blank.

The intent is to capture the degree of in-stent stenosis is present.

If "Stent Present" (field 1185) is marked "Yes," at least one vessel must have a stent present marked to avoid a missing data flag in the DQR.

If no documentation of in- stenosis, leave blank.

- Patent
- Stenosis \geq 50%
- Not documented

SEQ. #: 1450

Long Name: Fractional Flow Reserve (FFR) - RCA

Short Name: FFRRCA

Definition: Indicate the FFR in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the pressure difference across a coronary artery when FFR (field 1190) is "Yes," and has been performed and documented in at least one vessel.

If there is no FFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. A FFR of 0.70 indicates a higher level of stenosis than a FFR of 0.80.

SEQ. #: 1452

Long Name: Instantaneous Wave-Free Ration (iFR) - RCA

Short Name: IFRRCA

Definition: Indicate the iFR in this vessel at the time of this surgery.

Intent/Clarification:

If there is no iFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. iFR values of below 0.90 suggest flow limitation.

SEQ. #: 1455

Long Name: Percent Stenosis - Acute Marginal (AM)

Short Name: PctStenAM

Definition: Indicate the highest percent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture % stenosis for vessels with documented stenosis $\geq 50\%$
If "Native Artery % Stenosis Known" (field 1175) is marked "Yes," at least one vessel must have a percent stenosis marked to avoid a missing data flag in the DQR.

If there is no stenosis or no documentation or mention of a vessel, leave the selection blank

In instances where multiple lesions are present, enter the single highest percent stenosis noted in that vessel. When ranges are reported, such as 45- 50% for stenosis, **report as the highest percent in range, in this case 50%.**

FAQ September 2017:

General Statement When coding the % stenosis in a native coronary artery, code all the known percentages even if they are less than 50%. Understand that these fields will only for completion when at least one vessel has a stenosis greater than or equal to 50%.

SEQ. #: 1460

Long Name: Graft Stenosis - Acute Marginal (AM)

Short Name: GrftStenAM

Definition: Indicate the highest percent stenosis in this graft at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of stenosis in a graft if one is present.

If "Graft Present" (field 1180) is marked "Yes," at least one vessel must have a graft stenosis marked to avoid a missing data flag in the DQR.

If no stenosis, or no documentation of graft stenosis, leave blank.

- Patent
 - Stenosis $\geq 50\%$
 - 100% Occlusion
 - Not documented
- -----

SEQ. #: 1465

Long Name: Stent Stenosis - Acute Marginal (AM)

Short Name: StntStenAM

Definition: Indicate the highest percent of stent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of in-stent stenosis is present.

If no documentation of in-stent stenosis, leave blank.

If "Stent Present" (field 1185) is marked "Yes," at least one vessel must have a stent

present marked to avoid a missing data flag in the DQR.
If no documentation of in- stenosis, leave blank.

- Patent
- Stenosis \geq 50%
- Not documented

SEQ. #: 1470

Long Name: Fractional Flow Reserve (FFR) - Acute Marginal (AM)

Short Name: FFRAM

Definition: Indicate the FFR in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the pressure difference across a coronary artery when FFR (field 1190) is “Yes,” and has been performed and documented in at least one vessel. If there is no FFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. A FFR of 0.70 indicates a higher level of stenosis than a FFR of 0.80.

SEQ. #: 1472

Long Name: Instantaneous Wave-Free Ration (iFR) - Acute Marginal (AM)

Short Name: IFRAM

Definition: Indicate the iFR in this vessel at the time of this surgery.

Intent/Clarification:

Choose the **lowest** value documented in the medical record. iFR values of below 0.90 suggest flow limitation. If there is no iFR reported for this vessel, leave blank.

SEQ. #: 1475

Long Name: Percent Stenosis - Posterior Descending (PDA)

Short Name: PctStenPDA

Definition: Indicate the highest percent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture % stenosis for vessels with documented stenosis \geq 50% If “Native Artery % Stenosis Known” (field 1175) is marked “Yes,” at least one vessel must have a percent stenosis marked to avoid a missing data flag in the DQR. If there is no stenosis or no documentation or mention of a vessel, leave the selection blank

In instances where multiple lesions are present, enter the single highest percent stenosis noted in that vessel. When ranges are reported, such as 45- 50% for stenosis, **report as the highest percent in range, in this case 50%.**

FAQ September 2017:

General Statement When coding the % stenosis in a native coronary artery, code all the known percentages even if they are less than 50%. Understand that these fields will only for completion when at least one vessel has a stenosis greater than or equal to 50%.

SEQ. #: 1480

Long Name: Graft Stenosis - Posterior Descending (PDA)

Short Name: GraftStenPDA

Definition: Indicate the highest percent stenosis in this graft at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of stenosis in a graft if one is present.

If "Graft Present" (field 1180) is marked "Yes," at least one vessel must have a graft stenosis marked to avoid a missing data flag in the DQR.

If no stenosis, or no documentation of graft stenosis, leave blank.

- Patent
- Stenosis \geq 50%
- 100% Occlusion
- Not documented

SEQ. #: 1485

Long Name: Stent Stenosis - Posterior Descending (PDA)

Short Name: StntStenPDA

Definition: Indicate the highest percent of stent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

If no documentation of in-stent stenosis, leave blank.

The intent is to capture the degree of in-stent stenosis is present.

If "Stent Present" (field 1185) is marked "Yes," at least one vessel must have a stent present marked to avoid a missing data flag in the DQR.

If no documentation of in- stenosis, leave blank.

- Patent
 - Stenosis \geq 50%
 - Not documented
-
-

SEQ. #: 1490

Long Name: Fractional Flow Reserve (FFR) - Posterior Descending (PDA)

Short Name: FFRPDA

Definition: Indicate the FFR in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the pressure difference across a coronary artery when FFR (field 1190) is “Yes,” and has been performed and documented in at least one vessel. If there is no FFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. A FFR of 0.70 indicates a higher level of stenosis than a FFR of 0.80.

SEQ. #: 1492

Long Name: Instantaneous Wave-Free Ration (iFR) - Posterior Descending (PDA)

Short Name: IFRPDA

Definition: Indicate the iFR in this vessel at the time of this surgery.

Intent/Clarification:

If there is no iFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. iFR values of below 0.90 suggest flow limitation.

SEQ. #: 1495

Long Name: Percent Stenosis - Posterolateral (PLB)

Short Name: PctStenPLB

Definition: Indicate the highest percent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture % stenosis for vessels with documented stenosis $\geq 50\%$. If “Native Artery % Stenosis Known” (field 1175) is marked “Yes,” at least one vessel must have a percent stenosis marked to avoid a missing data flag in the DQR. If there is no stenosis or no documentation or mention of a vessel, leave the selection blank. In instances where multiple lesions are present, enter the single highest percent stenosis noted in that vessel. When ranges are reported, such as 45- 50% for stenosis, **report as the highest percent in range, in this case 50%.**

FAQ September 2017:

General Statement When coding the % stenosis in a native coronary artery, code all the known percentages even if they are less than 50%. Understand that these fields will only for completion when at least one vessel has a stenosis greater than or equal to 50%.

SEQ. #: 1500

Long Name: Graft Stenosis - Posterolateral (PLB)

Short Name: GrftStenPLB

Definition: Indicate the highest percent stenosis in this graft at the time of this surgery.

Intent/Clarification:

The intent is to capture the degree of stenosis in a graft if one is present.

If "Graft Present" (field 1180) is marked "Yes," at least one vessel must have a graft stenosis marked to avoid a missing data flag in the DQR.

If no stenosis, or no documentation of graft stenosis, leave blank.

- Patent
- Stenosis \geq 50%
- 100% Occlusion
- Not documented

SEQ. #: 1505

Long Name: Stent Stenosis - Posterolateral (PLB)

Short Name: StntStenPLB

Definition: Indicate the highest percent of stent stenosis in this vessel at the time of this surgery.

Intent/Clarification:

If no documentation of in-stent stenosis, leave blank.

The intent is to capture the degree of in-stent stenosis is present.

If "Stent Present" (field 1185) is marked "Yes," at least one vessel must have a stent present marked to avoid a missing data flag in the DQR.

If no documentation of in- stenosis, leave blank.

- Patent
- Stenosis \geq 50%
- Not documented

SEQ. #: 1510

Long Name: Fractional Flow Reserve (FFR) - Posterolateral (PLB)

Short Name: FFRPLB

Definition: Indicate the FFR in this vessel at the time of this surgery.

Intent/Clarification:

The intent is to capture the pressure difference across a coronary artery when FFR (field 1190) is "Yes," and has been performed and documented in at least one vessel.

If there is no FFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. A FFR of 0.70 indicates a higher level of stenosis than a FFR of 0.80.

SEQ. #: 1512

Long Name: Instantaneous Wave-Free Ratio (iFR) - Posterolateral (PLB)

Short Name: IFRPLB

Definition: Indicate the iFR in this vessel at the time of this surgery.

Intent/Clarification:

If there is no iFR reported for this vessel, leave blank. Choose the **lowest** value documented in the medical record. iFR values of below 0.90 suggest flow limitation.

SEQ. #: 1515

Long Name: Syntax Score Known

Short Name: SyntaxScrKnown

Definition: Indicate whether a syntax score is known.

Intent/Clarification:

The SYNTAX score is an angiographic grading tool to determine the complexity of coronary artery disease. It is not used routinely at all sites.

- Yes - a syntax score is documented
 - No - no syntax score is documented
-
-

SEQ. #: 1520

Long Name: Syntax Score

Short Name: SyntaxScr

Definition: Indicate syntax score documented prior to this surgery.

Intent/Clarification:

The SYNTAX score is an angiographic grading tool to determine the complexity of coronary artery disease. The SYNTAX score is the sum of the points assigned to each individual lesion identified in the coronary tree with >50% diameter narrowing in vessels > 1.5mm diameter.

Each segment is given a score of 1 or 2 based on the presence of disease and this score is then weighted based on a chart, with values ranging from 3.5 for the proximal left anterior descending artery (LAD) to 5.0 for left main, and 0.5 for smaller branches. The percent diameter stenosis is not a consideration in the SYNTAX score, only the presence of a stenosis from 50–99% diameter, <50% diameter narrowing or the total occlusion.

The SYNTAX score is a useful differentiator for the outcome of patients undergoing three-vessel PCI. The patients with the highest scores have the highest risk and the lowest scores, the lowest risk. The high scores indicate complex conditions and represent greatest risks to patients undergoing PCI. High scores have the worst prognosis for revascularization with PCI compared to coronary artery bypass graft

surgery (CABG).

When the Syntax score is reported in a range, code the highest value.

Reference: Sianos G, Morel MA, Kappetein AP, et al. The SYNTAX score: an angiographic tool grading the complexity of CAD. EuroInterv 2005; 1: 219-227

- Normal
- Abnormal
- Unavailable

SEQ. #: 1525

Long Name: Stress Test Performed

Short Name: StressTst

Definition: Indicate whether a stress test was performed prior to this surgery.

Intent/Clarification:

Indicate whether a stress test was performed within 6 months prior to this surgery. Types of stress tests include the following:

Standard Exercise Stress Test without imaging:

Treadmill Exercise Stress EKG

Stress Echocardiogram

Exercise Stress Echo with Doppler
Exercise Echo with Doppler
Pharmacologic Stress Echo with Doppler
Exercise Echo
Exercise Echo with Color Flow Doppler
Exercise Echo with Spectral Color Flow

Stress Testing with SPECT MPI

Nuclear Medicine Studies
Cardiac Scan - Infarct
Myocardial Perfusion - Rest/Stress
Myocardial Perfusion - Rest/Spect
Myocardial Perfusion - Rest/Stress/Spect
Myocardial Perfusion - Rest or Stress
PET Studies Heart, N-13 Blood Flow, Rest
Heart, N-13 Blood Flow, Stress
Myocardial Viability with Nuclear Perfusion

Stress Testing with CMR

MRI Studies
CMRI Dobutamine Stress
CMRI Adenosine Stress and Perfusion
CMRI Exercise Stress
CMRI Stress plus Flow Velocities with infusion
CMRI Stress plus Flow Velocities without infusion

SEQ. #: 1531

Long Name: Stress Test Result

Short Name: StrsTstRes

Definition: Indicate the results of the stress test.

Intent/Clarification:

- **Negative:** A stress test is negative when the electrocardiogram (ECG) is normal or not suggestive of ischemia. ECGs are not suggestive of ischemia when < 1 mm of horizontal or down sloping ST segment depression or elevation for ≥ 60 -80 milliseconds after the end of the QRS complex, either during or after exercise.
 - **Positive:** A stress test is positive when the electrocardiogram (ECG) suggests ischemia. ECGs suggestive of ischemia can be described as having ≥ 1 mm of horizontal or down sloping ST-segment depression or elevation for ≥ 60 -80 milliseconds after the end of the QRS complex, either during or after exercise. It is also be suggestive of ischemia if the patient had symptoms of ischemia (i.e. chest pain), arrhythmias, and/or a fall in blood pressure during or immediately after the procedure. If more than one study was performed with conflicting results and one study suggested coronary artery disease, code "Yes".
 - **Not documented**
-
-

SEQ. #: 1540

Long Name: Hemo Data-EF Done

Short Name: HDEFD

Definition: Indicate whether the Ejection Fraction was measured prior to the induction of anesthesia.

Intent/Clarification:

Some patients may not have had an LV Gram performed during cardiac catheterization due to existing clinical conditions. Ejection fraction (EF) and hemodynamic pressures may be obtained from other sources other than coronary angiogram, such as echo, or MUGA.

Because anesthesia can alter the values to be collected, do not collect data from intra-operative transesophageal echo (TEE) after the induction of anesthesia, unless you have no other source to collect the information.

Time Frame: Do not use results more than 6 months prior to this operation.

SEQ. #: 1545

Long Name: Hemo Data-EF

Short Name: HDEF

Definition: Indicate the percentage of the blood emptied from the left ventricle at the end of the contraction. Use the most recent determination prior to the surgical intervention documented on a diagnostic report.

Enter a percentage in the range of 1 - 99. If a qualitative description is reported, code the mean value for that range; i.e., normal (50-70%) is coded as 60%.

- Hyperdynamic: >70% (code 71%)
- Normal: 50%–70% (midpoint 60%)
- Mild dysfunction: 40%–49% (midpoint 45%)
- Moderate dysfunction: 30%–39% (midpoint 35%)
- Severe dysfunction: <30% (code 29%)

Note: If no diagnostic report is in the medical record, a value documented in the medical record is acceptable.

ACCF/AHA 2013

Intent/Clarification:

Use the most recent determination **prior to the induction** of anesthesia documented on a diagnostic report, regardless of the diagnostic procedure to obtain it.

If no diagnostic report specifying an ejection fraction (EF) is in the medical record, a value documented in the progress record is acceptable.

If there is no documentation of a pre-op EF, then it is acceptable to code the EF from the intra-op TEE prior to **incision**.

Use the surgeon's documentation if more than one value is reported as this was likely used to plan operative care.

Time Frame: Collect the last value closest to incision, not greater than **6 months**.

SEQ. #: 1555

Long Name: Hemo Data-Dimensions Available

Short Name: DimAvail

Definition: Indicate whether intracardiac dimensions are available.

Intent/Clarification:

Time Frame: Collect the last value closest to incision, not greater than **6 months** prior.

SEQ. #: 1560

Long Name: Hemo Data-LV End Systolic Dimension

Short Name: LVSD

Definition: Indicate LV End -Systolic Dimension in mm.

LV end systolic dimension is the same as left ventricular internal dimension in end systole (LVIDs)

Intent/Clarification:

During systole the left ventricle contracts pumping blood through the body. During

diastole the left ventricle relaxes and fills with blood again. The systolic dimension of the left ventricle demonstrates ventricular emptying and when compared to the end diastolic dimension, left ventricular performance is calculated.

SEQ. #: 1565

Long Name: Hemo Data-LV End-Diastolic Dimension

Short Name: LVEDD

Definition: Indicate the Left Ventricular End-Diastolic Dimension in mm. LV end diastolic dimension is the same as left ventricular internal dimension in end diastole (LVIDs)

Intent/Clarification:

During systole the ventricles contract pumping blood through the body. During diastole the ventricles relax and fill with blood again. The end-diastolic dimension of the left ventricle demonstrates ventricular filling and when compared to the end systolic dimension, left ventricular performance is calculated.

SEQ. #: 1570

Long Name: Hemo-PA Systolic Pressure Measured

Short Name: PASYSMeas

Definition: Indicate whether the PA systolic pressure was measured prior to ~~incision~~ **induction**.

Intent/Clarification:

Elevated pulmonary artery pressures are indicative of pulmonary hypertension, mitral valve disease and other pulmonary/cardiac diseases. Normal mean pulmonary artery pressure readings are between 9-17mm of pressure. If there are no PA pressures recorded or available from heart Cath –one may use PA pressure values from Swan Ganz Catheter inserted for surgery prior to induction of anesthesia.

SEQ. #: 1575

Long Name: Hemo-PA Systolic Pressure

Short Name: PASYS

Definition: Capture highest PA systolic pressure recorded prior to ~~incision~~ **induction**.

Intent/Clarification:

Elevated pulmonary artery pressures are indicative of pulmonary hypertension, mitral valve disease and other pulmonary/cardiac diseases. Normal mean pulmonary artery pressure readings are between 9-17mm of pressure.

If there are no PA pressures recorded or available from heart Cath one may use PA pressure values from Swan Ganz Catheter inserted for surgery prior to induction.

If more than one preoperative measurement is available, choose the **HIGHEST** PA systolic pressure recorded before induction.

If PA systolic pressure is not available it is acceptable to code the peak RV systolic pressure (RVSP). RVSP and PA systolic pressures will be the same as long as there is no pulmonary valve disease or outflow obstruction.

If more than one preoperative measurement is available, choose the **HIGHEST** PA systolic pressure recorded before induction.

If there is a preoperative echo, use those values **UNLESS** the diagnostic information from the TEE changes the procedure performed.

If there is no preop information, you may use the pre-incision intraoperative TEE.

FAQ August 2017: Please clarify, should the value be taken prior to induction or prior to incision?

Answer: The PA systolic value should be taken prior to induction of anesthesia.

SEQ. #: 1590

Long Name: VD-Insuff-Aortic

Short Name: VDInsufA

Definition: Indicate whether there is evidence of Aortic valve insufficiency/regurgitation. Enter the degree of insufficiency reported closest to incision and no more than 6 months prior to surgery.

Intent/Clarification:

Regurgitation/insufficiency is incompetence of the aortic valve or any of its valvular apparatus which allows diastolic blood flow to flow back into the left ventricular chamber. This may be a chronic or an acute condition.

Time Frame: Collect the last value closest to incision, not greater than **6 months** prior. Choose the **highest** level of valve dysfunction when there are differences in interpretation of the most recent study. Capture even if patient is not scheduled for valve repair and/or replacement when available.

- None
- Trivial/Trace
- Mild
- Moderate
- Severe
- Not documented

SEQ. #: 1591

Long Name: VD-Aortic Valve Eccentric Jet

Short Name: VDAVEccJet

Definition: Indicate whether aortic valve regurgitation is an eccentric jet.

Intent/Clarification:

- Yes
- No
- Not Documented

SEQ. #: 1595

Long Name: VD-Aortic

Short Name: VDAort

Definition: Indicate whether Aortic Valve disease is present.

Intent/Clarification:

Aortic valvular disease can be congenital or acquired and cause stenosis, regurgitation or both.

The valve should be coded as being diseased if there is mild, moderate or severe insufficiency.

SEQ. #: 1600

Long Name: VD-Stenosis-Aortic

Short Name: VDStenA

Definition: Indicate whether Aortic Stenosis is present.

Intent/Clarification:

The aortic valve controls the direction of blood flow from the left ventricle to the aorta. When in good working order, the aortic valve does not impede the flow of blood between these two spaces. Under some circumstances, the aortic valve becomes narrower than normal, impeding the flow of blood. This is known as aortic valve stenosis or aortic stenosis, often abbreviated as A.S.

AS is described as trace, mild, moderate or severe. Aortic valve stenosis may be caused by aging (leaflets become calcified, thick and stiff), birth defects (congenital bicuspid (2) leaflets) or other disease processes like rheumatic fever.

Capture any degree of aortic valve stenosis present, even if the patient is not scheduled for valve replacement, record if available.

- Yes
- No

Time Frame: Collect the last value closest to incision, not greater than **6 months**.

SEQ. #: 1605

Long Name: VD-Aortic Hemodynamic Data Available

Short Name: AoHemoDatAvail

Definition: Indicate whether aortic valve hemodynamic measurements are available.

Intent/Clarification:

Collect the last value closest to incision, not greater than **6 months**.

SEQ. #: 1610

Long Name: VD-Smallest Aortic Valve Area

Short Name: VDAoVA

Definition: Indicate the smallest documented aortic valve area (in cm squared).

Intent/Clarification:

The normal adult aortic valve opening is 3.0-4.0 (cm²). Aortic stenosis becomes hemodynamically significant when the area decreases to less than 2(cm²), as the systolic flow is impeded across the valve. If more than one aortic valve area is reported, choose the **smallest**.

SEQ. #: 1615

Long Name: VD-Aortic Gradient-Highest Mean

Short Name: VDGradA

Definition: Indicate the highest documented MEAN gradient (in mmHg) across the aortic valve.

Intent/Clarification:

When the aortic valve becomes stenotic, it causes a pressure gradient between the left ventricle (LV) and the aorta. The more constricted the valve, the higher the gradient between the LV and the aorta. For example, if the gradient is 20 mmHg, at peak systole, while the LV generates a pressure of 140 mmHg, the pressure that is transmitted to the aorta would only be 120 mmHg. A blood pressure cuff would measure a normal systolic blood pressure; the actual pressure generated by the LV would be considerably higher. In individuals with AS, the left ventricle (LV) has to work harder to overcome the increased afterload caused by the stenotic aortic valve and eject blood out of the LV. The more severe the aortic stenosis, the higher the gradient is between the left ventricular systolic pressures and the aortic systolic pressures.

SEQ. #: 1616

Long Name: VD - Maximum Aortic Jet Velocity (Vmax)

Short Name: VDVMax

Definition: Indicate the maximum aortic jet velocity

Intent/Clarification:

The antegrade systolic velocity across the narrowed aortic valve, or aortic jet velocity, is measured using continuous-wave (CW) Doppler (CWD) ultrasound. Velocity increases as stenosis severity increases.

SEQ. #: 1646

Long Name: VD-Aortic Valve Disease Primary Etiology

Short Name: VDAoPrimEt

Definition: Indicate the primary etiology of aortic valve disease.

Intent/Clarification:

There is no hierarchy, choose the primary etiology documented in the medical record. Primary etiology may also be identified at the time of the surgical procedure.

FAQ October 2017: My surgeon replaced a mitral valve in a patient with a history of Hodgkin's disease who is s/p chemotherapy and radiation therapy. The etiology is described as radiation induced heart disease. How should this be coded?

Answer: This should be coded as mixed etiology and would apply to aortic valve etiology as well. This will be added in the next upgrade. In addition to capturing the etiology, code mediastinal radiation in Seq. No. 495.

SEQ. #: 1647

Long Name: VD-Aortic Valve Disease Sievers Class

Short Name: VDAoSievers

Definition: Indicate the documented Sievers class

Intent/Clarification:

A systematic classification of bicuspid aortic valves: Three major types were identified: type 0 (no raphe), type 1 (one raphe), and type 2 (two raphes), followed by two supplementary characteristics, spatial position and function. These characteristics served to classify and codify the bicuspid aortic valves.

Raphe is a groove, seam or ridge in tissue typically marking the line where two halves fused in the embryo.

SEQ. #: 1680

Long Name: VD-Insuff-Mitral

Short Name: VDInsufM

Definition: Indicate whether there is evidence of Mitral valve insufficiency/regurgitation. Enter the degree of insufficiency reported closest to incision and no more than 6 months prior to surgery.

Intent/Clarification:

Mitral regurgitation/insufficiency may be an acute or chronic condition manifesting itself as increased left heart filling pressures which increase the left ventricular stroke volume

(amount of blood ejected from the Left Vent. with each heart beat). Over time, and depending upon the severity, MR can result in pulmonary edema and systemic volume overload. In chronic MR, Left Ventricular Hypertrophy may result. Mitral prolapse and rheumatic fever are the most common cause of MR. "Moderately severe" should be coded as "Severe".

Collect the last value closest to incision, not greater than **6 months**.

- None
- Trivial/Trace
- Mild
- Moderate
- Severe
- Not documented

SEQ. #: 1681

Long Name: VD-Mitral Valve Eccentric Jet

Short Name: VDMVEccJet

Definition: Indicate whether mitral valve regurgitation is an eccentric jet.

Intent/Clarification:

- Yes
- No
- Not Documented

SEQ. #: 1685

Long Name: VD-Mitral

Short Name: VDMit

Definition: Indicate whether Mitral valve disease is present.

Intent/Clarification:

The mitral valve is made up of the annulus, anterior and posterior leaflets, and chordae, which attach the leaflets to their respective papillary muscles. A normally functioning valve allows blood to flow unimpeded from the left atrium to the left ventricle during diastole and prevents regurgitation during systole. Normal mitral valve function is dependent not only on the integrity of the underlying valvular structure, but on that of the adjacent myocardium as well. Mitral valve disease is the most common form of heart valve disease in the United States, affecting 5 percent of the population and resulting in over 500,000 hospital admissions per year. There are two general forms of mitral valve disease: mitral regurgitation/insufficiency and mitral stenosis.

The valve should be coded as being diseased if there is mild, moderate or severe insufficiency.

- Yes
- No

SEQ. #: 1690

Long Name: VD-Stenosis-Mitral

Short Name: VDStenM

Definition: Indicate whether Mitral Stenosis is present.

Intent/Clarification:

Stenosis is the narrowing of the valve opening. Valve stenosis is most often caused by rheumatic fever, causing the leaflets to become rigid, stiff, and thick and/or fused reducing the amount of blood able to be ejected from the left atria into the left ventricle. Mitral stenosis (MS) causes blood to back up, dilate the left atria and create buildup of fluid in the lungs (congestive heart failure). Atrial fibrillation is a common arrhythmia in patients with MS.

Time Frame: Collect the last value closest to incision, not greater than **6 months**. Capture any degree of stenosis even if patient is not scheduled for valve repair and/or replacement when available.

- Yes
 - No
- -----

SEQ. #: 1695

Long Name: VD-Mitral Hemodynamic Data Available

Short Name: MiHemoDatAvail

Definition: Indicate whether mitral valve hemodynamic measurements are available.

Intent/Clarification:

- Yes
 - No
- -----

SEQ. #: 1700

Long Name: VD-Smallest Mitral Valve Area

Short Name: VDMVA

Definition: Indicate the smallest documented Mitral Valve Area.

Intent/Clarification:

The normal area of the mitral valve orifice is about 4 to 6 (cm²). Under normal conditions, a normal mitral valve will not impede the flow of blood from the left atrium to the left ventricle during (ventricular) diastole, and the pressures in the left atrium and the left ventricle during ventricular diastole will be equal. When the mitral valve area goes below 2.0 (cm²), the valve causes an impediment to the flow of blood into the left ventricle, creating a pressure gradient across the mitral valve.

Document the smallest valve area in square centimeters. If the cardiac Cath indicates a valve area of 2.0 and the echo report indicates 1.8, code 1.8.

SEQ. #: 1705

Long Name: VD-Mitral Gradient-Highest Mean

Short Name: VDGradM

Definition: Indicate the highest documented mean gradient (in mm Hg) across the mitral valve.

Intent/Clarification:

Mitral valve stenosis results from a narrowing of the mitral valve orifice when the valve is open. The high resistance across the stenotic mitral valve causes blood to back up into the left atrium thereby increasing LA pressure. This results in the left atrial (LA) pressure being much greater than left ventricular (LV) pressure during diastolic filling.

The gradient is highest during early diastole when the flow across the valve is highest. Normally, the pressure gradient across the valve is very small (a few mmHg); however, the pressure gradient can become quite high during severe stenosis (10-30 mmHg). If more than one gradient is documented in the record, capture the **HIGHEST** one.

SEQ. #: 1731

Long Name: VD-Mitral Valve Disease Primary Etiology

Short Name: VDMiPrimEt

Definition: Indicate the primary etiology of Mitral valve disease.

Intent/Clarification:

There is no hierarchy, choose the primary etiology as documented in the medical record. Primary etiology may not be identified until the time of the surgical procedure.

FAQ October 2017: My surgeon replaced a mitral valve in a patient with a history of Hodgkin's disease who is s/p chemotherapy and radiation therapy. The etiology is described as radiation induced heart disease. How should this be coded?

Answer: This should be coded as mixed etiology. This will be added in the next upgrade. In addition to capturing the etiology, code mediastinal radiation in Seq. No. 495.

SEQ. #: 1746

Long Name: VD-Mitral Valve Primary Lesion

Short Name: VDMiPrimLes

Definition: Indicate the primary mitral valve lesion.

Intent/Clarification:

There is no hierarchy, choose the primary lesion. Primary lesion may be identified at the time of the surgical procedure.

SEQ. #: 1775

Long Name: VD-Insuff-Tricuspid

Short Name: VDInsufT

Definition: Indicate whether there is evidence of Tricuspid valve insufficiency/regurgitation. Enter the degree of insufficiency reported closest to incision and no more than **6 months** prior to surgery.

Intent/Clarification:

Tricuspid regurgitation/insufficiency creates a backwards flow of blood across the tricuspid valve and causes enlargement of the right atrium and possibly atrial fibrillation. Capture even if patient is not scheduled for valve repair and/or replacement when available.

Time Frame: Collect the last value closest to incision, not greater than **6 months**.

Choose the highest level of valve dysfunction when there are differences in interpretation of the most recent study.

Capture even if patient is not scheduled for valve repair and/or replacement when available.

- None
- Trivial/Trace
- Mild
- Moderate
- Severe
- Not documented

SEQ. #: 1777

Long Name: VD-Tricuspid Annular Measurement Available

Short Name: VDTrAnnMeas

Definition: Indicate whether a tricuspid annular diameter measurement is available.

Intent/Clarification:

Tricuspid regurgitation (TR) occurs mainly from tricuspid annular dilation, which can result from left-sided heart failure from myocardial or valvular causes, right ventricular volume and pressure overload, or dilation of cardiac chambers.

Time Frame: Collect the last value closest to incision, not greater than **6 months**.

- Yes
- No

SEQ. #: 1778

Long Name: VD-Tricuspid Annulus Size (Diameter)

Short Name: VDTrAnnSize

Definition: Indicate tricuspid annular diameter in cm.

Intent/Clarification:

Normal values for Tricuspid annular diameter: 2-4 (cm²)

SEQ. #: 1780

Long Name: VD-Tricuspid

Short Name: VDTTr

Definition: Indicate whether Tricuspid Valve disease is present.

Intent/Clarification:

Tricuspid valve disease refers to abnormal function of the tricuspid valve.

Two types of tricuspid disease include:

Tricuspid regurgitation - the valve is leaky or doesn't close tight enough, causing blood to leak backwards across the valve.

Tricuspid stenosis - the valve leaflets are stiff and do not open widely enough, causing a restriction in the forward flow of blood.

There is no hierarchy, choose the primary etiology as documented in the medical record. Primary etiology may not be identified until the time of the surgical procedure.

Tricuspid disease should be captured if tricuspid insufficiency is mild, moderate or severe.

- Yes
 - No
-
-

SEQ. #: 1785

Long Name: VD-Stenosis-Tricuspid

Short Name: VDStenT

Definition: Indicate whether Tricuspid Stenosis is present.

Intent/Clarification:

The tricuspid valve is the largest of the four valves. Stenosis, over time, may create an enlarged right atrium, reducing the amount of blood flow into the right ventricle; thereby, reducing cardiac output. Prolonged or chronic tricuspid stenosis may cause systemic vascular congestion, manifested primarily in the liver. Capture even if patient is not scheduled for valve repair or replacement.

Time Frame: Collect the last value closest to incision, not greater than **6 months**.

Choose the highest level of valve dysfunction when there are differences in interpretation of the most recent study.

Capture even if patient is not scheduled for valve repair and/or replacement when available.

- Yes
 - No
-
-

SEQ. #: 1811

Long Name: VD-Tricuspid Valve Disease Primary Etiology

Short Name: VDTrPrimEt

Definition: Indicate the primary etiology of tricuspid valve disease.

Intent/Clarification:

There is no hierarchy, choose the primary etiology. Primary lesion may be identified at the time of the surgical procedure.

FAQ October 2017: My surgeon replaced a mitral valve in a patient with a history of Hodgkin's disease who is s/p chemotherapy and radiation therapy. The etiology is described as radiation induced heart disease. How should this be coded?

Answer: This should be coded as mixed etiology and would apply to tricuspid valve etiology. This will be added in the next upgrade. In addition to capturing the etiology, code mediastinal radiation in Seq. No. 495.

SEQ. #: 1820

Long Name: VD-Insuff-Pulmonic

Short Name: VDInsufP

Definition: Indicate whether there is evidence of Pulmonic valve insufficiency/regurgitation. Enter the degree of insufficiency reported closest to incision and no more than **6 months** prior to surgery.

Intent/Clarification:

Most common cause is from chronic pulmonary hypertension (noted by high PA pressures > 30mm Hg). Incompetent pulmonary leaflets allow blood to flow back into the Right Vent. Capture even if patient is not scheduled for valve repair and/or replacement

Time Frame: Collect the last value closest to incision, not greater than **6 months**. Enter the level of valve function associated with the highest risk (ie. worst performance) recorded in the chart. "Moderately severe" should be coded as "Severe".

Choose the **highest** level of valve dysfunction when there are differences in interpretation of the most recent study.

Capture even if patient is not scheduled for valve repair and/or replacement when available.

- None
- Trivial/Trace
- Mild
- Moderate
- Severe
- Not documented

SEQ. #: 1825

Long Name: VD-Pulmonic

Short Name: VDPulm

Definition: Indicate whether Pulmonic Valve disease is present.

Intent/Clarification:

The pulmonary valve is a valve between the heart and the artery that leads to the lungs. If valve regurgitation or insufficiency is present, blood is able to flow from the artery and back into the heart. Pulmonary stenosis reduces blood flow to the lungs and makes the right ventricle work harder. The condition can cause the right sided heart failure.

Pulmonary valve disease mostly occurs as a congenital abnormality but it can also be caused by conditions such as pulmonary hypertension, infective endocarditis or Marfan syndrome.

The valve should be coded as being diseased if there is mild, moderate or severe insufficiency.

- Yes
- No

SEQ. #: 1830**Long Name:** VD-Pulmonic-RVEDD Known**Short Name:** RVEDDKnown**Definition:** Indicate whether the Right Ventricular End-Diastolic Dimension (RVEDD) is available.**Intent/Clarification:**

- Yes
- No

SEQ. #: 1835**Long Name:** VD-Pulmonic-RVEDD Indexed To BSA**Short Name:** RVEDD**Definition:** Indicate (in cm squared) the RVEDD indexed to BSA.**Intent/Clarification:**

RVEDD may be called RVDD.

SEQ. #: 1840**Long Name:** VD-Stenosis-Pulmonic**Short Name:** VDStenP**Definition:** Indicate whether Pulmonic Stenosis is present.**Intent/Clarification:**

Pulmonary stenosis (PS) is often due to congenital malformation of the valve. As it restricts blood flow from the right ventricle into the pulmonary artery, patients experience extreme fatigue and palpitations. Severe PS may create a bluish tint to skin and is life threatening.

Choose **highest** level of valve dysfunction when there are differences in interpretation of

most recent study.

Capture even if patient is not scheduled for valve repair and/or replacement when available.

- Yes
- No

Time Frame: Collect the last value closest to incision, **not greater than 6 months**.

SEQ. #: 1845

Long Name: VD-Pulmonic Hemodynamic Data Available

Short Name: PuHemoDatAvail

Definition: Indicate whether pulmonary valve gradient is available.

Intent/Clarification:

Time Frame: Collect the last value closest to incision, not greater than **6 months**.

- Yes
 - No
-
-

SEQ. #: 1850

Long Name: VD-Pulmonic Gradient-Highest Mean

Short Name: VDGradP

Definition: Indicate highest mean PV gradient documented prior to incision.

Intent/Clarification:

Time Frame: Collect the last value closest to incision, not greater than **6 months**.

SEQ. #: 1855

Long Name: VD-Pulmonic Valve Disease Etiology

Short Name: VDPuEt

Definition: Indicate the etiology of pulmonary valve disease if known.

Intent/Clarification:

There is no hierarchy, choose the primary etiology. Primary lesion may be identified at the time of the surgical procedure.

FAQ October 2017: My surgeon replaced a mitral valve in a patient with a history of Hodgkin's disease who is s/p chemotherapy and radiation therapy. The etiology is described as radiation induced heart disease. How should this be coded?

Answer: This should be coded as mixed etiology and would apply to pulmonic valve etiology. This will be added in the next upgrade. In addition to capturing the etiology, code mediastinal radiation in Seq. No. 495.

Operative

SEQ. #: 1955

Long Name: Surgeon

Short Name: Surgeon

Definition: Indicate the name of the surgeon responsible for the patient's care.

This field must have controlled data entry where a user selects the surgeon name from a user list. This will remove variation in spelling, abbreviations and punctuation within the field.

Intent/Clarification:

Field must be populated. Missing data or information for a surgeon not on your current contract with the STS will cause your data file submission not to process.

SEQ. #: 1960

Long Name: Surgeon's National Provider Identifier

Short Name: SurgNPI

Definition: Indicate the individual-level National Provider Identifier of the surgeon performing the procedure.

For Non-US surgeons a unique identifier will be assigned by STS.

Intent/Clarification:

Field must be populated. Missing or inaccurate data will cause your data file submission not to process. It is crucial to enter the correct surgeon identifier since it may impact public reporting and physician quality reporting. This link provides an NPI search –

<https://nppes.cms.hhs.gov/#/>

SEQ. #: 1965

Long Name: Taxpayer Identification Number

Short Name: TIN

Definition: Indicate the Taxpayer Identification Number for the Taxpayer holder of record for the Surgeon's National Provider Identifier that performed the procedure. This may be an individual TIN or a group TIN depending on billing. This information is vital for PQRS reporting.

This field will be blank for Non-US participants

Intent/Clarification: -

SEQ. #: 1966

Long Name: STS Risk Calculator Score Discussed

Short Name: RiskDiscussed

Definition: Indicate whether the STS Risk Calculator score was discussed with the patient/family prior to surgery.

Intent/Clarification:

To meet this measure, discussion should take place between the surgeon and patient/family and be documented. STS risk models are available for CABG, AVR, AVR + CABG, MVR, MVR + CABG, MV Repair, MV Repair + CABG and calculated in vendor software or using the STS Risk Calculator. For all other procedures code NA. The Euroscore cannot be used to complete this field.

Use of STS Risk Calculator is the ONLY way to select "YES" to this question. This is a MIPS reported measure.

<http://www.sts.org/quality-research-patient-safety/quality/mips-reporting-database>

SEQ. #: 1970

Long Name: Incidence

Short Name: Incidenc

Definition: Indicate if this is the patient's:

- first surgery
- first re-op surgery
- second re-op surgery
- third re-op surgery
- fourth or more re-op surgery

Intent/Clarification:

For the purposes of this field surgery is defined as cardiothoracic surgical procedures performed on the heart, great vessels or major pericardial procedures, with or without cardiopulmonary bypass (CPB). The key distinction is surgical entry into the pericardial space. A pericardiectomy or pericardial window would qualify as surgery. Ascending aortic and arch procedures also qualify. A surgical descending thoracic aortic aneurysmectomy does not involve entry into the pericardial space and does not qualify. Similarly, catheter based procedures such as TAVR, TEVAR, mitral-clip, are endovascular procedures and are not classified as prior surgery. Also include lung procedures utilizing CPB or tracheal procedures utilizing CPB. Reoperation increases risk due to presence of scar tissue or adhesions.

The intent of this field is to capture the incidence of the procedure that the patient is about to go through during the current hospitalization, as compared to those procedures prior to this hospitalization. First operative means the patient has never had any surgical procedure on the heart and/or great vessels. Note: previous surgical intervention increases risk for morbidity and mortality and severity of disease process.

SEQ. #: 1975

Long Name: Status

Short Name: Status

Definition: Indicate the clinical status of the patient prior to entering the operating room.

Intent/Clarification:

- **Elective**- The patient's cardiac function has been stable in the days or weeks prior to the operation. The procedure could be deferred without increased risk of compromised cardiac outcome.

- **Urgent** - Procedure required during same hospitalization in order to minimize chance of further clinical deterioration. Examples include but are not limited to: Worsening or sudden chest pain, CHF, acute myocardial infarction (AMI), anatomy, IABP, unstable angina (USA) with intravenous (IV) nitroglycerin (NTG) or rest angina.
 - Any of the conditions that require that the patient remain in the hospital until surgery can take place, but the patient is able to wait for surgery until the next available OR schedule time. Delay in the operation may be necessitated by attempts to improve the patient's condition, availability of a spouse or parent for informed consent, availability of blood products, or the availability of results of essential laboratory procedures or tests. **There is no hierarchy - choose the primary reason the procedure is urgent.**
 - If a patient has severe aortic and mitral valve stenosis, but also has symptoms such as dyspnea on exertion (DOE), paroxysmal nocturnal dyspnea (PND), congestion on x-ray or pedal edema that has been treated as CHF, code "CHF" as the most appropriate choice.
 - Valve dysfunction is defined as a structural failure with that valve. For prosthetic valves – fractured leaflet, thrombus formation, pannus development which impedes flow through the valve orifice, or valvular dehiscence (coming loose or disconnected at the suture line). Native valve dysfunction includes papillary rupture or torn leaflet. Rupture or dissection during cardiac cath; Perforation, tamponade following cardiac cath-does not include stent closure.

- **Emergent** - Patients requiring emergency operations will have ongoing, refractory (difficult, complicated, and/or unmanageable) unremitting cardiac compromise, with or without hemodynamic instability, and not responsive to any form of therapy except cardiac surgery. An emergency operation is one in which there should be no delay in providing operative intervention. Patients requiring emergency operations will have ongoing, refractory (difficult, complicated, and/or unmanageable) cardiac compromise, with or without hemodynamic instability, and not responsive to any form of therapy except cardiac surgery. Hemodynamic picture of shock that is being chemically or mechanically supported. (IV inotrope or IABP to maintain cardiac output [CO]. Requires intubation and ventilation for pulmonary edema. The patient is extending an MI and requires immediate surgery. The patient continues to show signs of ongoing ischemia, i.e. EKG changes. Acute native valve dysfunction i.e. as acute papillary muscle rupture or torn leaflet. Prosthetic

valve dysfunction is defined as a structural failure with that valve-fractured or torn leaflet, thrombus formation, pannus development which impedes flow through the valve orifice, or valvular dehiscence (coming loose or disconnected at the suture line). Acute dissection secondary to trauma or dissection secondary to progression of disease. Rupture or dissection during cardiac cath; perforation, tamponade following cardiac cath.

- If a patient presents with a scenario that does not fit into a definite category; it is reasonable to code the reason that most closely matches the patient's presentation.
- **Emergent/Salvage** - The patient is undergoing CPR en-route to the OR prior to anesthesia induction or has ongoing ECMO to maintain life.

SEQ. #: 1990

Long Name: Urgent Or Emergent Reason

Short Name: UrgEmergRsn

Definition: Choose one reason from the list below that best describes why this operation was considered urgent or emergent.

Intent/Clarification:

See list for options. There may be multiple reasons, choose one that best describes this patient's clinical state.

SEQ. #: 1995

Long Name: Previously Attempted Case Canceled

Short Name: PCancCase

Definition: Indicate whether this case was previously attempted during this admission and cancelled or aborted after patient entered the operating room.

Intent/Clarification:

To capture occasions when the patient goes to the operating room with the intention to perform a cardiac surgery but the case is canceled.

SEQ. #: 2000

Long Name: Previously Attempted Canceled Case Date

Short Name: PCancCaseDt

Definition: Enter date previously attempted case was cancelled.

Intent/Clarification:

Date must be during this hospital admission. Required date format: mm/dd/yyyy.

SEQ. #: 2005

Long Name: Previously Attempted Canceled Case Timing

Short Name: PCancCaseTmg

Definition: Indicate at what point previously attempted case was cancelled or aborted.

Intent/Clarification:

The intent is to capture the timing associated with cancelling the case:

- Prior to Induction of Anesthesia
 - After Induction, Prior to Incision
 - After Incision Made
-
-

SEQ. #: 2010

Long Name: Previously Attempted Canceled Case Reason

Short Name: PCancCaseRsn

Definition: Indicate the reason why the previously attempted case was cancelled or aborted.

Intent/Clarification:

The intent is to capture the reason for cancelling the case:

- Anesthesiology event - Includes airway, line insertion and medication issues encountered during induction
 - Cardiac arrest - Patient deterioration unrelated to induction
 - Equipment/supply issue - Device malfunction or supply issue including devices and blood products needed for surgery but not available
 - Access issue – Unable to gain access for lines and/or surgical exposure
 - Unanticipated tumor – Tumor discovered at time of surgery
 - Donor organ unacceptable – Organs for transplant found to be unacceptable
 - Abnormal labs – Lab results could increase risk of surgery and/or require intervention prior to surgery
 - Other – Reason not specified above
-
-

SEQ. #: 2015

Long Name: Previously Attempted Cancelled Case Procedure - CABG

Short Name: PCancCaseCAB

Definition: Indicate whether the plan for the previously attempted procedure included coronary artery bypass grafting.

Intent/Clarification:

The intent is to capture if the intended procedure for the cancelled case was Coronary Artery Bypass.

SEQ. #: 2020

Long Name: Previously Attempted Canceled Case Procedure - Mechanical Assist Device

Short Name: PCancCaseMech

Definition: Indicate whether the plan for the previously attempted procedure included implanting or explanting a mechanical assist device.

Intent/Clarification:

The intent is to capture if the intended procedure for the cancelled case was a mechanical assist device.

SEQ. #: 2025

Long Name: Previously Attempted Canceled Case Procedure - Other Non-Cardiac

Short Name: PCancCaseONC

Definition: Indicate whether the plan for the previously attempted procedure included any other non-cardiac procedure.

Intent/Clarification:

The intent is to capture if the intended procedure for the cancelled case was any other non-cardiac procedure.

SEQ. #: 2030

Long Name: Previously Attempted Canceled Case Procedure - Valve, Surgical

Short Name: PCancCaseValSur

Definition: Indicate whether the plan for the previously attempted procedure included a surgical valve procedure.

Intent/Clarification:

The intent is to capture if the intended procedure for the cancelled case was a surgical valve procedure.

SEQ. #: 2035

Long Name: Previously Attempted Canceled Case Procedure - Valve, Transcatheter

Short Name: PCancCaseValTrans

Definition: Indicate whether the plan for the previously attempted procedure included a transcatheter valve procedure.

Intent/Clarification:

The intent is to capture if the intended procedure for the cancelled case was a transcatheter valve procedure.

SEQ. #: 2040

Long Name: Previously Attempted Canceled Case Procedure - Other Cardiac

Short Name: PCancCaseOC

Definition: Indicate whether the plan for the previously attempted procedure included any other cardiac procedure.

Intent/Clarification:

The intent is to capture if the intended procedure for the cancelled case was any other cardiac procedure.

SEQ. #: 2050

Long Name: Current Case Canceled

Short Name: CCancCase

Definition: Indicate whether the current case was canceled or aborted after patient entered the operating room.

Intent/Clarification:

The intent is to capture if the current case that was cancelled occurred after the patient entered the operating room.

SEQ. #: 2055

Long Name: Current Case Canceled Timing

Short Name: CCancCaseTmg

Definition: Indicate at what point the current case was canceled or aborted.

Intent/Clarification:

The intent is to capture the timing associated with cancelling the case:

- Prior to Induction of Anesthesia
 - After Induction, Prior to Incision
 - After Incision Made
-
-

SEQ. #: 2060

Long Name: Current Case Canceled Reason

Short Name: CCancCaseRsn

Definition: Indicate the reason why the current case was canceled or aborted.

Intent/Clarification:

The intent is to capture the reason for cancelling the case:

- **Anesthesiology event** - Includes airway, line insertion and medication issues encountered during induction
 - **Cardiac arrest** - Patient deterioration unrelated to induction
 - **Equipment/supply issue** - Device malfunction or supply issue including devices and blood products needed for surgery but not available
 - **Access issue** – Unable to gain access for lines and/or surgical exposure
 - **Unanticipated tumor** – Tumor discovered at time of surgery
 - **Donor organ unacceptable** – Organs for transplant found to be unacceptable
 - **Abnormal labs** – Lab results could increase risk of surgery and/or require intervention prior to surgery
 - **Other** – Reason not specified above
-
-

SEQ. #: 2065

Long Name: Current Case Canceled Procedure - CABG

Short Name: CCancCaseCAB

Definition: Indicate whether the plan for the current procedure included coronary artery bypass grafting.

Intent/Clarification:

The intent is to capture if the intended procedure for the cancelled case included Coronary Artery Bypass.

SEQ. #: 2075

Long Name: Current Case Canceled Procedure - Mechanical Assist Device

Short Name: CCancCaseMech

Definition: Indicate whether the plan for the current procedure included implanting or explanting a mechanical assist device.

Intent/Clarification:

The intent is to capture if the intended procedure for the cancelled case was a mechanical assist device.

SEQ. #: 2080

Long Name: Current Case Canceled Procedure - Other Non-cardiac

Short Name: CCancCaseONC

Definition: Indicate whether the plan for the current procedure included any other non-cardiac procedure.

Intent/Clarification:

The intent is to capture if the intended procedure for the cancelled case was any other non-cardiac procedure.

SEQ. #: 2085

Long Name: Current Case Canceled Procedure - Valve, Surgical

Short Name: CCancCaseValSur

Definition: Indicate whether the plan for the previously attempted procedure included a surgical valve procedure.

Intent/Clarification:

The intent is to capture if the intended procedure for the cancelled case was a surgical valve procedure.

SEQ. #: 2090

Long Name: Current Case Canceled Procedure - Valve, Transcatheter

Short Name: CCancCaseValTrans

Definition: Indicate whether the plan for the previously attempted procedure included a transcatheter valve procedure.

Intent/Clarification:

The intent is to capture if the intended procedure for the cancelled case was a transcatheter valve procedure.

SEQ. #: 2095

Long Name: Current Case Canceled Procedure - Other Cardiac

Short Name: CCancCaseOC

Definition: Indicate whether the plan for the current procedure included any other cardiac procedure.

Intent/Clarification:

The intent is to capture if the intended procedure for the cancelled case was any other cardiac procedure.

SEQ. #: 2100

Long Name: Operative Approach

Short Name: OPApp

Definition: Indicate the initial operative approach.

Intent/Clarification:

The intent is to capture the **initial** operative approach:

- Full conventional sternotomy
- Partial sternotomy
- Transverse sternotomy (includes clamshell)
- RIGHT OR LEFT parasternal incision
- Sub-xiphoid

- Sub-Costal
- Left Thoracotomy
- Right Thoracotomy
- Bilateral Thoracotomy
- Limited (mini) Thoracotomy, right (transapical TAVR)
- Limited (mini) Thoracotomy, left
- Limited (mini) Thoracotomy, bilateral
- Thoracoabdominal Incision
- Percutaneous
- Port Access
- Other
- None (cancelled case)

Commonly used approaches for the following devices:

- **Impella 2.0**
 - Percutaneous femoral
 - Percutaneous iliac
- **Impella 5.0**
 - Percutaneous femoral
 - Open femoral
 - Open aorta
 - Open iliac
- **VA ECMO**
 - Percutaneous femoral
 - Open Femoral
 - Percutaneous carotid
 - Percutaneous subclavian
 - Open subclavian

SEQ. #: 2105

Long Name: Operative Approach Converted

Short Name: ApproachCon

Definition: Indicate whether the operative approach was converted during the procedure.

Intent/Clarification:

The intent is to capture whether the approach was converted and whether that was part of the initial surgical plan.

- Yes, planned
- Yes, unplanned
- No

SEQ. #: 2110

Long Name: Robot Used

Short Name: Robotic

Definition: Indicate whether a robot was used during cardiac surgery.

Intent/Clarification:

The intent is to whether any portion of the procedure was used during the surgical procedure.

- Yes
 - No
-
-

SEQ. #: 2115

Long Name: Robot Use Time Frame

Short Name: RobotTim

Definition: Indicate the time frame of robotic use.

Intent/Clarification:

The intent is to the extent of the procedure where the robot was used.

- Used for entire operation
 - Used for part of the operation
-
-

SEQ. #: 2120

Long Name: CAB

Short Name: OpCAB

Definition: Indicate whether coronary artery bypass grafting was done.

Intent/Clarification:

The intent is to capture procedures where bypass grafts were constructed to native coronary arteries.

- Yes, planned*
- Yes, unplanned due to surgical complication*
- Yes, unplanned due to unsuspected disease or anatomy*
- No

***If yes, complete Section J.**

FAQ September 2017: Pt with previous CAB in January of this year now presents with Aortic Dissection. The dissection was repaired and cadaver vein graft extensions were used to reimplant the PDA/PL grafts. The OM and Diag grafts were occluded by the dissection. Should this case include a CAB?

Answer: No, a distal coronary artery graft is not constructed using the cadaver vein.

FAQ October 2017: Following a David procedure, due to bleeding at the right coronary anastomosis, a Kay Zubiato reconstruction was required to reconstruct the ostium of the RCA using a segment of saphenous vein. How is this coded?

Answer: Code CAB; Yes, unplanned due to surgical complication.

SEQ. #: 2125

Long Name: Valve

Short Name: OpValve

Definition: Indicate whether a surgical procedure was done on the Aortic, Mitral, Tricuspid or Pulmonic valves.

Intent/Clarification:

The intent is to capture procedures where valve procedures were performed.

- Yes*
- No

***If yes, complete Section K.**

SEQ. #: 2126

Long Name: Surgeon Input for Valve Surgery Data Abstraction

Short Name: OpValSurgInput

Definition: Indicate whether the surgeon provided input for the valve surgery data abstraction.

Intent/Clarification:

Indicates that the data manager confirms that because of interaction with the surgeon the data included reflects the pathology and the procedure performed.

SEQ. #: 2128

Long Name: Aorta Procedure Performed

Short Name: AortProc

Definition: Indicate whether a procedure was performed on the aorta.

Intent/Clarification:

The intent is to capture procedures where procedures were performed involving the aorta.

- Yes, planned*
- Yes, unplanned due to surgical complication*
- Yes, unplanned due to unsuspected disease or anatomy*
- No

***If Yes, complete Section M2**

SEQ. #: 2129

Long Name: Surgeon Input for Aortic Surgery Data Abstraction

Short Name: AortProcSurgInput

Definition: Indicate whether the surgeon provided input for the aortic surgery data abstraction.

Intent/Clarification:

Indicates that the data manager confirms that because of interaction with the surgeon the data included reflects the pathology and the procedure performed.

- Yes
- No

SEQ. #: 2140

Long Name: Other Card

Short Name: OpOCard

Definition: Indicate whether another cardiac procedure was done (other than CABG and/or Valve procedures).

Intent/Clarification:

The intent is to capture procedures where procedures were performed involving the aorta.

- Yes*
- No

***If Yes, complete Section M.**

Do not include isolated ECMO, Impella or IABP insertions here.

FAQ September 2017: If a patient has an extensive atrial fibrillation procedure and an LAA clipping, how would we include the atrial ligation/exclusion method and device?
Answer: Multiple sequence numbers will be required to code the ligation/exclusion and device.

First capture sequence numbers 2140 and 2145 to open section M and M-1.

Then capture the atrial appendage procedure in sequence 4050; then ligation/exclusion method in sequence 4051 and Model in sequence number 4052 as well as the UDI, if available, in sequence 4053.

Code the extensive ablation procedure in section M-1 as described by your physician including lesion number 7 to complete the LAA.

FAQ October 2017: A patient has a four vessel bypass. During the procedure he also has an endarterectomy and vein patch angioplasty of the circumflex (no bypass). Do I code that a coronary bypass graft was done? Do I capture this as Other Cardiac thus throwing this out of the isolated CABG category?

Answer: In order to open the child fields in the coronary bypass grid to complete the fields for endarterectomy you must code 'Yes'. Leave proximal site blank. Do not code as 'Other Cardiac'.

SEQ. #: 2145

Long Name: Atrial Fibrillation Procedure Performed

Short Name: AFibProc

Definition: Indicate whether an atrial fibrillation procedure was performed.

Intent/Clarification:

The intent is to capture when atrial fibrillation procedures were performed.

- Yes*
- No

***If Yes, complete Section M1.**

FAQ September 2017: If a patient has an extensive atrial fibrillation procedure and an LAA clipping how would we include the atrial ligation/exclusion method and device?

Answer: Multiple sequence numbers will be required to code the ligation/exclusion and device.

First capture sequence numbers 2140 and 2145 to open section M and M-1.

Then capture the atrial appendage procedure in sequence 4050; then ligation/exclusion method in sequence 4051 and Model in sequence number 4052 as well as the UDI, if available, in sequence 4053.

Code the extensive ablation procedure in section M-1 as described by your physician including lesion number 7 to complete the LAA.

SEQ. #: 2146

Long Name: Surgeon Input for Other Cardiac Afib Data Abstraction

Short Name: AFibProcSurgInput

Definition: Indicate whether the surgeon provided input for the other cardiac Afib procedure data abstraction.

Intent/Clarification:

Indicates that the data manger confirms that because of interaction with the surgeon the data included reflects the pathology and the procedure performed.

- Yes*
- No

SEQ. #: 2155

Long Name: Other Non Card

Short Name: OpONCard

Definition: Indicate whether a non-cardiac procedure was done.

Intent/Clarification:

The intent is to capture when non-cardiac procedures were performed.

SEQ. #: 2195

Long Name: CPT-1 Code # 1

Short Name: CPT1Code1

Definition: Indicate the first CPT procedure code (CPT-1) pertaining to the surgery for which the data collection form was initiated.

Intent/Clarification:

There is no STS list. Use whichever CPT codes were entered for procedures performed during this operation. Consult with your Billing/Coding Department, if applicable.

SEQ. #: 2200

Long Name: CPT-1 Code # 2

Short Name: CPT1Code2

Definition: Indicate, if applicable, the second CPT procedure code (CPT-1) pertaining to the surgery for which the data collection form was initiated.

Intent/Clarification:

There is no STS list. Use whichever CPT codes were entered for procedures performed during this operation. Consult with your Billing/Coding Department, if applicable.

SEQ. #: 2205

Long Name: CPT-1 Code # 3

Short Name: CPT1Code3

Definition: Indicate, if applicable, the third CPT procedure code (CPT-1) pertaining to the surgery for which the data collection form was initiated.

Intent/Clarification:

There is no STS list. Use whichever CPT codes were entered for procedures performed during this operation. Consult with your Billing/Coding Department, if applicable.

SEQ. #: 2210

Long Name: CPT-1 Code # 4

Short Name: CPT1Code4

Definition: Indicate, if applicable, the fourth CPT procedure code (CPT-1) pertaining to the surgery for which the data collection form was initiated.

Intent/Clarification:

There is no STS list. Use whichever CPT codes were entered for procedures performed during this operation. Consult with your Billing/Coding Department, if applicable.

SEQ. #: 2215

Long Name: CPT-1 Code # 5 **Short Name:** CPT1Code5

Definition: Indicate, if applicable, the fifth CPT procedure code (CPT-1) pertaining to the surgery for which the data collection form was initiated.

Intent/Clarification:

There is no STS list. Use whichever CPT codes were entered for procedures performed during this operation. Consult with your Billing/Coding Department, if applicable.

SEQ. #: 2220

Long Name: CPT-1 Code # 6

Short Name: CPT1Code6

Definition: Indicate, if applicable, the sixth CPT procedure code (CPT-1) pertaining to the surgery for which the data collection form was initiated.

Intent/Clarification:

There is no STS list. Use whichever CPT codes were entered for procedures performed during this operation. Consult with your Billing/Coding Department, if applicable.

SEQ. #: 2225

Long Name: CPT-1 Code # 7

Short Name: CPT1Code7

Definition: Indicate, if applicable, the seventh CPT procedure code (CPT-1) pertaining to the surgery for which the data collection form was initiated.

Intent/Clarification:

There is no STS list. Use whichever CPT codes were entered for procedures performed during this operation. Consult with your Billing/Coding Department, if applicable.

SEQ. #: 2230

Long Name: CPT-1 Code # 8

Short Name: CPT1Code8

Definition: Indicate, if applicable, the eighth CPT procedure code (CPT-1) pertaining to the surgery for which the data collection form was initiated.

Intent/Clarification:

There is no STS list. Use whichever CPT codes were entered for procedures performed during this operation. Consult with your Billing/Coding Department, if applicable.

SEQ. #: 2235

Long Name: CPT-1 Code # 9

Short Name: CPT1Code9

Definition: Indicate, if applicable, the ninth CPT procedure code (CPT-1) pertaining to the surgery for which the data collection form was initiated.

Intent/Clarification:

There is no STS list. Use whichever CPT codes were entered for procedures performed during this operation. Consult with your Billing/Coding Department, if applicable.

SEQ. #: 2240

Long Name: CPT-1 Code # 10

Short Name: CPT1Code10

Definition: Indicate, if applicable, the tenth CPT procedure code (CPT-1) pertaining to the surgery for which the data collection form was initiated.

Intent/Clarification:

There is no STS list. Use whichever CPT codes were entered for procedures performed during this operation. Consult with your Billing/Coding Department, if applicable.

SEQ. #: 2245

Long Name: OR Entry Date And Time

Short Name: OREntryDT

Definition: Indicate the date and time, to the nearest minute (using 24-hour clock), that the patient entered the operating room. If the procedure was performed in a location other than the OR, record the time when the sterile field, or its equivalent, was set up.

Intent/Clarification:

The intent is to capture the actual date and time the patient physically enters the operating room. For emergency procedures done outside the OR, this may be an estimated time.

Required date format: mm/dd/yyyy

Required time format: hh:mm (0-24 hour clock)

SEQ. #: 2250

Long Name: OR Exit Date And Time

Short Name: ORExitDT

Definition: Indicate the date and time, to the nearest minute (using 24-hour clock), that the patient exits the operating room. If the procedure was performed in a location other than the OR, record the time when the sterile field, or its equivalent, was taken down.

Intent/Clarification:

The intent is to capture the actual date and time the patient physically leaves the operating room. This field is used to calculate post-operative ventilation time and therefore prolonged ventilation.

Required date format: mm/dd/yyyy.

Required time format: hh:mm (0-24 hour clock)

SEQ. #: 2251

Long Name: General Anesthesia

Short Name: GenAnes

Definition: Indicate whether general anesthesia was used during this procedure.

Intent/Clarification:

The intent is to clarify whether general anesthesia was used.

SEQ. #: 2252

Long Name: Procedural Sedation

Short Name: ProcSed

Definition: Indicate whether the procedure was performed under sedation (also referred to as “moderate sedation” or “conscious sedation”) and not general anesthesia.

Intent/Clarification:

The intent is to identify whether sedation was used, moderate or conscious, instead of general anesthesia.

SEQ. #: 2253

Long Name: Intubation

Short Name: Intubate

Definition: Indicate the status of intubation.

Intent/Clarification:

The intent is to identify whether the patient required endotracheal or tracheal intubation.

SEQ. #: 2255

Long Name: Initial Intubation Date And Time

Short Name: IntubateDT

Definition: Indicate the date (mm/dd/yyyy) and time (hh:mm) (24 hour clock) ventilatory support started. Date in the format mm/dd/yyyy.

Intent/Clarification:

The following guidelines apply:

- Capture the intubation closest to the surgical start time. If the patient was intubated upon admission and remained intubated until the surgical start time, capture this intubation's date and time.
- If the patient was admitted intubated (intubated at another institution) and remained continually intubated until the surgical start time, capture the patient's admission date and time.
- If the patient was admitted with a tracheostomy in place without ventilator support, capture the date and time closest to the surgical start time that ventilator support was initiated.
- If the patient was admitted with a tracheostomy in place and was receiving chronic ventilator support, capture the admission date and time.
- If the intubation date and time is otherwise unknown, enter the date and time the patient entered the operating room.
- Do not alter the previously established date and time that ventilator support was initiated for scenarios including, but not limited to, interruptions in ventilator support due to accidental extubation/de-cannulation, elective tube change etc.

Required date format: mm/dd/yyyy

Required time format: hh:mm (0-24 hour clock)

SEQ. #: 2260

Long Name: Initial Extubation Date And Time

Short Name: ExtubateDT

Definition: Indicate the date (mm/dd/yyyy) and time (hh:mm) (24 hour clock) ventilatory support initially ceased after surgery.

Intent/Clarification:

The following guidelines apply:

- Capture extubation time closest to the surgical stop time.
- If the patient has a tracheostomy and is separated from the mechanical ventilator postoperatively within the hospital admission, capture the date and time of separation from the mechanical ventilator closest to the surgical stop time.
- If the patient expires while intubated or cannulated and on the ventilator, capture the date and time of expiration as extubation time.
- If patient is discharged on chronic ventilator support, capture the date and time of discharge.

Required date format: mm/dd/yyyy

Required time format: hh:mm (0-24 hour clock)

SEQ. #: 2265

Long Name: Skin Incision Start Date And Time

Short Name: SIStartDT

Definition: Indicate the date and time, to the nearest minute (using 24-hour clock), that the first skin incision, or its equivalent, was made.

Intent/Clarification:

Use the first incision, i.e. vein harvest incision; for the skin incision date and time. Do not code access site stab wounds.

Required date format: mm/dd/yyyy

Required time format: hh:mm (0-24 hour clock)

SEQ. #: 2270

Long Name: Skin Incision Stop Date And Time

Short Name: SIStopDT

Definition: Indicate the date and time, to the nearest minute (using 24-hour clock), that the skin incision was closed, or its equivalent. If the patient leaves the operating room with an open incision, collect the time that the dressings were applied to the incision.

Intent/Clarification:

Use the documented time the incision was closed.

Required date format: mm/dd/yyyy

Required time format: hh:mm (0-24 hour clock)

SEQ. #: 2275

Long Name: Anesthesia End Date and Time

Short Name: AnesEndDT

Definition: Indicate the anesthesia end time documented in the medical record. The definition of anesthesia end time is when the anesthesiologist is no longer in personal attendance, that is, when the patient is safely placed under post-anesthesia supervision.

Intent/Clarification:

The time may be in the recovery room or intensive care unit; when it is documented that anesthesia care has ended. This field will be referenced for selecting the peak post op glucose (4550) Anesthesia end time should be captured from the anesthesia record.

Required date format: mm/dd/yyyy

Required time format: hh:mm (0-24 hour clock)

SEQ. #: 2280

Long Name: Appropriate Antibiotic Selection

Short Name: AbxSelect

Definition: Indicate if there was documentation of an order for a first generation or second generation cephalosporin prophylactic antibiotic, documentation that it was given preoperatively or in the event of a documented allergy an alternate antibiotic choice is ordered and administered.

Intent/Clarification: Refer to the antibiotic guidelines on the STS website.

<http://www.sts.org/resources-publications/clinical-practice-credentialing-guidelines/antibiotic-guidelines>

SEQ. #: 2285

Long Name: Appropriate Antibiotic Administration Timing

Short Name: AbxTiming

Definition: Indicate whether prophylactic antibiotics were administered within one hour of surgical incision or start of procedure if no incision required (two hours if receiving Vancomycin or fluoroquinolone).

The surgical incision time is the time of the first incision, regardless of location.

Intent/Clarification:

The documented time the antibiotic started must be prior to the documented time of the first surgical incision. Refer to antibiotic guidelines on the STS website.

<http://www.sts.org/resources-publications/clinical-practice-credentialing-guidelines/antibiotic-guidelines>

SEQ. #: 2290

Long Name: Appropriate Antibiotic Discontinuation

Short Name: AbxDisc

Definition: Indicate whether the prophylactic antibiotics were ordered to be discontinued OR were discontinued within 48 hours after surgery end time.

Determining the timeframe (within 48 hours) begins at the "surgical end time".

Intent/Clarification:

Refer to antibiotic guidelines on the STS website.

<http://www.sts.org/resources-publications/clinical-practice-credentialing-guidelines/antibiotic-guidelines>

SEQ. #: 2295

Long Name: Additional Intraoperative Prophylactic Antibiotic Dose

Short Name: AddIntraopPAnti

Definition: Indicate whether an additional prophylactic antibiotic dose was given in the operating room.

Intent/Clarification:

Refer to antibiotic guidelines on the STS website.

<http://www.sts.org/resources-publications/clinical-practice-credentialing-guidelines/antibiotic-guidelines>

SEQ. #: 2296

Long Name: Temperature Measured

Short Name: TempMeas

Definition: Indicate whether the patient's core temperature was measured during the procedure.

Intent/Clarification:

SEQ. #: 2300

Long Name: Lowest Temperature

Short Name: LwstTemp

Definition: Record the patient's lowest core temperature in the operating room in degrees centigrade.

Intent/Clarification:

The intent is to capture the lowest documented temperature intraoperatively. The source of the documentation may be Esophageal, CPB venous return, Bladder, Nasopharyngeal, Tympanic, Rectal, or Other.

SEQ. #: 2305

Long Name: Lowest Temperature Source

Short Name: LwstTempSrc

Definition: Indicate the source where the lowest core temperature was measured.

Intent/Clarification:

Temperatures are typically documented on perfusion record or anesthesia record. Venous temperatures on CPB are most common and always available, however not as

accurate. Sources may be Esophageal, CPB venous return, Bladder, Nasopharyngeal, Tympanic, Rectal, Other, or Unknown.

SEQ. #: 2310

Long Name: Lowest Intra-op Hemoglobin

Short Name: LwstIntraHemo

Definition: Enter the lowest measured hemoglobin recorded in the operating room. Do not enter calculated values.

Intent/Clarification:

If you do not have measured lab values you may use calculated values.

**Note that Hemoglobin (Hgb) should always be less than the Hematocrit (Hct).

SEQ. #: 2315

Long Name: Lowest Hematocrit

Short Name: LwstHct

Definition: Enter the lowest measured hematocrit recorded in the operating room. Do not enter calculated values.

Intent/Clarification:

If you do not have measured lab values you may use calculated values.

**Note that Hemoglobin (Hgb) should always be less than the Hematocrit (Hct).

SEQ. #: 2320

Long Name: Highest Intra-op Glucose

Short Name: HighIntraGlu

Definition: Enter the highest glucose recorded in the operating room.

Intent/Clarification:

Typically documented in laboratory tests, anesthesia record, or perfusion record.

SEQ. #: 2325

Long Name: CPB Utilization

Short Name: CPBUtil

Definition: Indicate the level of CPB or coronary perfusion used during the procedure.

Intent/Clarification:

- **None:** No CPB or coronary perfusion used during the procedure.

- **Combination:** With or without CPB and/or with or without coronary perfusion at any time during the procedure (capture conversions from off-pump to on-pump only):
 - At start of procedure: No CPB/No Coronary Perfusion -> conversion to -> CPB
 - At start of procedure: No CPB/No Coronary Perfusion -> conversion to -> Coronary perfusion
 - At start of procedure: No CPB/No Coronary Perfusion -> conversion to -> Coronary perfusion -> conversion to -> CPB
 - **Full CPB** or coronary perfusion was used for the entire procedure
-
-

SEQ. #: 2330

Long Name: CPB Utilization - Combination Plan

Short Name: CPBCmb

Definition: Indicate whether the combination procedure from off-pump to on-pump was a planned or an unplanned conversion.

Intent/Clarification:

To capture if the operation was intended to be an off pump case and, for some clinical reason, required cardiopulmonary bypass to complete the operation.

- **Planned** - The surgeon intended to treat with any of the combination options described in "CPB utilization"
 - **Unplanned** - The surgeon did not intend to treat with any of the combination options described in "CPB utilization"
-
-

SEQ. #: 2335

Long Name: CPB Utilization - Unplanned Combination Reason

Short Name: CPBCmbR

Definition: Indicate the reason that the procedure required the initiation of CPB and/or coronary perfusion.

Intent/Clarification:

To capture the reason that caused the procedure to require the initiation of cardiopulmonary bypass:

- Exposure/visualization
 - Bleeding
 - Inadequate size and/or diffuse disease of the distal vessel
 - Hemodynamic instability (hypotension/arrhythmias)
 - Conduit quality and/or trauma
 - Other
-
-

SEQ. #: 2340

Long Name: Cannulation - Arterial Cannulation Site - Aortic

Short Name: CanArtStAort

Definition: Indicate whether the arterial cannulation site included the aorta.

Intent/Clarification:

The arterial cannulation site was the aorta.

SEQ. #: 2345

Long Name: Cannulation - Arterial Cannulation Site - Femoral

Short Name: CanArtStFem

Definition: Indicate whether the arterial cannulation site included a femoral artery.

Intent/Clarification:

The arterial cannulation site was the femoral artery.

SEQ. #: 2350

Long Name: Cannulation - Arterial Cannulation Site - Axillary

Short Name: CanArtStAx

Definition: Indicate whether the arterial cannulation site included an axillary artery.

Intent/Clarification:

The arterial cannulation site was the axillary artery.

SEQ. #: 2355

Long Name: Cannulation - Arterial Cannulation Site - Innominate

Short Name: CanArtStInn

Definition: Indicate whether the arterial cannulation site included an innominate artery.

Intent/Clarification:

The arterial cannulation site was the innominate artery.

SEQ. #: 2360

Long Name: Cannulation - Arterial Cannulation Site - Other

Short Name: CanArtStOth

Definition: Indicate whether the arterial cannulation site included any other artery.

Intent/Clarification:

There was any other arterial cannulation site.

SEQ. #: 2365

Long Name: Cannulation - Venous Cannulation Site - Femoral

Short Name: CanVenStFem

Definition: Indicate whether the venous (inflow) cannulation site included a femoral vein.

Intent/Clarification:

The venous cannulation site was the femoral vein.

SEQ. #: 2370

Long Name: Cannulation - Venous Cannulation Site - Jugular

Short Name: CanVenStJug

Definition: Indicate whether the venous (inflow) cannulation site included a jugular vein.

Intent/Clarification:

The venous cannulation site was the jugular vein.

SEQ. #: 2375

Long Name: Cannulation - Venous Cannulation Site - Right Atrial

Short Name: CanVenStRtA

Definition: Indicate whether the venous (inflow) cannulation site included the right atrium.

Intent/Clarification:

The venous cannulation site was the right atrium.

SEQ. #: 2380

Long Name: Cannulation - Venous Cannulation Site - Left Atrial

Short Name: CanVenStLfA

Definition: Indicate whether the venous (inflow) cannulation site included the left atrium.

Intent/Clarification:

The venous cannulation site was the left atrium.

SEQ. #: 2385

Long Name: Cannulation - Venous Cannulation Site - Pulmonary Vein

Short Name: CanVenStPulm

Definition: Indicate whether the venous (inflow) cannulation site included a pulmonary vein.

Intent/Clarification:

The venous cannulation site was the pulmonary vein.

SEQ. #: 2390

Long Name: Cannulation - Venous Cannulation Site - Caval/Bicaval

Short Name: CanVenStBi

Definition: Indicate whether the venous (inflow) cannulation site included the superior and/or inferior vena cava.

Intent/Clarification:

The venous cannulation site was the superior and/or inferior vena cava.

SEQ. #: 2395

Long Name: Cannulation - Venous Cannulation Site - Other

Short Name: CanVenStOth

Definition: Indicate whether the venous (inflow) cannulation site included any other site.

Intent/Clarification:

Any other venous cannulation site was used.

SEQ. #: 2400

Long Name: Cardiopulmonary Bypass Time

Short Name: PerfusTm

Definition: Indicate the total number of minutes that systemic return is diverted into the cardiopulmonary bypass (CPB) circuit and returned to the systemic system. This time period (Cardiopulmonary Bypass Time) includes all periods of cerebral perfusion and sucker bypass. This time period (Cardiopulmonary Bypass Time) excludes any circulatory arrest and modified ultrafiltration periods. If more than one period of CPB is required during the surgical procedure, the sum of all the CPB periods will equal the total number of CPB minutes.

Intent/Clarification:

The total time in minutes. This information can be obtained from the perfusion record or in the Surgeon's dictation.

SEQ. #: 2405

Long Name: Circulatory Arrest

Short Name: CircArr

Definition: Indicate whether or not circulatory arrest was utilized during the procedure.

Intent/Clarification:

Circulatory arrest is defined as the complete cessation of blood flow to the patient. Circulatory arrest is a surgical technique that involves cooling the body of the patient and stopping blood circulation and is not the same as coronary-pulmonary bypass time. It is used in cardiac surgery to allow operation on the aortic arch and in neurosurgery to repair some brain aneurysms.

SEQ. #: 2410

Long Name: Circulatory Arrest Time Without Cerebral Perfusion

Short Name: DHCATm

Definition: Indicate the total number of minutes of deep hypothermic circulatory arrest without cerebral perfusion. If more than one period of circulatory arrest is required during this surgical procedure, the sum of these periods is equal to the total duration of circulatory arrest.

Intent/Clarification:

If more than one period of circulatory arrest with cerebral perfusion is required during this surgical procedure, the sum of these periods is equal to the total duration of circulatory arrest without cerebral perfusion.

SEQ. #: 2415

Long Name: Circulatory Arrest With Cerebral Perfusion

Short Name: CPerfUtil

Definition: Indicate whether circulatory arrest with cerebral perfusion was performed.

Intent/Clarification:

Selective cerebral perfusion is a technique that involves providing blood flow and metabolic support to the brain while the blood flow to the rest of the body is stopped during circulatory arrest. This approach is commonly used during complex surgery that requires circulatory arrest. It offers more protection for the brain and minimizes the risk of stroke and other serious complications.

SEQ. #: 2420

Long Name: Cerebral Perfusion Time

Short Name: CPerfTime

Definition: Indicate the total number of minutes cerebral perfusion was performed. This would include antegrade and/or retrograde cerebral perfusion strategies.

Intent/Clarification:

If more than one period of circulatory arrest with cerebral perfusion was used, add the times for the total circulatory arrest with cerebral perfusion time.

SEQ. #: 2425**Long Name:** Cerebral Perfusion Type**Short Name:** CPerfTyp**Definition:** Indicate type of cerebral perfusion utilized.**Intent/Clarification:**

Indicate the type of cerebral perfusion:

- Antegrade
 - Retrograde
 - Both antegrade and retrograde
-
-

SEQ. #: 2426**Long Name:** Total Circulatory Arrest Time**Short Name:** TotCircArrTm**Definition:** Calculated variable measuring circulatory arrest without cerebral perfusion time plus any cerebral perfusion time.**Intent/Clarification:**

This value will be automatically generated by the software. It will total the number of minutes of circulatory arrest without cerebral perfusion + the total number of minutes of circulatory arrest with cerebral perfusion.

SEQ. #: 2430**Long Name:** Aortic Occlusion**Short Name:** AortOccl**Definition:** Indicate the technique of aortic occlusion used.**Intent/Clarification:**

Identify the method used to prevent blood from circulating through the heart and to allow the delivery of cardioplegia into the aortic root to arrest the heart. In procedures where cardioplegia is not administered for myocardial protection, but a cross clamp is applied to isolated diseased sections of the aorta (i.e. descending thoracic or thoracoabdominal aneurysm repairs) the appropriate response to aortic occlusion is aortic cross clamp. You should populate the cross clamp time field with the appropriate minutes of cross clamp time. The Cardioplegia field would be equal to None.

Externally, the aortic cross clamp is used. Internally, balloon occlusion is used. Choose one of the following:

- None - beating heart
 - None - fibrillating heart
 - Aortic Cross clamp
 - Balloon Occlusion
-
-

SEQ. #: 2435

Long Name: Cross Clamp Time (min)

Short Name: XClampTm

Definition: Indicate the total number of minutes that the coronary circulation is mechanically isolated from systemic circulation, either by an aortic cross clamp or systemic circulatory arrest.

Intent/Clarification:

Example: For the following two operations: (1) "Transplant, Heart", and (2) "Transplant, Heart and Lung", the field "Cross Clamp Time" will be defined as the cross clamp time of the **donor** heart. Therefore, these two operations represent the only operations where the field "Cross Clamp Time" can be greater than the field "Cardiopulmonary Bypass Time".

SEQ. #: 2440

Long Name: Cardioplegia Delivery

Short Name: CplegiaDeliv

Definition: Indicate the delivery method of cardioplegia if used.

Intent/Clarification:

Cardioplegia is a solution that is used to cause the heart to arrest as documented by the surgeon or perfusionist. Refer to the perfusion record or Surgeon's dictation.

- Non, if not used
 - Antegrade
 - Retrograde
 - Both
-
-

SEQ. #: 2445

Long Name: Cardioplegia Type

Short Name: CplegiaType

Definition: Indicate the type of cardioplegia used.

Intent/Clarification:

Choose one of following:

- **Blood** (If any blood is contained in the solution, any ratio). Includes the following solutions:
 - Combination of blood +St. Thomas solution (i.e.Plegisol)

- DelNido cardioplegia
 - Microplegia
 - **Crystalloid** (If solution is **only** crystalloid)
 - **Both** (If both types of solutions are used) Use “Both” if two different solutions were used during the procedure, 1 with blood and 1 crystalloid
 - **Other**
-
-

SEQ. #: 2450

Long Name: Cerebral Oximetry Used

Short Name: CerOxUsed

Definition: Indicate whether cerebral oximetry was used.

Intent/Clarification:

Cerebral oximetry is similar to pulse oximetry in that it uses differences in light absorption between oxygenated and deoxygenated hemoglobin to measure regional oxygen saturation.

SEQ. #: 2490

Long Name: Diffuse Aortic Calcification (Porcelain Aorta)

Short Name: ConCalc

Definition: Indicate whether diffuse or concentric calcification of the aorta was discovered preoperatively or intraoperatively using imaging or palpation.

Intent/Clarification:

The intent is to capture when and if concentric calcification is discovered. This may impact the surgeon’s approach to cannulation.

Concentric calcification is the same as circumferential calcification and is often described as a porcelain aorta.

SEQ. #: 2495

Long Name: Assessment of Ascending Aorta/Arch

Short Name: AsmtAscAA

Definition: Indicate whether the Ascending Aorta/Arch was evaluated for atheroma or plaque during surgery using TEE or epi-aortic ultrasound. (Not intended for assessment of aneurysmal disease or dissection.)

Intent/Clarification:

Do not capture descending calcification. The intent is to evaluate the area of the aorta that will be cannulated, clamped or otherwise manipulated during the case. Calcification or atheroma in this area can predispose the patient to stroke. Include descriptions of aortic root as ascending calcification.

- Yes
- No
- Not reported

SEQ. #: 2497

Long Name: Method of Assessment of Aorta Plaque

Short Name: AsmtAoDxMeth

Definition: Indicate the method of assessing the highest grade of atheroma or plaque in the ascending aorta.

Intent/Clarification:

Do not capture descending calcification. The intent is to evaluate the area of the aorta that will be cannulated, clamped or otherwise manipulated during the case. Calcification or atheroma in this area can predispose the patient to stroke. Include descriptions of aortic root as ascending calcification.

Indicate the method of assessment: Epiaortic ultrasound is an intraoperative evaluation. TEE can be performed pre and/or intraoperatively. Some patients may have preoperative evaluation by CT scan. MRI/A is another modality.

SEQ. #: 2500

Long Name: Assessment of Aorta Plaque

Short Name: AsmtAoDx

Definition: Indicate highest grade of atheroma or plaque in the ascending aorta.

Intent/Clarification:

Choose one of following:

- Normal Aorta/No or minimal plaque
- Extensive intimal thickening
- Protruding Atheroma < 5 mm
- Protruding Atheroma ≥ 5 mm
- Mobile Plaques
- Not Documented

This will be found intraoperatively in the surgeon's dictation or the anesthesia record.

SEQ. #: 2505

Long Name: Aortic Condition Altered Plan

Short Name: AsmtAPIn

Definition: Indicate whether aortic assessment changed cannulation strategy or surgical plan.

Intent/Clarification:

This assessment can assist the surgeon with selection of optimal site for cannulation of ascending aorta or may prompt decision to select alternate arterial cannulation site or an off pump approach.

SEQ. #: 2510

Long Name: Intraop Blood Products Refused

Short Name: IBldProdRef

Definition: Indicate whether the patient or family refused blood products.

Intent/Clarification: Identify if the patient refused blood or blood products prior to surgery. This may be found in the history and physical, surgical consultation or in a specific consent/refusal form.

Sequence number 2510 is the parent field to sequence number 2515.

SEQ. #: 2515

Long Name: Intraop Blood Products

Short Name: IBldProd

Definition: Indicate whether blood products were transfused any time intraoperatively during the initial surgery. Intraoperatively is defined as any blood started inside of the OR.

Intent/Clarification:

Intraoperatively is defined as any blood started inside of the OR.

For these Intraop Blood Product data fields, the intent is to ONLY collect blood products that were transfused any time intraoperatively during the INITIAL SURGERY. This includes RBCs, FFP, Platelets or Cryoprecipitate.

Sequence number 2515 is the parent field to sequence numbers 2520, 2525, 2530, and 2535.

SEQ. #: 2520

Long Name: Intraop Blood Products - RBC Units

Short Name: IBdRBCU

Definition: Indicate the number of units of packed red blood cells that were transfused intraoperatively. Do not include autologous, cell-saver, pump-residual or chest tube recirculated blood.

Intent/Clarification:

Do not include autologous, cell-saver, pump-residual or chest tube recirculated blood.

SEQ. #: 2525

Long Name: Intraop Blood Products - FFP Units

Short Name: IBdFFPU

Definition: Indicate the number of units of fresh frozen plasma that were transfused intraoperatively.

Intent/Clarification:

This can be found in the EMR, anesthesia or operative record or blood transfusion records.

SEQ. #: 2530

Long Name: Intraop Blood Products - Platelet Units

Short Name: IBdPlatU

Definition: Indicate the number of units of platelets that were transfused intraoperatively.

Count the dose pack as one unit. A dose pack may consist of 4, 6, 8, 10, or any number of donor platelets obtained. The number of units coded is not volume dependent.

Intent/Clarification:

The number of units of platelets transfused during the surgical procedure while the patient was in the OR.

Platelets can be aggregated from several donors or be designated as single donor platelets. It is imperative that each site understand their institution's definition for Random Donor Platelets (RDP) and Single Donor Platelets (SDP).

SDP or Platelet Pheresis count as one unit. One unit is comprised of platelets derived from a single donor. The number of units is not volume dependent.

SEQ. #: 2535

Long Name: Intraop Blood Products - Cryo Units

Short Name: IBdCryoU

Definition: Indicate the number of units of cryoprecipitate that were transfused intraoperatively. One bag of cryo = one unit.

The number of units is not volume dependent.

Intent/Clarification:

This can be found in the EMR, anesthesia or operative record or blood transfusion records.

SEQ. #: 2545

Long Name: Intraop Clotting Factors

Short Name: IntraClotFact

Definition: Indicate whether clotting factors were administered intraoperatively.

Intent/Clarification:

Include clotting factors other than those mentioned above. Other clotting factors may include: Factor VIIa, FEIBA (Anti-Inhibitor Coagulant Complex), or Composite (Platelet-rich Plasma)

- Yes, Factor VIIa
- Yes, FEIBA (Anti-Inhibitor Coagulant Complex)
- Yes, Composite, includes Platelet-rich Plasma
- No

SEQ. #: 2546

Long Name: Intraop Prothrombin Complex Concentrate

Short Name: IntraopProComCon

Definition: Indicate whether prothrombin complex concentrate (i.e.K-Centra) was given intraoperatively.

Intent/Clarification: -

SEQ. #: 2550

Long Name: Intraop Antifibrinolytic Medications - Epsilon Amino-Caproic Acid

Short Name: IMedEACA

Definition: Indicate whether the patient received Epsilon Amino-Caproic Acid in the operating room.

Intent/Clarification: -

SEQ. #: 2555

Long Name: Intraop Antifibrinolytic Medications - Tranexamic Acid

Short Name: IMedTran

Definition: Indicate whether the patient received Tranexamic Acid in the operating room.

Intent/Clarification: -

SEQ. #: 2560

Long Name: Intraop TEE post procedure

Short Name: InOpTEE

Definition: Indicate whether intraoperative TEE was performed following procedure.

Intent/Clarification: Indicate if a transesophageal echocardiogram (TEE) was performed intraoperatively following the procedure after the patient is removed from Cardiopulmonary Bypass prior to OR Exit time.

A TEE is performed by passing a small tube thru the patient's mouth into the esophagus to typically assess the efficiency of the patient's heart valves and ejection fraction (efficiency of the left ventricle). At this point of the surgery it is done to assess the valves and to obtain ejection fraction.

SEQ. #: 2565

Long Name: Post Repair TEE Aortic Insufficiency

Short Name: PRepAR

Definition: Indicate the highest level of aortic insufficiency/ regurgitation found on post CPB intraop TEE. Mild-to-Moderate should be coded as moderate; moderate to severe should be coded as severe. Amount of AR should be the LAST ASSESSMENT before leaving the operating room. For example: if patient has aortic repair, separates from CPB and finds moderate AR, surgeon goes back on and re-fixes, comes off and finds no AR, it should be recorded as none.

Intent/Clarification: Indicate the level of aortic insufficiency obtained by the intraoperative post-procedure TEE prior to the patient leaving the OR after the surgical procedure is complete. Obtain the amount closest to OR Exit Time.

Insufficiency is also called regurgitation (AR) in which the valve does not seal properly and allows too much blood to return to the left ventricle after diastole. This causes an increase in preload of the left ventricle.

Choices include:

- None
- Trivial/Trace
- Mild
- Moderate
- Severe
- Not Documented

If the valve is reported as "normal" code "none"

Choose not documented for the highest level of insufficiency if the test was performed and insufficiency is not reported.

SEQ. #: 2566

Long Name: Aortic Gradient - Post Repair Mean

Short Name: PRepAGradM

Definition: Indicate the mean aortic valve gradient on TEE in the OR after the procedure

Intent/Clarification: Record the mean aortic valve gradient obtained from the TEE intraoperatively post-procedure. Record the one closest to OR Exit Time.

The aortic mean gradient is the mean of the amount of pressure across the aortic valve and should be reported as a pressure in millimeters of Mercury (mmHg) with a typical range of <20 to >50mmHg.

SEQ. #: 2567

Long Name: Post Repair Aortic Paravalvular Leak

Short Name: PRepAPVL

Definition: Indicate whether there was an aortic paravalvular leak noted on TEE in the OR after the procedure

Intent/Clarification: Indicate if any amount of leakage was identified around the aortic valve intraoperatively post-procedure. Obtain the amount closest to OR Exit Time.

Choices are:

- None
- Trivial/Trace
- Mild
- Moderate
- Severe
- Not Documented

FAQ August 2017: Is this field intended to be coded for all valves or just when a new valve is implanted?

Answer: This field should be coded only for current and/or prior prosthetic valves. It can be coded when a new valve is implanted or for a valve that was implanted during a previous operation. DCRI will omit these fields from your DQR when no prosthetic valve is present.

SEQ. #: 2570

Long Name: Post Repair TEE Mitral Insufficiency

Short Name: PRepMR

Definition: Indicate the highest level of mitral insufficiency/ regurgitation found on post CPB intraop TEE. Mild-to-Moderate should be coded as moderate; moderate to severe should be coded as severe. Amount of MR should be the LAST ASSESSMENT before leaving the operating room. For example: if patient has mitral repair, separates from CPB and finds moderate MR, surgeon goes back on and re-fixes, comes off and finds no MR, it should be recorded as none.

Intent/Clarification: Indicate the level of mitral valve insufficiency obtained by the intraoperative post-procedure TEE prior to the patient leaving the OR after the surgical procedure is complete. Obtain the amount closest to OR Exit Time.

Insufficiency is also called mitral regurgitation (MR) in which the valve does not seal properly and allows too much blood to return to the left atrium.

Choices include:

- None
- Trivial/Trace
- Mild
- Moderate
- Severe
- Not Documented

If the valve is reported as “normal” code “none”

Choose not documented for the highest level of insufficiency if the test was performed and insufficiency is not reported.

SEQ. #: 2571

Long Name: Mitral Gradient - Post Repair Mean

Short Name: PRepMGradM

Definition: Indicate the mean mitral valve gradient on TEE in the OR after the procedure

Intent/Clarification: Record the mean mitral valve gradient obtained from the TEE intraoperatively post-procedure. Record the one closest to OR Exit Time.

The mitral mean gradient is the mean of the amount of pressure across the mitral valve and should be reported as a pressure in millimeters of Mercury (mmHg) with a typical range of <5 to >10mmHg.

SEQ. #: 2572

Long Name: Post Repair Mitral Paravalvular Leak

Short Name: PRepMPVL

Definition: Indicate whether there was a mitral paravalvular leak noted on TEE in the OR after the procedure

Intent/Clarification: Indicate if any amount of leakage was identified around the mitral valve intraoperatively post-procedure. Obtain the amount closest to OR Exit Time.

Choices are:

- None
- Trivial/Trace
- Mild
- Moderate
- Severe

- Not Documented

FAQ August 2017: Is this field intended to be coded for all valves or just when a new valve is implanted?

Answer: This field should be coded only for current and/or prosthetic valves. It can be coded when a new valve is implanted or for a valve that was implanted during a previous operation. DCRI will omit these fields from your DQR when no prosthetic valve is present.

SEQ. #: 2575

Long Name: Post Repair TEE Tricuspid Insufficiency

Short Name: PRepTR

Definition: Indicate the highest level of tricuspid insufficiency/regurgitation found on post CPB intraop TEE. Mild-to-Moderate should be coded as moderate; moderate to severe should be coded as severe. Amount of TR should be the LAST ASSESSMENT before leaving the operating room.

Intent/Clarification: Indicate the level of tricuspid valve insufficiency obtained by the intraoperative post-procedure TEE prior to the patient leaving the OR after the surgical procedure is complete. Obtain the amount closest to OR Exit Time.

Insufficiency is also called tricuspid regurgitation (TR) in which the valve does not seal properly and allows too much blood to return to the right atrium.

Choices include:

- None
- Trivial/Trace
- Mild
- Moderate
- Severe
- Not Documented

If the valve is reported as “normal” code “none”

Choose not documented for the highest level of insufficiency if the test was performed and insufficiency is not reported.

SEQ. #: 2576

Long Name: Tricuspid Gradient - Post Repair Mean

Short Name: PRepTGradM

Definition: Indicate the mean tricuspid valve gradient on TEE in the OR after the procedure

Intent/Clarification: Record the mean tricuspid valve gradient obtained from the TEE intraoperatively post-procedure. Record the one closest to OR Exit Time.

The tricuspid mean gradient is the mean of the amount of pressure across the tricuspid valve and should be reported as a pressure in millimeters of Mercury (mmHg) with a typical range of <5 or ≥5.

SEQ. #: 2577

Long Name: Post Repair Tricuspid Paravalvular Leak

Short Name: PRepTPVL

Definition: Indicate whether there was a tricuspid paravalvular leak noted on TEE in the OR after the procedure

Intent/Clarification: Indicate if any amount of leakage was identified around the tricuspid valve intraoperatively post-procedure. Obtain the amount closest to OR Exit Time.

Choices are:

- None
- Trivial/Trace
- Mild
- Moderate
- Severe
- Not Documented

FAQ August 2017: Is this field intended to be coded for all valves or just when a new valve is implanted?

Answer: This field should be coded only for current and/or prior prosthetic valves. It can be coded when a new valve is implanted or for a valve that was implanted during a previous operation. DCRI will omit these fields from your DQR when no prosthetic valve is present.

SEQ. #: 2581

Long Name: Ejection Fraction Measured Post Procedure

Short Name: PPEFMeas

Definition: Indicate whether the ejection fraction was measured after the procedure.

Intent/Clarification: Indicate if an ejection fraction was obtained intraoperatively post-procedure.

Ejection fraction (EF) indicates the efficiency of the left ventricle (ability to pump blood sufficiently to the rest of the body). It compares the amount of blood in the left ventricle at the end of systole (when the ventricle is fuller) to the end of diastole (after the ventricle contracted and should be less full). Issues effecting the left ventricles pumping ability include preload (the amount of blood deposited into the ventricle prior to diastole), afterload (amount of pressure the ventricle has to pump against typically high as a result of elevated systemic venous pressure), ventricular hypertrophy (the enlargement of the ventricle which results in stretching of the ventricle causing decreased contractility and

is a usually a result of congestive heart failure), and valvular insufficiency. Ejection fraction is typically reported in a percentage (1-99%) or described with words.

SEQ. #: 2582

Long Name: Ejection Fraction Post Procedure

Short Name: PPEF

Definition: Indicate the percentage of the blood emptied from the left ventricle at the end of the contraction. Use the most recent determination after the procedure documented on a diagnostic report.

Enter a percentage in the range of 1 - 99. If a percentage range is reported, report a whole number using the "mean" (i.e., 50-55% is reported as 53%).

Note: If no diagnostic report is in the medical record, a value documented in the medical record is acceptable.

ACCF/AHA 2013

Intent/Clarification: Record the mean ejection fraction (EF) closest to OR Exit Time. If a qualitative description is reported, code the mean value for that range; i.e., normal (50-70%) is coded as 60%.

- Hyperdynamic: >70% (code 71%)
- Normal: 50%–70% (midpoint 60%)
- Mild dysfunction: 40%–49% (midpoint 45%)
- Moderate dysfunction: 30%–39% (midpoint 35%)
- Severe dysfunction: <30% (code 29%)

Note: If no diagnostic report is in the medical record, a value documented in the medical record is acceptable.

SEQ. #: 2606

Long Name: Planned Post Procedure PCI

Short Name: PPPlannedPCI

Definition: Indicate whether the procedure was followed by a planned PCI.

Intent/Clarification: Indicate if the patient returned to the Cath Lab any time after OR EXIT Time and before discharge for a percutaneous coronary intervention (PCI) that was planned prior to coronary, valve, or aorta surgery. To be considered “planned” this would need to be indicated in the Medical Provider’s preoperative notes.

A percutaneous coronary intervention (PCI) is understood to be any procedure where entry to the vascular system is obtain thru percutaneous access (a needle poked thru the skin). A catheter is then inserted and a guide wire is thru passed thru the vascular system to the heart. Dye is then injected and pictures of the heart vessels are obtained as the dye flows thru via fluoroscopy (x-ray). An intervention is then performed to “open up” a vessel(s) if a blockage is recognized where dye flow was decreased. This can be done either thru angioplasty (a ballooning of the vessel to allow more blood to flow) or an angioplasty with stent placement (a ballooning of the vessel to allow more blood to

pass followed by placement of a stent to help the vessel to remain open). Either “angioplasty” or “angioplasty with stent placement” should be captured here.

Coronary Bypass

SEQ. #: 2626

Long Name: Internal Mammary Artery Used

Short Name: IMAUsed

Definition: Indicate whether an internal mammary artery conduit was used

Intent/Clarification:

To capture the use of an internal mammary artery to construct one or more distal anastomoses: LIMA, RIMA, both or none. IMA may be used as a free or in-situ graft; pedicle, skeletonized.

FAQ October 2017: Patient had a non-STEMI with urgent CABG. LIMA and RIMA were initially anastomosed to the LAD and Diagonal but prior to leaving the OR, they had insufficient flow and vein grafts were used. The RIMA was a T-graft to the Diagonal from the free LIMA. The surgeon documented “sluggish flow in the LIMA after harvesting as a pedicle, therefore it was utilized as a free graft. He documented this was probably due to stenosis at the takeoff from the left subclavian artery. The chest was closed and initial vitals were stable. Prior to leaving the OR, EKG and hemodynamic changes were noted. Therefore, the chest was reopened and the patient was placed back on CPB. The flow via the IMA grafts were poor so the surgeon removed them and used venous grafts to the LAD and Diagonal.” How do I code IMA used?

Answer: Code no, an IMA was not used.

SEQ. #: 2627

Long Name: Reason for No IMA

Short Name: NoIMARsn

Definition: Indicate PRIMARY reason Internal Mammary artery was not used as documented in medical record.

Intent/Clarification:

Choose from the following reasons:

- Subclavian stenosis*
- Previous cardiac or thoracic surgery
- Previous mediastinal radiation
- Emergent or salvage procedure
- No (BYPASSABLE) LAD disease - This can include a clean LAD, diffusely diseased LAD or other condition resulting in the LAD not being bypassed.

- Other – The National Quality Forum (NQF) does not consider this exclusion for measure purposes.

Other is not an acceptable exclusion in the NQF endorsed measure and will have a negative impact on the star rating.

FAQ October 2017: Patient had a non-STEMI with urgent CABG. LIMA and RIMA were initially anastomosed to the LAD and Diagonal but prior to leaving the OR, they had insufficient flow and vein grafts were used. The RIMA was a T-graft to the Diagonal from the free LIMA. The surgeon documented sluggish flow in the LIMA after harvesting as a pedicle, therefore it was utilized as a free graft. He documented this was probably due to stenosis at the takeoff from the left subclavian artery. The chest was closed and initial vitals were stable. Prior to leaving the OR, EKG and hemodynamic changes were noted. Therefore, the chest was reopened and the patient was placed back on CPB. The flow via the IMA grafts were poor so the surgeon removed them and used venous grafts to the LAD and Diagonal. If I must code “no” IMA used, what reason for no IMA should I use?

Answer: Code ‘Other’. Subclavian stenosis cannot be considered an exclusion because the IMA was initially used as a free graft.

SEQ. #: 2628

Long Name: IMA Dist Anast #

Short Name: NumIMADA

Definition: Indicate the total number of distal anastomoses done using IMA grafts.

Intent/Clarification:

To collect the total number of anastomoses constructed using an IMA conduit. More than one anastomosis can be constructed from each IMA; the IMA may be used as a pedicle graft or a free graft. A pedicle graft remains connected at its proximal origin and requires only a distal anastomosis.

SEQ. #: 2629

Long Name: Left IMA Used

Short Name: LeftIMA

Definition: Indicate whether the left internal mammary was used

Intent/Clarification:

The left IMA was used to construct one or more anastomosis; pedicle or skeletonized.

SEQ. #: 2630

Long Name: Left IMA Harvest Technique

Short Name: LIMAHarvTech

Definition: Indicate the harvest technique used for the left internal mammary

Intent/Clarification:

Indicate the technique used to harvest an IMA:

- **Direct vision** (open) - Standard method; through full or partial sternotomy. IMA harvest with the chest open using a standard retractor.
- **Thoracoscopy** - Endoscopy used for the entire IMA harvest.
- **Combination** - Both thoracoscopy and direct vision used for IMA harvest.
- **Robotic assist** - Robot was used to harvest IMA.

SEQ. #: 2631

Long Name: Right IMA Used

Short Name: RightIMA

Definition: Indicate whether the right internal mammary was used

Intent/Clarification:

The right IMA was used to construct one or more anastomosis; pedicle or skeletonized.

SEQ. #: 2632

Long Name: Right IMA Harvest Technique

Short Name: RIMAHarvTech

Definition: Indicate the harvest technique used for the right internal mammary

Intent/Clarification:

Indicate the technique used to harvest an IMA:

- **Direct vision** (open) -Standard method; through full or partial sternotomy. IMA harvest with the chest open using a standard retractor.
- **Thoracoscopy** - Endoscopy used for the entire IMA harvest.
- **Combination** - Both Thoracoscopy and direct vision used for IMA harvest.
- **Robotic assist** - Robot was used to harvest IMA.

SEQ. #: 2633

Long Name: Radial Artery Used

Short Name: RadialArtUsed

Definition: Indicate whether a radial artery conduit was used.

Intent/Clarification:

The radial artery was used to construct one or more anastomosis.

SEQ. #: 2634

Long Name: Radial Dist Anast #

Short Name: NumRadDA

Definition: Indicate the total number of distal anastomoses done using radial artery grafts.

Intent/Clarification:

To collect the total number of distal anastomoses constructed using a radial artery. More than one anastomosis can be constructed from each radial artery.

SEQ. #: 2635

Long Name: Radial Dist Anast Harvest Technique

Short Name: RadHTech

Definition: Indicate the technique used to harvest the radial artery(s).

Intent/Clarification:

The technique used to harvest the radial artery (ies):

- Endoscopic
 - Direct vision (open) - Standard method; through full or partial radial harvest
 - Both - Both endovascular and direct vision used for radial artery harvest
-
-

SEQ. #: 2636

Long Name: Radial Artery Harvest and Preparation Time

Short Name: RadHarvPrepTm

Definition: Indicate the total time for radial artery harvest and preparation.

Intent/Clarification:

It is important to quantify the harvest and prep times to track resource utilization and provide objective data for RUC (Specialty Society Relative Value Scale Update Committee or Relative Value Update Committee, an American Medical Association group involved in health care pricing) surveys and coding. This is important because these values determine the rate at which Medicare and other payers reimburse for procedures.

SEQ. #: 2637

Long Name: Venous Conduit(s) Used

Short Name: VenousCondUsed

Definition: Indicate whether a venous conduit was used

Intent/Clarification:

A venous conduit was used to construct one or more anastomosis.

SEQ. #: 2638

Long Name: Dist Anast - Vein #

Short Name: DistVein

Definition: Indicate the total number of distal anastomoses with venous conduits.

Intent/Clarification:

Distal anastomosis refers to the connection between the bypass graft (conduit) and coronary artery. Record the total number of venous anastomoses constructed using a venous conduit connection to a coronary artery. More than one anastomosis can be constructed from a single vein. Saphenous veins are used as free grafts to bypass any coronary artery.

SEQ. #: 2639

Long Name: Dist Anast - Vein Harvest Technique

Short Name: DistVeinHTech

Definition: Indicate the technique used to harvest the vein graft(s).

Intent/Clarification:

The technique(s) used to harvest the vein grafts:

- Endoscopic
 - Direct vision (open) - Through full or partial vein harvest
 - Both - Both endoscopic and direct vision used to harvest the vein grafts
 - Cryopreserved - Cryopreserved veins harvested from a donor, typically commercially supplied
-
-

SEQ. #: 2640

Long Name: Saphenous Vein Harvest And Preparation Time

Short Name: SaphHarPrepTm

Definition: Indicate the total time for saphenous vein harvest and preparation.

Intent/Clarification:

It is important to quantify the harvest and prep times to track resource utilization and provide objective data for RUC, (Specialty Society Relative Value Scale Update Committee or Relative Value Update Committee, an American Medical Association group involved in health care pricing) surveys and coding. This is important because

these values determine the rate at which Medicare and other payers reimburse for procedures.

SEQ. #: 2641

Long Name: Other Arterial Distal Anastomoses #

Short Name: NumOArtD

Definition: Indicate the number of arterial distal anastomoses that were used, other than radial or IMA.

Intent/Clarification:

Any other arterial conduit was used to construct one or more anastomosis; i.e. inferior epigastric artery.

SEQ. #: 2650

Long Name: Number Of Distal Anastomoses With Arterial-Venous Composite Conduits

Short Name: NumArtVenComp

Definition: Indicate the number of distal anastomoses with arterial-venous composite conduits

Intent/Clarification:

To capture grafts constructed from artery and venous composite. The venous component of the composite graft is anastomosed to the coronary artery. (i.e. an IMA or radial artery is lengthened with a segment of saphenous vein). The vein segment is attached to the coronary artery. Alternatively, a composite graft can be constructed as a “Y” or “T” with one limb going to one coronary artery and the other limb going to a different coronary site. The **arterial** segment provides inflow. When part of the IMA was used as part of the composite graft, code 2626 yes as the internal mammary artery was used. Composite grafts that include an IMA meet the NQF measure for IMA use.

SEQ. #: 2651

Long Name: Number Of Distal Anastomoses With Venous-Arterial Composite Conduits

Short Name: NumVenArtComp

Definition: Indicate the number of distal anastomoses with venous-arterial composite conduits

Intent/Clarification:

To capture the number of grafts (if any) constructed from venous and arterial composite. The arterial component of the composite graft is anastomosed to the coronary artery. i.e., the IMA or radial artery which has been anastomosed to the coronary artery is too short to reach the aorta and is lengthened with a segment of saphenous vein. Alternatively, a composite graft can be constructed as a “Y” or “T” with one limb going to one coronary artery and the other limb going to a different coronary site. The **venous** segment provides inflow. When part of the IMA was used as part of the composite graft,

code 2626 yes as the internal mammary artery was used. Composite grafts that include an IMA meet the NQF measure for IMA use.

SEQ. #: 2652

Long Name: Number Of Distal Anastomoses With Arterial-Arterial Composite Conduits

Short Name: NumArtArtComp

Definition: Indicate the number of distal anastomoses with arterial-arterial composite conduits

Intent/Clarification:

To capture the number of grafts (if any) constructed from arterial and arterial composite; i.e. IMA plus radial artery. The concepts are similar to seq# 2651 and 2652, except the composite is composed of arteries only. When part of the IMA was used as part of the composite graft, code 2626 yes as the internal mammary artery was used. Composite grafts that include an IMA meet the NQF measure for IMA use.

SEQ. #: 2710

Long Name: Proximal Technique

Short Name: ProxTech

Definition: Indicate the technique employed for proximal graft anastomosis.

Intent/Clarification:

The intent is to determine various methods used to perform proximal anastomosis which may have an impact on the risk of stroke/ embolization from aortic intima. If more than one technique was used for proximal grafts, choose the highest level of occlusion used.

- Single Cross Clamp
 - Partial Occlusion Clamp
 - Anastomotic Assist Device – such as Cyclone, Enclose, Cardica Passport, Heart String, etc.
 - None (isolated in-situ mammary)
-
-

SEQ. #: 2730

Long Name: CAB Distal Site 01

Short Name: CABDistSite01

Definition: Indicate distal insertion site of bypass.

Intent/Clarification:

- These order does not matter, include up to 10 grafts. One graft = one distal insertion.
- Left Main - Left Main
- Prox LAD - Proximal Left Anterior Descending

- Mid LAD - Middle Left Anterior Descending
- Distal LAD - Distal Left Anterior Descending
- Diagonal 1 - First Diagonal
- Diagonal 2 - Second Diagonal
- Diagonal 3 - Third Diagonal
- Circumflex - Circumflex
- Obtuse Marginal 1 - First Obtuse Marginal
- Obtuse Marginal 2 - Second Obtuse Marginal
- Obtuse Marginal 3 - Third Obtuse Marginal
- Ramus - Ramus Intermedius
- RCA - Right Coronary Artery
- Acute Marginal (AM) - Acute Marginal
- Posterior Descending (PDA) - Posterior Descending Artery
- Posterolateral (PLB) - Posterolateral Branch
- Other - Any other site

FAQ September 2017: Pt with previous CAB in January of this year now presents with Aortic Dissection. The dissection was repaired and cadaver vein graft extensions were used to reimplant the PDA/PL grafts. The OM and Diag grafts were occluded by the dissection. How is this captured in the CAB grid?

Answer: Do not complete the CAB grid, a distal coronary artery graft is not constructed using the cadaver vein.

SEQ. #: 2740

Long Name: CAB Proximal Site 01

Short Name: CABProximalSite01

Definition: Indicate proximal site of the bypass graft.

Intent/Clarification:

- In Situ Mammary
- Ascending aorta
- Descending aorta
- Subclavian artery
- Innominate artery
- T-graft off SVG
- T-graft off Radial
- T-graft off LIMA
- T-graft off RIMA
- Natural Y vein graft
- Other

SEQ. #: 2750

Long Name: CAB Conduit 01

Short Name: CABConduit01

Definition: Indicate the conduit type used.

Intent/Clarification:

- Vein graft
- In Situ LIMA
- In Situ RIMA
- Free IMA
- Radial artery
- Other arteries, homograft
- Synthetic graft
- Composite artery-vein

FAQ August 2017: What is a composite artery vein?

Answer: A composite is an extension of the IMA with a vein to allow the reach its intended distal coronary target. Using composite does not exclude the IMA from the composite measure for IMA usage.

SEQ. #: 2755

Long Name: CAB Distal Position 01

Short Name: CABDistPos01

Definition: Indicate anastomotic position.

Intent/Clarification:

End to side: the end of the graft is inserted into the side of the target vessel

• **Sequential (side to side):** sometimes called a jump graft, the side of the graft is inserted into the side of the target vessel and the end of the graft is inserted elsewhere on that vessel or on another target vessel.

SEQ. #: 2760

Long Name: CAB Endarterectomy 01

Short Name: CABEndArt01

Definition: Indicate whether endarterectomy was performed.

Intent/Clarification:

Endarterectomy is a surgical procedure to remove the atheromatous plaque material, or blockage, in the lining of an artery constricted by the buildup of soft/hardening deposits. It is carried out by separating (peeling) the plaque from the arterial wall.

SEQ. #: 2765

Long Name: CAB Vein Patch Angioplasty 01

Short Name: CABVeinPatAng01

Definition: Indicate whether a vein/**arterial** patch angioplasty was performed.

Intent/Clarification:

A patch constructed from a piece of vein, suturing the vein around the arteriotomy to reconstruct the coronary artery without narrowing it. . A bypass graft is then often, but not always, placed into the vein patch reconstruction to provide new inflow.

NOTE: this explanatory note is valid for all subsequent fields related to CAB vein patch angioplasty

FAQ October 2017: The surgeon completed a patch angioplasty using the radial artery instead of a vein, how is this coded?

Answer: Code as vein patch angioplasty. This can apply to either vein or arterial patch.

SEQ. #: 2770

Long Name: CAB 02 **Short Name:** CAB02

Definition: Indicate whether a second Coronary Artery Bypass graft was done.

FAQ October 2017: A patient has a four vessel bypass. During the procedure he also has an endarterectomy and vein patch angioplasty of the circumflex (no bypass). Do I code that a coronary bypass graft was done? Do I capture this as Other Cardiac thus throwing this out of the isolated CABG category?

Answer: In order to open the child fields in the coronary bypass grid to complete the fields for endarterectomy you must code 'Yes'. Leave proximal site blank. Do not code as 'Other Cardiac'.

SEQ. #: 2790

Long Name: CAB Distal Site 02

Short Name: CABDistSite02

Definition: Indicate distal insertion site of bypass.

FAQ October 2017: A patient has a four vessel bypass. During the procedure he also has an endarterectomy and vein patch angioplasty of the circumflex (no bypass). How do I capture the distal insertion site?

Answer: Code distal insertion site circumflex.

SEQ. #: 2800

Long Name: CAB Proximal Site 02

Short Name: CABProximalSite02

Definition: Indicate proximal site of the bypass graft.

FAQ October 2017: A patient has a four vessel bypass. During the procedure he also has an endarterectomy and vein patch angioplasty of the circumflex (no bypass). How do I capture the proximal site?

Answer: Leave proximal site blank.

SEQ. #: 2810

Long Name: CAB Conduit 02

Short Name: CABConduit02

Definition: Indicate the conduit type used.

FAQ October 2017: A patient has a four vessel bypass. During the procedure he also has an endarterectomy and vein patch angioplasty of the circumflex (no bypass). How do I capture conduit type?

Answer: Leave conduit blank.

SEQ. #: 2815

Long Name: CAB Distal Position 02

Short Name: CABDistPos02

Definition: Indicate anastomotic position.

FAQ October 2017: A patient has a four vessel bypass. During the procedure he also has an endarterectomy and vein patch angioplasty of the circumflex (no bypass). How do I capture anastomotic position?

Answer: Leave anastomotic position blank.

SEQ. #: 2820

Long Name: CAB Endarterectomy 02

Short Name: CABEndArt02

Definition: Indicate whether endarterectomy was performed.

FAQ October 2017: A patient has a four vessel bypass. During the procedure he also has an endarterectomy and vein patch angioplasty of the circumflex (no bypass). How do I capture endarterectomy performed?

Answer: Code yes, an endarterectomy was performed.

SEQ. #: 2825

Long Name: CAB Vein Patch Angioplasty 02

Short Name: CABVeinPatAng02

Definition: Indicate whether a vein/**arterial** patch angioplasty was performed.

FAQ October 2017: A patient has a four vessel bypass. During the procedure he also has an endarterectomy and vein patch angioplasty of the circumflex (no bypass). How do I capture vein/arterial patch angioplasty?

Answer: Code yes, a vein patch angioplasty was constructed.

FAQ October 2017: The surgeon completed a patch angioplasty using the radial artery instead of a vein, how is this coded?

Answer: Code as vein patch angioplasty. This can apply to either vein or arterial patch.

SEQ. #: 2830

Long Name: CAB 03

Short Name: CAB03

Definition: Indicate whether a third Coronary Artery Bypass graft was done.

SEQ. #: 2850

Long Name: CAB Distal Site 03

Short Name: CABDistSite03

Definition: Indicate distal insertion site of bypass.

SEQ. #: 2860

Long Name: CAB Proximal Site 03

Short Name: CABProximalSite03

Definition: Indicate proximal site of the bypass graft.

SEQ. #: 2870

Long Name: CAB Conduit 03

Short Name: CABConduit03

Definition: Indicate the conduit type used.

SEQ. #: 2875

Long Name: CAB Distal Position 03

Short Name: CABDistPos03

Definition: Indicate anastomotic position.

SEQ. #: 2880

Long Name: CAB Endarterectomy 03

Short Name: CABEndArt03

Definition: Indicate whether endarterectomy was performed.

SEQ. #: 2885

Long Name: CAB Vein Patch Angioplasty 03

Short Name: CABVeinPatAng03

Definition: Indicate whether a vein/**arterial** patch angioplasty was performed.

FAQ October 2017: The surgeon completed a patch angioplasty using the radial artery instead of a vein, how is this coded?

Answer: Code as vein patch angioplasty. This can apply to either vein or arterial patch.

SEQ. #: 2890

Long Name: CAB 04

Short Name: CAB04

Definition: Indicate whether a fourth Coronary Artery Bypass graft was done.

SEQ. #: 2910

Long Name: CAB Distal Site 04

Short Name: CABDistSite04

Definition: Indicate distal insertion site of bypass.

SEQ. #: 2920

Long Name: CAB Proximal Site 04

Short Name: CABProximalSite04

Definition: Indicate proximal site of the bypass graft.

SEQ. #: 2930

Long Name: CAB Conduit 04

Short Name: CABConduit04

Definition: Indicate the conduit type used.

SEQ. #: 2935
Long Name: CAB Distal Position 04
Short Name: CABDistPos04
Definition: Indicate anastomotic position.

SEQ. #: 2940
Long Name: CAB Endarterectomy 04
Short Name: CABEndArt04
Definition: Indicate whether endarterectomy was performed.

SEQ. #: 2945
Long Name: CAB Vein Patch Angioplasty 04
Short Name: CABVeinPatAng04
Definition: Indicate whether a vein/**arterial** patch angioplasty was performed.

FAQ October 2017: The surgeon completed a patch angioplasty using the radial artery instead of a vein, how is this coded?
Answer: Code as vein patch angioplasty. This can apply to either vein or arterial patch.

SEQ. #: 2950
Long Name: CAB 05
Short Name: CAB05
Definition: Indicate whether a fifth Coronary Artery Bypass graft was done.

SEQ. #: 2970
Long Name: CAB Distal Site 05
Short Name: CABDistSite05
Definition: Indicate distal insertion site of bypass.

SEQ. #: 2980
Long Name: CAB Proximal Site 05
Short Name: CABProximalSite05
Definition: Indicate proximal site of the bypass graft.

SEQ. #: 2990
Long Name: CAB Conduit 05
Short Name: CABConduit05
Definition: Indicate the conduit type used.

SEQ. #: 2995
Long Name: CAB Distal Position 05
Short Name: CABDistPos05
Definition: Indicate anastomotic position.

SEQ. #: 3000
Long Name: CAB Endarterectomy 05
Short Name: CABEndArt05
Definition: Indicate whether endarterectomy was performed.

SEQ. #: 3005
Long Name: CAB Vein Patch Angioplasty 05
Short Name: CABVeinPatAng05
Definition: Indicate whether a vein/**arterial** patch angioplasty was performed.

FAQ October 2017: The surgeon completed a patch angioplasty using the radial artery instead of a vein, how is this coded?

Answer: Code as vein patch angioplasty. This can apply to either vein or arterial patch.

SEQ. #: 3010
Long Name: CAB 06
Short Name: CAB06
Definition: Indicate whether a sixth Coronary Artery Bypass graft was done.

SEQ. #: 3030
Long Name: CAB Distal Site 06
Short Name: CABDistSite06
Definition: Indicate distal insertion site of bypass.

SEQ. #: 3040
Long Name: CAB Proximal Site 06
Short Name: CABProximalSite06
Definition: Indicate proximal site of the bypass graft.

SEQ. #: 3050
Long Name: CAB Conduit 06
Short Name: CABConduit06
Definition: Indicate the conduit type used.

SEQ. #: 3055
Long Name: CAB Distal Position 06
Short Name: CABDistPos06
Definition: Indicate anastomotic position.

SEQ. #: 3060
Long Name: CAB Endarterectomy 06
Short Name: CABEndArt06
Definition: Indicate whether endarterectomy was performed.

SEQ. #: 3065
Long Name: CAB Vein Patch Angioplasty 06
Short Name: CABVeinPatAng06
Definition: Indicate whether a vein/**arterial** patch angioplasty was performed.

FAQ October 2017: The surgeon completed a patch angioplasty using the radial artery instead of a vein, how is this coded?

Answer: Code as vein patch angioplasty. This can apply to either vein or arterial patch.

SEQ. #: 3070
Long Name: CAB 07
Short Name: CAB07
Definition: Indicate whether a seventh Coronary Artery Bypass graft was done.

SEQ. #: 3090
Long Name: CAB Distal Site 07
Short Name: CABDistSite07
Definition: Indicate distal insertion site of bypass.

SEQ. #: 3100
Long Name: CAB Proximal Site 07
Short Name: CABProximalSite07
Definition: Indicate proximal site of the bypass graft.

SEQ. #: 3110
Long Name: CAB Conduit 07
Short Name: CABConduit07
Definition: Indicate the conduit type used.

SEQ. #: 3115
Long Name: CAB Distal Position 07
Short Name: CABDistPos07
Definition: Indicate anastomotic position.

SEQ. #: 3120
Long Name: CAB Endarterectomy 07
Short Name: CABEndArt07
Definition: Indicate whether endarterectomy was performed.

SEQ. #: 3125
Long Name: CAB Vein Patch Angioplasty 07
Short Name: CABVeinPatAng07
Definition: Indicate whether a vein/**arterial** patch angioplasty was performed.

FAQ October 2017: The surgeon completed a patch angioplasty using the radial artery instead of a vein, how is this coded?

Answer: Code as vein patch angioplasty. This can apply to either vein or arterial patch.

SEQ. #: 3130

Long Name: CAB 08

Short Name: CAB08

Definition: Indicate whether an eighth Coronary Artery Bypass graft was done.

SEQ. #: 3150

Long Name: CAB Distal Site 08

Short Name: CABDistSite08

Definition: Indicate distal insertion site of bypass.

SEQ. #: 3160

Long Name: CAB Proximal Site 08

Short Name: CABProximalSite08

Definition: Indicate proximal site of the bypass graft.

SEQ. #: 3170

Long Name: CAB Conduit 08

Short Name: CABConduit08

Definition: Indicate the conduit type used.

SEQ. #: 3175

Long Name: CAB Distal Position 08

Short Name: CABDistPos08

Definition: Indicate anastomotic position.

SEQ. #: 3180

Long Name: CAB Endarterectomy 08

Short Name: CABEndArt08

Definition: Indicate whether endarterectomy was performed.

SEQ. #: 3185

Long Name: CAB Vein Patch Angioplasty 08

Short Name: CABVeinPatAng08

Definition: Indicate whether a vein/**arterial** patch angioplasty was performed.

FAQ October 2017: The surgeon completed a patch angioplasty using the radial artery instead of a vein, how is this coded?

Answer: Code as vein patch angioplasty. This can apply to either vein or arterial patch.

SEQ. #: 3190

Long Name: CAB 09

Short Name: CAB09

Definition: Indicate whether a ninth Coronary Artery Bypass graft was done.

SEQ. #: 3210

Long Name: CAB Distal Site 09

Short Name: CABDistSite09

Definition: Indicate distal insertion site of bypass.

SEQ. #: 3220

Long Name: CAB Proximal Site 09

Short Name: CABProximalSite09

Definition: Indicate proximal site of the bypass graft.

SEQ. #: 3230

Long Name: CAB Conduit 09

Short Name: CABConduit09

Definition: Indicate the conduit type used.

SEQ. #: 3235

Long Name: CAB Distal Position 09

Short Name: CABDistPos09

Definition: Indicate anastomotic position.

SEQ. #: 3240

Long Name: CAB Endarterectomy 09

Short Name: CABEndArt09

Definition: Indicate whether endarterectomy was performed.

SEQ. #: 3245

Long Name: CAB Vein Patch Angioplasty 09

Short Name: CABVeinPatAng09

Definition: Indicate whether a vein/**arterial** patch angioplasty was performed.

FAQ October 2017: The surgeon completed a patch angioplasty using the radial artery instead of a vein, how is this coded?

Answer: Code as vein patch angioplasty. This can apply to either vein or arterial patch.

SEQ. #: 3250

Long Name: CAB 10

Short Name: CAB10

Definition: Indicate whether a tenth Coronary Artery Bypass graft was done.

SEQ. #: 3270

Long Name: CAB Distal Site 10

Short Name: CABDistSite10

Definition: Indicate distal insertion site of bypass.

SEQ. #: 3280

Long Name: CAB Proximal Site 10

Short Name: CABProximalSite10

Definition: Indicate proximal site of the bypass graft.

SEQ. #: 3290

Long Name: CAB Conduit 10

Short Name: CABConduit10

Definition: Indicate the conduit type used.

SEQ. #: 3295

Long Name: CAB Distal Position 10
Short Name: CABDistPos10
Definition: Indicate anastomotic position.

SEQ. #: 3300
Long Name: CAB Endarterectomy 10
Short Name: CABEndArt10
Definition: Indicate whether endarterectomy was performed.

SEQ. #: 3305
Long Name: CAB Vein Patch Angioplasty 10
Short Name: CABVeinPatAng10
Definition: Indicate whether a vein/**arterial** patch angioplasty was performed.

FAQ October 2017: The surgeon completed a patch angioplasty using the radial artery instead of a vein, how is this coded?

Answer: Code as vein patch angioplasty. This can apply to either vein or arterial patch.

Valve Surgery

SEQ. #: 3310
Long Name: Valve Prosthesis Explant
Short Name: ValExp
Definition: Indicate whether a prosthetic valve or annuloplasty was explanted during this procedure.

Intent/Clarification:

The intent is to capture as much information as possible about explanted devices. This will assist with post market device surveillance and provide information on device longevity. Having this information will help surgeons and patients make informed decisions on device selection.

Code the valve explant even if the sewing cuff is retained.

Do not code a valve explant if a valve is implanted and explanted during the same operation due to the fact the valve did not work or fit.

SEQ. #: 3315

Long Name: Valve Prosthesis Explant Position

Short Name: ValExpPos

Definition: Indicate the location of the first explanted prosthetic valve or annuloplasty device.

Intent/Clarification:

- Aortic
 - Mitral
 - Tricuspid
 - Pulmonic
-
-

SEQ. #: 3320

Long Name: Valve Explant Type

Short Name: ValExpTyp

Definition: Indicate the first type of valve device explanted or enter unknown.

Intent/Clarification:

- Mechanical Valve
 - Leaflet clip
 - Bioprosthetic Valve
 - Transcatheter Device
 - Homograft
 - Other
 - Annuloplasty Device
 - Unknown
-
-

SEQ. #: 3325

Long Name: Valve Explant Etiology

Short Name: ValExpEt

Definition: Indicate the primary reason for explanting valve device.

Intent/Clarification:

Choose the most critical reason that the patient is having the valve explanted.

- Endocarditis
- Failed repair
- Hemolysis: Valve causes destruction of red blood cells.
- Incompetence
- Pannus: Mobility of the leaflets obstructed or impaired by a membrane of tissue.
- Para-valvular leak: Leak around the valve
- Prosthetic deterioration
- Sizing/positioning issue: Valve size or position is suboptimal
- Stenosis
- Thrombosis
- Other

- Unknown

When coding the replacement of a calcified homograft code prosthetic deterioration.

SEQ. #: 3330

Long Name: Valve Explant Device Known

Short Name: ValExpDevKnown

Definition: Indicate whether the type of explanted valve device is known.

Intent/Clarification:

Information is available to identify the explanted valve device. This may include the patient's device card from the manufacturer.

SEQ. #: 3335

Long Name: Valve Explant Device

Short Name: ValExpDev

Definition: Indicate the model number of the first prosthesis explanted.

Intent/Clarification:

Choose the device type from the device list.

SEQ. #: 3340

Long Name: Valve Explant Unique Device Identifier (UDI)

Short Name: ValExpUDI

Definition: Indicate the device UDI if available, otherwise leave blank.

Intent/Clarification:

This is a unique identifier that will be on each valve. It may not be available immediately. If not available leave blank. Please refer to the FDA web site for detailed explanation of UDI.

https://www.fda.gov/MedicalDevices/DeviceRegulationandGuidance/UniqueDeviceIdentification/default.htm?utm_source=Members%E2%80%90Only+Updates&utm_campaign=c7c1e8c870%E2%80%90%20Proposed_Rules_7_5_2012&utm_medium=email

SEQ. #: 3350

Long Name: Second Valve Prosthesis Explant

Short Name: ValExp2

Definition: Indicate whether a second prosthetic valve or annuloplasty was explanted during this procedure.

Intent/Clarification:

In the event that more than one device is explanted, capture both. Code the valve explanted even if the sewing cuff is retained. Do not code a valve explant if a valve is implanted and explanted during the same operation due to the fact the valve did not work or fit.

SEQ. #: 3355

Long Name: Second Valve Prosthesis Explant Position

Short Name: ValExpPos2

Definition: Indicate the location of the second explanted prosthetic valve or annuloplasty.

Intent/Clarification:

- Aortic
 - Mitral
 - Tricuspid
 - Pulmonic
-
-

SEQ. #: 3360

Long Name: Second Valve Explant Type

Short Name: ValExpTyp2

Definition: Indicate the second type of valve device explanted or enter unknown.

Intent/Clarification:

- Mechanical Valve
 - Leaflet clip
 - Bioprosthetic Valve
 - Transcatheter Device
 - Homograft
 - Other
 - Annuloplasty Device
 - Unknown
-
-

SEQ. #: 3365

Long Name: Second Valve Explant Etiology

Short Name: ValExpEt2

Definition: Indicate the primary reason for explanting valve device.

Intent/Clarification:

Choose the most critical reason that the patient had their valve replaced.

- Endocarditis
- Failed repair
- Hemolysis: Valve causes destruction of red blood cells.
- Incompetence
- Pannus: Mobility of the leaflets obstructed or impaired by a membrane of tissue.
- Para-valvular leak: Leak around the valve
- Prosthetic deterioration
- Sizing/positioning issue: Valve size or position is suboptimal
- Stenosis
- Thrombosis
- Other
- Unknown

SEQ. #: 3370

Long Name: Second Valve Explant Device Known

Short Name: ValExpDevKnown2

Definition: Indicate whether the type of explanted valve device is known.

Intent/Clarification:

Information is available to identify the explanted valve device.

SEQ. #: 3375

Long Name: Second Valve Explant Device

Short Name: ValExpDev2

Definition: Indicate the model number of the second prosthesis explanted.

Intent/Clarification:

Choose the device type from the device list.

SEQ. #: 3380

Long Name: Second Valve Explant Device Unique Device Identifier (UDI)

Short Name: ValExpDevUDI

Definition: Indicate the device UDI if available, otherwise leave blank.

Intent/Clarification:

This is a unique identifier that will be on each valve. It may not be available immediately. If not available leave blank. Please refer to the FDA web site for detailed explanation of UDI.

https://www.fda.gov/MedicalDevices/DeviceRegulationandGuidance/UniqueDeviceIdentification/default.htm?utm_source=Members%E2%80%90Only+Updates&utm_campaign=c7c1e8c870%E2%80%90%20Proposed_Rules_7_5_2012&utm_medium=email

SEQ. #: 3390

Long Name: VS-Aortic Valve

Short Name: VSAV

Definition: Indicate whether an aortic valve procedure was performed.

Intent/Clarification:

- Yes, planned
 - Yes, unplanned due to surgical complication
 - Yes, unplanned due to unsuspected disease or anatomy
 - No
-
-

SEQ. #: 3395

Long Name: VS-Aortic Valve Procedure

Short Name: VSAVPr

Definition: Indicate the type of procedure that was performed on the aortic valve and/or ascending aorta.

Intent/Clarification:

Options include:

- Replacement
- Repair / Reconstruction
- ~~Root replacement with valved conduit (Bentall) (Remove)~~
- ~~Replacement AV and insertion aortic non-valved conduit in supra-coronary position (Remove)~~
- ~~Replacement AV and major root reconstruction/debridement with valved conduit (Remove)~~
- ~~Resuspension AV without replacement of ascending aorta (Remove)~~
- ~~Resuspension AV with replacement of ascending aorta (Remove)~~
- ~~Apico-aortic conduit (Aortic valve bypass) (Remove)~~
- ~~Autograft with pulmonary valve (Ross procedure) (Remove)~~
- ~~Homograft Root Replacement (Remove)~~
- ~~Valve sparing root reimplantation (David) (Remove)~~
- ~~Valve sparing root remodeling (Yacoub) (Remove)~~
- ~~Valve Sparing root reconstruction (Florida Sleeve) (Remove)~~

FAQ September 2017: Surgeon performed the following mitral valve procedure “anterior mitral leaflet endarterectomy/decalcification” done in conjunction with an Aortic Valve Replacement. How is this documented under the options provided for MV repair?

Answer: No, anterior mitral leaflet endarterectomy/decalcification is considered part of the AVR and should not be coded as a mitral valve procedure.

SEQ. #: 3400

Long Name: VS-Aortic Transcatheter Valve Replacement

Short Name: VSTCV

Definition: Indicate whether the aortic valve replacement was done using a transcatheter valve device.

Intent/Clarification:

Transcatheter Aortic Valve Replacement (TAVR) technology is designed to allow some patients, who may not be candidates for conventional open-heart valve replacement surgery due to excessive risk, to obtain a life-saving valve.

Catheter based access is obtained through an artery.

If you participate in the TVT registry you may opt to submit transcatheter cases to the STS adult cardiac surgery registry in addition to the TVT registry, but it is not required.

SEQ. #: 3405

Long Name: VS-Transcatheter Valve Replacement Approach

Short Name: VSTCVR

Definition: Indicate transcatheter valve replacement approach.

Intent/Clarification:

TAVR devices may be implanted via multiple vascular approaches:

- Transapical
 - Transaxillary
 - Transfemoral
 - Transaortic
 - Subclavian
 - Other
-
-

SEQ. #: 3407

Long Name: VS-Aortic Surgical Valve Replacement

Short Name: VSAVSurgRep

Definition: Indicate whether the aortic valve replacement was done using a surgical procedure.

Intent/Clarification:

An open surgical valve procedure was performed.

SEQ. #: 3408**Long Name:** VS-Aortic Surgical Valve Replacement Device Type**Short Name:** VSAVSurgType**Definition:** Indicate the type of device used to surgically replace the aortic valve.**Intent/Clarification:**

Choose the device type:

- Mechanical
 - Bioprosthetic
 - Surgeon fashioned pericardium (Ozaki)
 - Other
-
-

SEQ. #: 3409**Long Name:** VS-Aortic Surgical Bioprosthetic Replacement Valve Type**Short Name:** VSAVSurgBioT**Definition:** Indicate the type of bioprosthetic device used to surgically replace the aortic valve.**Intent/Clarification:**

If bioprosthetic, choose valve type:

- Stented
 - Stentless subcoronary valve only
 - Sutureless/rapid deployment
-
-

SEQ. #: 3410**Long Name:** VS-Aortic Valve Repair - Commissural Suture Annuloplasty**Short Name:** VSAVRComA**Definition:** Indicate whether the aortic valve repair procedure included a commissural annuloplasty.

Intent/Clarification: Sometimes referred to as “subcommissural annuloplasty”. Identifies repairs involving placement of pledgeted mattress sutures across the upper portion of the commissural post to improve leaflet coaptation. These annuloplasty sutures are contained with the inside of the aorta, in contrast to the sutures for commissural resuspension (seq# 3425).

SEQ. #: 3411

Long Name: VS-Aortic Valve Repair - External Suture Annuloplasty

Short Name: VSAVRExSutAn

Definition: Indicate whether the aortic valve repair procedure included an external suture annuloplasty.

Intent/Clarification: To identify placement of the annuloplasty suture outside the right/left commissure, passing the needle through the septal myocardium.

SEQ. #: 3415

Long Name: VS-Aortic Valve Repair - Leaflet Plication

Short Name: VSAVRLPlic

Definition: Indicate whether the aortic valve repair procedure included leaflet plication.

Intent/Clarification: To identify repair with central plication stitches, shortening the leaflet free-edge length for the correction of leaflet prolapse.

SEQ. #: 3416

Long Name: VS-Aortic Valve Repair - Nodular Release

Short Name: VSAVRNodRel

Definition: Indicate whether the aortic valve repair procedure included nodular release.

Intent/Clarification: -

SEQ. #: 3420

Long Name: VS-Aortic Valve Repair - Leaflet Free Edge Reinforcement (PTFE) Suture

Short Name: VSAVRPTFE

Definition: Indicate whether the aortic valve repair procedure included leaflet free edge reinforcement (PTFE) suture.

Intent/Clarification: The free edge reinforcement technique is performed by using suture passed in running fashion over and over along the entire length of the free margin.

SEQ. #: 3425

Long Name: VS-Aortic Valve Repair - Leaflet Commissural Resuspension Suture

Short Name: VSAVRComRS

Definition: Indicate whether the aortic valve repair procedure included leaflet commissural resuspension suture.

Intent/Clarification: A commissural resuspension suture is a pledgeted mattress suture placed at the top end of the commissural post. The stitch is placed transmurally, so that one pledget is on the inside of the aorta and the other pledget is on the outside of the aorta. This suture has the effect of compressing all aortic layers together and is often used in repair of aortic dissections.

SEQ. #: 3430

Long Name: VS-Aortic Valve Repair - Division of Fused Leaflet Raphe

Short Name: VSAVRRaphe

Definition: Indicate whether the aortic valve repair procedure included division of fused leaflet raphe.

Intent/Clarification: The division of the raphe (the two commissures or hinge points that are fused) in bicuspid valves.

SEQ. #: 3435

Long Name: VS-Aortic Valve Repair - Ring Annuloplasty

Short Name: VSAVRRingA

Definition: Indicate whether the aortic valve repair procedure included a ring annuloplasty.

Intent/Clarification: Describes a ring sewn around the base to the annulus to reshape it and provide support. Rings may be flexible or rigid.

SEQ. #: 3436

Long Name: VS-Aortic Valve Repair - Ring Annuloplasty - Type

Short Name: VSAVRRingATy

Definition: Indicate the type of ring annuloplasty that was used in this procedure.

Intent/Clarification:

- External ring
 - Internal ring
-
-

SEQ. #: 3440

Long Name: VS-Aortic Valve Repair - Leaflet Resection Suture

Short Name: VSAVRLResect

Definition: Indicate whether the aortic valve repair procedure included leaflet resection.

Intent/Clarification: Sutures places to mark the edges of the resection.

SEQ. #: 3441

Long Name: VS-Aortic Valve Repair - Leaflet Shaving

Short Name: VSAVRLeafShav

Definition: Indicate whether the aortic valve repair procedure included leaflet shaving.

Intent/Clarification: Removing the growth

SEQ. #: 3445

Long Name: VS-Aortic Valve Repair - Leaflet Pericardial Patch

Short Name: VSAVRLPPatch

Definition: Indicate whether the aortic valve repair procedure included leaflet pericardial patch.

Intent/Clarification: A pericardial patch can be used to repair larger perforations in the valve leaflets

SEQ. #: 3450

Long Name: VS-Aortic Valve Repair - Leaflet Debridement

Short Name: VSAVRDeb

Definition: Indicate whether the aortic valve repair procedure included leaflet debridement.

Intent/Clarification: A debridement technique can be used to remove small leaflet lesions such as LambI's excrescence, fibroelastomas and small calcific deposits. When tumors such as fibroelastoma or myxoma are removed, also code in seq # 4115.

SEQ. #: 3455

Long Name: VS-Aortic Valve Repair - Repair of Peri-prosthetic Leak

Short Name: VSAVRPeriLeak

Definition: Indicate whether the aortic valve repair procedure included repair of a Peri-prosthetic leak.

Intent/Clarification: Leak of a previously place valve prosthesis. A periprosthetic leak occurs because of gap between the valve sewing ring and the native annulus. Repair of such a leak may mandate removal of the entire valve and re-replacement. This is not the intent of this field. Rather, this field pertains to the repair of a peri-prosthetic leak with one or more repair sutures without needing to remove the existing prosthesis.

SEQ. #: 3460

Long Name: VS-Aortic Proc-Aortic Annular Enlargement

Short Name: AnlrEnl

Definition: Indicate whether an annular enlargement procedure was performed on the Aortic Valve. An aortic annular enlargement is defined as incision of the aortic annulus to enlarge the aortic orifice. Annular enlargement techniques include but are not limited to Manougian, Konno and Nicks.

Intent/Clarification:

Enlargement of the aortic annulus during aortic valve replacement permits insertion of a larger prosthetic valve or allows for optimal positioning. The enlarging procedure typically employs a patch of either pericardium or Dacron. In the classic Nick's or Manougian, the patch extends across the annulus (an aorto-annuloplasty). A patch that extends down to but not across the actual annulus (supra-annular aortoplasty) is considered a modification of the Nick's or Manougian and is coded as Nick's or Manougian as appropriate.

SEQ. #: 3461

Long Name: VS-Aortic Proc-Aortic Annular Enlargement With Patch - Technique

Short Name: AnlrEnlTech

Definition: Indicate the technique used for the aortic annular enlargement procedure.

Intent/Clarification:

Intended to capture whether a Nicks-Nunez, Manougian, Konno, Other or Unknown was performed utilizing patch material.

SEQ. #: 3462

Long Name: VS-Aortic Root Procedure

Short Name: VSAVRoot

Definition: Indicate whether an aortic root procedure was performed during this operation.

Intent/Clarification: For AV surgery involving the aortic root, also complete section M2.

SEQ. #: 3463

Long Name: VS-Aortic Root Procedure With Coronary Ostial Reimplantation (Bentall)

Short Name: VSAVRootOReimp

Definition: Indicate whether the root replacement procedure included coronary Ostial Reimplantation (Bentall).

Intent/Clarification: The coronary ostia are reimplanted following the replacement of the aortic root.

SEQ. #: 3464

Long Name: VS-Aortic Root Procedure With Coronary Ostial Reimplantation (Bentall) - Type

Short Name: VSAVRootORempTy

Definition: Indicate the type of device used for root replacement.

Intent/Clarification:

SEQ. #: 3465

Long Name: VS-Aortic Root Procedure With Coronary Ostial Reimplantation - Bioprosthetic Type

Short Name: VSAVRepBioTy

Definition: Indicate the type of bioprosthetic device used during the aortic root replacement with coronary Ostial Reimplantation

Intent/Clarification:

SEQ. #: 3466

Long Name: VS-Aortic Valve Sparing Root Operation Performed

Short Name: VSAVSparRt

Definition: Indicate whether a valve sparing root operation was performed.

Intent/Clarification:

SEQ. #: 3467

Long Name: VS-Aortic Valve Sparing Root Operation

Short Name: VSAVSparRtOp

Definition: Indicate the type of aortic valve sparing root operation that was performed.

Intent/Clarification:

SEQ. #: 3468

Long Name: VS-Aortic Valve Major Root Reconstruction

Short Name: VSAVRootRecon

Definition: Indicate whether the procedure included aortic valve major root reconstruction / debridement with or without pericardial patch.

Intent/Clarification:

SEQ. #: 3469

Long Name: VS-Aortic Valve Patch

Short Name: VSAVPat

Definition: Indicate whether a patch was used

Intent/Clarification:

SEQ. #: 3470

Long Name: VS-Aortic Valve Patch Type

Short Name: VSAVPatTy

Definition: Indicate the type of patch used

Intent/Clarification:

SEQ. #: 3472

Long Name: VS-Aortic Valve Implant

Short Name: AorticImplant

Definition: Indicate whether an aortic valve or valve repair device was implanted.

Intent/Clarification:

FAQ August 2017: When will the updated valve/VAD list be published?

Answer: The STS will not be publishing a Valve/VAD list. It is the responsibility of your vendor to maintain the lists in the drop-downs in your software.

SEQ. #: 3480

Long Name: VS-Aortic Proc-Implant Model Number

Short Name: VSAoIm

Definition: Indicate the name of the prosthesis implanted. The names provided include the manufacturer's model number with "xx" substituting for the device size.

Intent/Clarification:

Choose the device type from the device list.

SEQ. #: 3485

Long Name: VS-Aortic Proc-Imp-Size

Short Name: VSAoImSz

Definition: Indicate the Aortic implant size.

Intent/Clarification:

SEQ. #: 3490

Long Name: VS-Aortic Proc-Imp - Unique Device Identifier (UDI)

Short Name: VSAoImUDI

Definition: Indicate the device UDI if available, otherwise leave blank.

Intent/Clarification:

This is a unique identifier that will be on each patch. It may not be available immediately. If not available leave blank. Please refer to the FDA web site for detailed explanation of UDI.

https://www.fda.gov/MedicalDevices/DeviceRegulationandGuidance/UniqueDeviceIdentification/default.htm?utm_s%20our%20ce=Members%E2%80%90Only+Updates&utm_campaign=c7c1e8c870%E2%80%90%20Proposed_Rules_7_5_2012&utm_medium=email

SEQ. #: 3495

Long Name: VS-Mitral Valve

Short Name: VSMV

Definition: Indicate whether a mitral valve procedure was performed.

Intent/Clarification:

- Yes, planned
- Yes, unplanned due to surgical complication
- Yes, unplanned due to unsuspected disease or anatomy
- No

SEQ. #: 3500

Long Name: VS-Mitral Valve Procedure

Short Name: VSMVPr

Definition: Indicate the type of procedure that was performed on the mitral valve.

Intent/Clarification:

- Repair
- Replacement

FAQ September 2017: Surgeon performed the following mitral valve procedure “anterior mitral leaflet endarterectomy/decalcification” done in conjunction with an Aortic Valve Replacement. How is this documented under the options provided for MV repair?

Answer: No, anterior mitral leaflet endarterectomy/decalcification is considered part of the AVR and should not be coded as a mitral valve procedure.

SEQ. #: 3501

Long Name: VS-Mitral Valve - Repair Approach

Short Name: VSMVRepApp

Definition: Indicate the approach that was used to repair the Mitral Valve.

Intent/Clarification:

- Transcatheter
- Surgical

SEQ. #: 3505

Long Name: VS-Mitral Valve Repair - Annuloplasty

Short Name: VSMitRAnnulo

Definition: Indicate whether the mitral valve repair procedure included an annuloplasty.

Intent/Clarification:

SEQ. #: 3510

Long Name: VS-Mitral Valve Repair - Leaflet Resection

Short Name: VSMitRLeafRes

Definition: Indicate whether the mitral valve repair procedure included a leaflet resection.

Intent/Clarification:

SEQ. #: 3515

Long Name: VS-Mitral Leaflet Resection Type

Short Name: VSLeafResTyp

Definition: Indicate the type of leaflet resection.

Intent/Clarification:

- Triangular
- Quadrangular
- Other

SEQ. #: 3517

Long Name: VS-Mitral Repair Leaflet - Anterior Resection

Short Name: VSLeafAntRes

Definition: Indicate whether anterior MV leaflet resection was performed

Intent/Clarification:

- Yes
- No

SEQ. #: 3518

Long Name: VS-Mitral Repair Leaflet - Anterior Resection - Location Documented

Short Name: VSLeafAntResLocD

Definition: Indicate whether the location of the anterior resection was documented.

Intent/Clarification:

- Yes
- No

SEQ. #: 3519

Long Name: VS-Mitral Repair Leaflet - Anterior Resection - A1

Short Name: VSLeafAntResA1

Definition: Indicate whether the anterior leaflet resection included location A1

Intent/Clarification:

SEQ. #: 3520

Long Name: VS-Mitral Repair Leaflet - Anterior Resection - A2

Short Name: VSLeafAntResA2

Definition: Indicate whether the anterior leaflet resection included location A2

Intent/Clarification:

SEQ. #: 3521

Long Name: VS-Mitral Repair Leaflet - Anterior Resection - A3

Short Name: VSLeafAntResA3

Definition: Indicate whether the anterior leaflet resection included location A3

Intent/Clarification:

SEQ. #: 3522

Long Name: VS-Mitral Repair Leaflet - Posterior Resection

Short Name: VSLeafPostRes

Definition: Indicate whether posterior MV leaflet resection was performed

Intent/Clarification:

- Yes
 - No
- -----

SEQ. #: 3523

Long Name: VS-Mitral Repair Leaflet - Posterior Resection - Location Documented

Short Name: VSLeafPostResLocD

Definition: Indicate whether posterior MV leaflet resection location was documented

Intent/Clarification:

SEQ. #: 3524

Long Name: VS-Mitral Repair Leaflet - Posterior Resection - P1

Short Name: VSLeafPostResP1

Definition: Indicate whether the posterior leaflet resection included location P1

Intent/Clarification:

SEQ. #: 3525

Long Name: VS-Mitral Repair Leaflet - Posterior Resection - P2

Short Name: VSLeafPostResP2

Definition: Indicate whether the posterior leaflet resection included location P2

Intent/Clarification:

SEQ. #: 3526

Long Name: VS-Mitral Repair Leaflet - Posterior Resection - P3

Short Name: VSLeafPostResP3

Definition: Indicate whether the posterior leaflet resection included location P3

Intent/Clarification:

SEQ. #: 3527

Long Name: VS-Mitral Repair Leaflet - Commissure Resection

Short Name: VSLeafComRes

Definition: Indicate whether resection of the mitral commissure was performed

Intent/Clarification:

- Yes
 - No
- -----

SEQ. #: 3528

Long Name: VS-Mitral Repair Leaflet - Commissure Resection - Location

Short Name: VSLeafComResLoc

Definition: Indicate the location of the mitral commissure resection

Intent/Clarification:

- Medial
 - Lateral
 - Both
 - Not Documented
- -----

SEQ. #: 3532

Long Name: VS-Mitral Valve Repair - Neochords (PTFE)

Short Name: VSMitRPTFE

Definition: Indicate whether the mitral valve repair procedure included neochords (PTFE).

Intent/Clarification: Intended to replace damaged chordae by delivering artificial chordae tendineae.

SEQ. #: 3534

Long Name: VS-Mitral Valve Repair - Anterior Neochords

Short Name: VSNeoAnt

Definition: Indicate whether anterior neochords were placed

Intent/Clarification:

- Yes
- No

SEQ. #: 3535

Long Name: VS-Mitral Valve Repair - Anterior Neochords - Location Documented

Short Name: VSNeoAntLocD

Definition: Indicate whether location of anterior neochord placement was documented

Intent/Clarification:

- Yes
- No

SEQ. #: 3536

Long Name: VS-Mitral Valve Repair - Anterior Neochords - A1

Short Name: VSNeoAntA1

Definition: Indicate whether neochord location included location A1

Intent/Clarification:

SEQ. #: 3537

Long Name: VS-Mitral Valve Repair - Anterior Neochords - A2

Short Name: VSNeoAntA2

Definition: Indicate whether neochord location included location A2

Intent/Clarification:

SEQ. #: 3538

Long Name: VS-Mitral Valve Repair - Anterior Neochords - A3

Short Name: VSNeoAntA3

Definition: Indicate whether neochord location included location A3

Intent/Clarification:

SEQ. #: 3539

Long Name: VS-Mitral Valve Repair - Posterior Neochords

Short Name: VSNeoPost

Definition: Indicate whether posterior neochords were placed

Intent/Clarification:

- Yes
 - No
-
-

SEQ. #: 3540

Long Name: VS-Mitral Valve Repair - Posterior Neochords - Location Documented

Short Name: VSNeoPostLocD

Definition: Indicate whether location of posterior neochord placement was documented

Intent/Clarification:

- Yes
 - No
-
-

SEQ. #: 3541

Long Name: VS-Mitral Valve Repair - Posterior Neochords - P1

Short Name: VSNeoPostP1

Definition: Indicate whether posterior neochord location included location P1

Intent/Clarification:

SEQ. #: 3542

Long Name: VS-Mitral Valve Repair - Posterior Neochords - P2

Short Name: VSNeoPostP2

Definition: Indicate whether posterior neochord location included location P2

Intent/Clarification:

SEQ. #: 3543

Long Name: VS-Mitral Valve Repair - Posterior Neochords - P3

Short Name: VSNeoPostP3

Definition: Indicate whether posterior neochord location included location P3

Intent/Clarification:

SEQ. #: 3544

Long Name: VS-Mitral Valve Repair - Commissure Neochords

Short Name: VSNeoCom

Definition: Indicate whether commissural neochords were placed

Intent/Clarification:

- Yes
 - No
-
-

SEQ. #: 3545

Long Name: VS-Mitral Valve Repair - Commissure Neochords - Location

Short Name: VSNeoComLoc

Definition: Indicate location of commissural neochord placement

Intent/Clarification:

- Medial
 - Lateral
 - Both
 - Not Documented
-
-

SEQ. #: 3550

Long Name: VS-Mitral Valve Repair - Chordal / Leaflet Transfer

Short Name: VSMitRChord

Definition: Indicate whether the mitral valve repair procedure included a chordal / leaflet transfer.

Intent/Clarification:

- Yes
 - No
-
-

SEQ. #: 3551

Long Name: VS-Mitral Valve Repair - Chordal Leaflet Transfer - Anterior

Short Name: VSChorLfAnt

Definition: Indicate whether chordal leaflet transfer was anterior

Intent/Clarification:

- Yes
- No

SEQ. #: 3552

Long Name: VS-Mitral Valve Repair - Chordal Leaflet Transfer - Anterior Location Documented

Short Name: VSChorLfAntLocD

Definition: Indicate whether location of anterior chordal leaflet transfer was documented

Intent/Clarification:

- Yes
 - No
-
-

SEQ. #: 3553

Long Name: VS-Mitral Valve Repair - Chordal Leaflet Transfer - Anterior - A1

Short Name: VSChorLfAntA1

Definition: Indicate whether anterior chordal leaflet transfer location was A1

Intent/Clarification:

SEQ. #: 3554

Long Name: VS-Mitral Valve Repair - Chordal Leaflet Transfer - Anterior - A2

Short Name: VSChorLfAntA2

Definition: Indicate whether anterior chordal leaflet transfer location was A2

Intent/Clarification:

SEQ. #: 3555

Long Name: VS-Mitral Valve Repair - Chordal Leaflet Transfer - Anterior - A3

Short Name: VSChorLfAntA3

Definition: Indicate whether anterior chordal leaflet transfer location was A3

Intent/Clarification:

SEQ. #: 3556

Long Name: VS-Mitral Valve Repair - Chordal Leaflet Transfer - Posterior

Short Name: VSChorLfPost

Definition: Indicate whether chordal leaflet transfer was posterior

Intent/Clarification:

- Yes
- No

SEQ. #: 3557

Long Name: VS-Mitral Valve Repair - Chordal Leaflet Transfer - Posterior Location Documented

Short Name: VSChorLfPostLocD

Definition: Indicate whether location of posterior chordal leaflet transfer was documented

Intent/Clarification:

- Yes
- No

SEQ. #: 3558

Long Name: VS-Mitral Valve Repair - Chordal Leaflet Transfer - Posterior - P1

Short Name: VSChorLfPostP1

Definition: Indicate whether posterior chordal leaflet transfer location was P1

Intent/Clarification:

SEQ. #: 3559

Long Name: VS-Mitral Valve Repair - Chordal Leaflet Transfer - Posterior - P2

Short Name: VSChorLfPostP2

Definition: Indicate whether posterior chordal leaflet transfer location was P2

Intent/Clarification:

SEQ. #: 3560

Long Name: VS-Mitral Valve Repair - Chordal Leaflet Transfer - Posterior - P3

Short Name: VSChorLfPostP3

Definition: Indicate whether posterior chordal leaflet transfer location was P3

Intent/Clarification:

SEQ. #: 3561

Long Name: VS-Mitral Valve Repair - Chordal Leaflet Transfer - Commissure

Short Name: VSChorLfCom

Definition: Indicate whether chordal leaflet transfer was commissural

Intent/Clarification:

- Yes
 - No
-
-

SEQ. #: 3562

Long Name: VS-Mitral Valve Repair - Chordal Leaflet Transfer - Commissure Location

Short Name: VSChorLfComLoc

Definition: Indicate location of commissural leaflet transfer

Intent/Clarification:

- Medial
 - Lateral
 - Both
 - Not Documented
-
-

SEQ. #: 3565

Long Name: VS-Mitral Valve Repair - Folding Plasty

Short Name: VSMitRFold

Definition: Indicate whether the mitral valve repair procedure included folding plasty.

Intent/Clarification:

SEQ. #: 3566

Long Name: VS-Mitral Valve Repair - Sliding Plasty

Short Name: VSMitRSlidP

Definition: Indicate whether the mitral valve repair procedure included a sliding plasty.

Intent/Clarification:

SEQ. #: 3567

Long Name: VS-Mitral Valve Repair - Annular Decalcification / Debridement

Short Name: VSMitRADecalc

Definition: Indicate whether the mitral valve repair procedure included an annular decalcification / debridement.

Intent/Clarification:

SEQ. #: 3568

Long Name: VS-Mitral Valve Repair - Leaflet Extension / Replacement / Patch

Short Name: VSMitRLeafERP

Definition: Indicate whether the mitral valve repair procedure included a leaflet extension / replacement / patch.

Intent/Clarification:

SEQ. #: 3569

Long Name: VS-Mitral Valve Repair - Leaflet Extension / Replacement / Patch - Location

Short Name: VSMitRLeafERPLoc

Definition: Indicate the location of the mitral leaflet extension/replacement patch

Intent/Clarification:

SEQ. #: 3570

Long Name: VS-Mitral Valve Repair - Edge To Edge Repair

Short Name: VSMitREdge

Definition: Indicate whether the mitral valve repair procedure included an edge to edge repair.

Intent/Clarification:

Edge-to-edge repair is a surgical approximation of the mitral valve leaflets, sometimes called the Alfieri procedure or Bow Tie procedure.

SEQ. #: 3580

Long Name: VS-Mitral Valve Repair - Mitral Commissurotomy

Short Name: VSMitRMitComm

Definition: Indicate whether the mitral valve repair procedure included a mitral commissurotomy.

Intent/Clarification: Disruption of the components of a commissure fused as a result of valvular disease.

SEQ. #: 3585

Long Name: VS-Mitral Valve Repair - Mitral Commissuroplasty

Short Name: VSMitRMitCplasty

Definition: Indicate whether the mitral valve repair procedure included a mitral commissuroplasty.

Intent/Clarification:

SEQ. #: 3590

Long Name: VS-Mitral Valve Repair - Mitral Cleft Repair (Scallop Closure)

Short Name: VSMitRMitCleft

Definition: Indicate whether the mitral valve repair procedure included a mitral cleft repair.

Intent/Clarification:

SEQ. #: 3591

Long Name: VS-Mitral Valve Repair - Paraprosthesis Leak Repair

Short Name: VSMitParaprosLeak

Definition: Indicate whether there was repair of a mitral paraprosthesis leak

Intent/Clarification: Leak of a previously placed valve prosthesis.

SEQ. #: 3600

Long Name: VS-Mitral Repair Attempted

Short Name: MitralIntent

Definition: Indicate whether a Mitral Valve Repair was attempted prior to the Mitral Valve Replacement.

Intent/Clarification:

- Yes
 - No
- -----

SEQ. #: 3605

Long Name: VS-Mitral Chordal Preservation

Short Name: VSChorPres

Definition: Indicate whether native chords were preserved.

Intent/Clarification:

- Anterior
- Posterior

- Both
- None

SEQ. #: 3610

Long Name: VS-Mitral Transcatheter Valve Replacement

Short Name: VSTCVMit

Definition: Indicate whether the mitral valve replacement was done using a transcatheter valve device.

Intent/Clarification:

SEQ. #: 3615

Long Name: VS-Mitral Implant

Short Name: MitralImplant

Definition: Indicate whether a mitral valve or valve device was implanted.

Intent/Clarification:

- Yes
- No

FAQ August 2017: When will the updated valve/VAD list be published?

Answer: The STS will not be publishing a Valve/VAD list. It is the responsibility of your vendor to maintain the lists in the drop-downs in your software.

SEQ. #: 3620

Long Name: VS-Mitral Implant - Type

Short Name: MitralImplantTy

Definition: Indicate the type of mitral valve or valve device implanted.

Intent/Clarification:

- Mechanical Valve
 - Bioprosthetic valve
 - Annuloplasty device
 - Mitral Leaflet clip
 - Transcatheter device
 - Surgically implanted transcatheter device
 - Other
-
-

SEQ. #: 3625

Long Name: VS-Mitral Proc-Implant Model Number

Short Name: VSMilm

Definition: Indicate the model number of the device implanted. The names provided include the manufacturer's model number with "xx" substituting for the device size.

Intent/Clarification:

Choose the device type from the device list.

SEQ. #: 3630

Long Name: VS-Mitral Proc-Imp-Size

Short Name: VSMilmSz

Definition: Indicate the Mitral implant size.

Intent/Clarification:

SEQ. #: 3635

Long Name: VS-Mitral Proc-Imp-Unique Device Identifier (UDI)

Short Name: VSMilmUDI

Definition: Indicate the device UDI if available, otherwise leave blank.

Intent/Clarification:

This is a unique identifier that will be on each valve. It may not be available immediately. If not available leave blank. Please refer to the FDA web site for detailed explanation of UDI.

https://www.fda.gov/MedicalDevices/DeviceRegulationandGuidance/UniqueDeviceIdentification/default.htm?utm_source=Members%E2%80%90Only+Updates&utm_campaign=c7c1e8c870%E2%80%90%20Proposed_Rules_7_5_2012&utm_medium=email

SEQ. #: 3640

Long Name: VS-Tricuspid Valve

Short Name: VSTV

Definition: Indicate whether a tricuspid valve procedure was performed.

Intent/Clarification:

- Yes, planned
- Yes, unplanned due to surgical complication
- Yes, unplanned due to unsuspected disease or anatomy
- No

SEQ. #: 3646

Long Name: VS-Tricuspid Repair

Short Name: VSTrRepair

Definition: Indicate whether tricuspid repair was performed

Intent/Clarification:

- Yes
 - No
-
-

SEQ. #: 3647

Long Name: VS-Tricuspid Repair - Annuloplasty

Short Name: VSTrRepAnnulo

Definition: Indicate whether the tricuspid repair included an annuloplasty

Intent/Clarification:

- Yes
 - No
-
-

SEQ. #: 3648

Long Name: VS-Tricuspid Repair - Annuloplasty Type

Short Name: OpTricusAnTy

Definition: Indicate type of annuloplasty procedure.

Intent/Clarification:

- Pericardium
 - Suture
 - Prosthetic ring
 - Prosthetic band
 - Other
-
-

SEQ. #: 3649

Long Name: VS-Tricuspid Repair - Leaflet Resection

Short Name: VSTrLeafRes

Definition: Indicate whether the tricuspid repair included leaflet resection

Intent/Clarification:

- Yes
- No

SEQ. #: 3650

Long Name: VS-Tricuspid Replacement

Short Name: VSTrReplace

Definition: Indicate whether tricuspid replacement was performed

Intent/Clarification:

- Yes
- No

FAQ August 2017: When will the updated valve/VAD list be published?

Answer: The STS will not be publishing a Valve/VAD list. It is the responsibility of your vendor to maintain the lists in the drop-downs in your software.

SEQ. #: 3652

Long Name: VS-Tricuspid Transcatheter Valve Replacement

Short Name: VSTCVTri

Definition: Indicate whether the tricuspid valve replacement was done using a transcatheter valve device.

Intent/Clarification:

- Yes
 - No
-
-

SEQ. #: 3653

Long Name: VS-Tricuspid Valvectomy **Short Name:** VSTrValvec

Definition: Indicate whether tricuspid valvectomy was performed

Intent/Clarification:

Intended to capture procedures where the tricuspid valve is removed.

SEQ. #: 3660

Long Name: VS-Tricuspid Implant **Short Name:** TricuspidImplant

Definition: Indicate whether a tricuspid valve or device was implanted.

Intent/Clarification:

- Yes
 - No
-
-

SEQ. #: 3665

Long Name: VS-Tricuspid Implant - Type

Short Name: TricusImplantTy

Definition: Indicate the type of tricuspid valve or valve device implanted.

Intent/Clarification:

- Mechanical valve
- Annuloplasty device
- Bioprosthetic valve
- Transcatheter device
- Homograft
- Other

SEQ. #: 3670

Long Name: VS-Tricuspid Proc-Implant Model Number

Short Name: VSTrlm

Definition: Indicate the model number of the prosthesis implanted. The names provided include the manufacturer's model number with "xx" substituting for the device size.

Intent/Clarification:

Choose the device type from the device list.

SEQ. #: 3675

Long Name: VS-Tricuspid Proc-Imp-Size

Short Name: VSTrlmSz

Definition: Indicate the Tricuspid implant size.

Intent/Clarification:

SEQ. #: 3680

Long Name: VS-Tricuspid Proc-Imp-Unique Device Identifier (UDI)

Short Name: VSTrlmUDI

Definition: Indicate the device UDI if available, otherwise leave blank.

Intent/Clarification:

This is a unique identifier that will be on each valve. It may not be available immediately. If not available leave blank. Please refer to the FDA web site for detailed explanation of UDI.

https://www.fda.gov/MedicalDevices/DeviceRegulationandGuidance/UniqueDeviceIdentification/default.htm?utm_source=our%20ce=Members%E2%80%90Only+Updates&utm

SEQ. #: 3685

Long Name: VS-Pulmonic Valve

Short Name: VSPV

Definition: Indicate whether a pulmonic valve procedure was performed.

Intent/Clarification:

- Yes, planned
 - Yes, unplanned due to surgical complication
 - Yes, unplanned due to unsuspected disease or anatomy
 - No
-
-

SEQ. #: 3690

Long Name: VS-Pulmonic Proc-Procedure

Short Name: OpPulm

Definition: Indicate the type of procedure that was performed on the pulmonic valve.

Intent/Clarification:

- Replacement
 - Reconstruction
 - Valvectomy
-
-

SEQ. #: 3695

Long Name: VS-Pulmonic Transcatheter Valve Replacement

Short Name: VSTCVPu

Definition: Indicate whether the pulmonic valve replacement was done using a transcatheter valve device.

Intent/Clarification:

- Yes
 - No
-
-

SEQ. #: 3700

Long Name: VS-Pulmonic Implant

Short Name: PulmonicImplant

Definition: Indicate whether a pulmonic valve or device was implanted.

Intent/Clarification:

- Yes
- No

FAQ August 2017: When will the updated valve/VAD list be published?

Answer: The STS will not be publishing a Valve/VAD list. It is the responsibility of your vendor to maintain the lists in the drop-downs in your software.

SEQ. #: 3701

Long Name: VS-Pulmonic - Type Of Implant

Short Name: VSPuTypImp

Definition: Indicate the type of pulmonic implant

Intent/Clarification:

- Surgeon Fashioned
 - Commercially Supplied
-
-

SEQ. #: 3702

Long Name: VS-Pulmonic - Surgeon Fashioned Implant Material

Short Name: VSPulmpMat

Definition: Indicate the material used to fashion the pulmonic implant

Intent/Clarification: Unlike conventional valve replacement, measured and crafted to meet specific dimensions of the annulus.

- PTFE (Gore-Tex)
 - Pericardium
 - Other
-
-

SEQ. #: 3705

Long Name: VS-Pulmonic Implant - Type **Short Name:** PulmonicImplantTy

Definition: Indicate the type of pulmonic valve or valve device implanted.

Intent/Clarification:

- Mechanical valve
 - Annuloplasty device
 - Bioprosthetic valve
 - Transcatheter device
 - Homograft
 - Other
-
-

SEQ. #: 3710

Long Name: VS-Pulmonic Proc-Implant Model Number

Short Name: VSPulm

Definition: Indicate the model number of the prosthesis implanted. The names provided include the manufacturer's model number with "xx" substituting for the device size.

Intent/Clarification:

Choose the device type from the device list.

SEQ. #: 3715

Long Name: VS-Pulmonic Proc-Imp-Size

Short Name: VSPulmSz

Definition: Indicate the Pulmonic implant size.

Intent/Clarification:

SEQ. #: 3720

Long Name: VS-Pulmonic Proc-Imp-Unique Device Identifier **Short Name:**

VSPulmUDI

Definition: Indicate the device UDI if available, otherwise leave blank.

Intent/Clarification:

This is a unique identifier that will be on each valve. It may not be available immediately. If not available leave blank. Please refer to the FDA web site for detailed explanation of UDI.

https://www.fda.gov/MedicalDevices/DeviceRegulationandGuidance/UniqueDeviceIdentification/default.htm?utm_source=Members%E2%80%90Only+Updates&utm_campaign=c7c1e8c870%E2%80%90%20Proposed_Rules_7_5_2012&utm_medium=email

Mechanical Cardiac Assist Devices

SEQ. #: 3725

Long Name: IABP

Short Name: IABP

Definition: Indicate whether the patient was placed on an Intra-Aortic Balloon Pump (IABP).

Intent/Clarification:

IABP is a device inserted into the descending thoracic aorta distal to the left subclavian and proximal to the renal arteries used to increase coronary blood flow and decrease work of the left ventricle. Balloon catheter inflates and deflates rapidly in conjunction with cardiac cycle. Inflation of the balloon partially obstructs the aorta, diverting more blood into coronary arteries. Deflation of the balloon just prior to systole, allows blood to be more easily ejected by the left ventricle. This applies to IABP devices in at the time of surgery, not previously placed and removed devices.

- Yes
- No

SEQ. #: 3730

Long Name: IABP-When Inserted

Short Name: IABPWhen

Definition: Indicate when the IABP was inserted.

Intent/Clarification:

Identify when the IABP was inserted as it relates to the cardiac operation.

- **Preop** refers to the IABP placement in the Cath lab or in the ICU prior to patient entering the operating room.
- **Intraop** refers to insertion of the IABP during the cardiac operation (after the patient has entered the operating room and before the patient leaves the operating room).
- **Postop** refers to insertion of the IABP after the patient has left the operating room.

SEQ. #: 3735

Long Name: IABP-Indication

Short Name: IABPInd

Definition: Indicate the primary reason for inserting the IABP.

Intent/Clarification:

The reason for inserting an IABP as it relates to the cardiac operation. Choose one of the following:

- Hemodynamic instability (hypotension/shock)
- Procedural support
- Unstable angina
- Cardiopulmonary Bypass (CPB) weaning failure
- Prophylactic
- Other

SEQ. #: 3745

Long Name: Catheter Based Assist Device Used

Short Name: CathBasAssist

Definition: Indicate whether the patient was placed on a catheter based assist device (e.g., Impella).

Intent/Clarification:

Catheter based assist devices offer short term minimally invasive circulatory support. Catheter Based Assist Devices are only captured in this section and are not included in section L.2 Ventricular Assist Devices. Examples include Impella, Tandem Heart. Do not capture devices inserted and removed prior to the operation.

SEQ. #: 3755

Long Name: Catheter Based Assist Type

Short Name: CathBasAssistTy

Definition: Indicate the type of catheter based assist device.

Intent/Clarification:

- RV (Right Ventricular)
 - LV (Left Ventricular)
 - BiVAD (Biventricular)
-
-

SEQ. #: 3760

Long Name: Catheter Based Assist Device When Inserted

Short Name: CathBasAssistWhen

Definition: Indicate when the catheter based assist device was inserted.

Intent/Clarification:

Identify when the assist device was inserted as it relates to the cardiac operation.

- **Preop** refers to the assist device placement in the Cath lab or in the ICU prior to patient entering the operating room.
- **Intraop** refers to insertion of the assist device during the cardiac operation (after the patient has entered the operating room and before the patient leaves the operating room).
- **Postop** refers to insertion of the assist device after the patient has left the operating room.
- **Non-operative** refers to patients who have a catheter based assist initiated by a CT surgeon but are not having a CT surgery procedure. These may be for victims of near drowning, influenza, amniotic fluid embolus. Stand-alone

procedures are not mandatory to collect, however if your surgeon(s) elects to track these, use this harvest code.

SEQ. #: 3765

Long Name: Catheter Based Assist Device Indication

Short Name: CathBasAssistInd

Definition: Indicate the primary reason for inserting the device.

Intent/Clarification:

The goal is to identify the reason the device was inserted.

- Hemodynamic Instability
 - Cardiopulmonary Bypass (CPB) weaning failure
 - PCI Failure
 - Procedural support
 - Other
-
-

SEQ. #: 3775

Long Name: Extracorporeal Membrane Oxygenation

Short Name: ECMO

Definition: Indicate whether the patient was placed on ECMO.

Intent/Clarification:

ECMO, which stands for Extracorporeal Membrane Oxygenation, functions as a replacement for a critically ill patient's heart and lungs. It is used to support a patient who is awaiting surgery, or to give vital organs time to recover from heart surgery or disease. It can also be used to rewarm victims of hypothermia or drowning.

ECMO initiation may be done in the OR or at the bedside in the ICU.

- Venous-Venous
 - Venous-Arterial
 - Venous-venous converted to Venous-arterial
 - No (ECMO not initiated)
-
-

SEQ. #: 3780

Long Name: ECMO When Initiated

Short Name: ECMOWhen

Definition: Indicate when patient was placed on ECMO.

Intent/Clarification:

- **Preop** refers to placement in the Cath lab or in the ICU prior to patient entering the operating room.

- **Intraop** refers to insertion during the cardiac operation.
 - **Postop** refers to insertion after the patient has left the operating room.
 - **Non-Operative** refers to patients who have ECMO initiated by a CT surgeon but are not having a CT surgery procedure. Stand-alone procedures are not mandatory to collect, however if your surgeon(s) elects to track these, use this harvest code.
-
-

SEQ. #: 3785

Long Name: ECMO Indication

Short Name: ECMOInd

Definition: Indicate clinical indication for placing patient on ECMO.

Intent/Clarification:

The intent is to capture the indication for ECMO

- Cardiac Failure
 - Respiratory Failure
 - Hypothermia (**Intended to capture rewarming for hypothermic/drowning victims**)
 - Rescue/salvage (**Intended to capture cath lab or other disasters requiring support**)
 - Other
-
-

Ventricular Assist Devices

SEQ. #: 3790

Long Name: VAD-Patient Admitted With VAD

Short Name: PrevVAD

Definition: Indicate if at the time of this procedure, the patient has a VAD in place that was inserted during a previous admission or from an outside hospital.

Intent/Clarification:

SEQ. #: 3795

Long Name: Previous VAD Facility

Short Name: PrevVADF

Definition: Indicate if the previously implanted assist device was implanted at another facility.

Intent/Clarification:

SEQ. #: 3800

Long Name: Previous VAD Insertion Date

Short Name: PrevVADD

Definition: Indicate insertion date of previous VAD.

Intent/Clarification:

Required date format: mm/dd/yyyy.

SEQ. #: 3805

Long Name: Previous VAD Indication

Short Name: PrevVADIn

Definition: Specify indication for VAD insertion.

Intent/Clarification:

- **Bridge to Transplantation:** Includes those patients who are supported with a VAD until a heart transplant is possible.
 - **Bridge to Recovery:** Includes those patients who are expected to have ventricular recovery. (i.e. Myocarditis patients, viral cardiomyopathies, AMI w/ revascularization, and post-transplant reperfusion injury).
 - **Destination:** Includes those patients where a heart transplant is not an option. The VAD is placed for permanent life sustaining support.
 - **Post Cardiectomy Ventricular Failure:** Includes those postcardiotomy patients who receive a VAD because of failure to separate from the heart-lung machine. Postcardiotomy refers to those patients with the inability to wean from cardiopulmonary bypass secondary to left, right, or biventricular failure.
 - **Device Malfunction:** Includes those patients who are currently VAD supported and are experiencing device failure.
 - **End of (device) Life:** Mechanical device pump has reached functional life expectancy and requires replacement.
 - **Salvage:** Moribund patients unresponsive to medical interventions.
-
-

SEQ. #: 3810

Long Name: Previous VAD Type

Short Name: PrevVADTy

Definition: Indicate type of VAD previously inserted.

Intent/Clarification:

- Right VAD (RVAD) - Right Ventricular Assist Device
 - Left VAD (LVAD) - Left Ventricular Assist Device
 - Biventricular VAD (BiVAD) - Biventricular Assist Device
 - Total Artificial Heart (TAH)
- -----

SEQ. #: 3815

Long Name: Previous VAD Device Model Number

Short Name: PrevVADDevice

Definition: Indicate Previous VAD device.

Intent/Clarification:

Choose the device type from the device list.

SEQ. #: 3820

Long Name: Previous VAD Unique Device Identifier (UDI)

Short Name: PrevVADUDI

Definition: Indicate the device UDI if available, otherwise leave blank.

Intent/Clarification:

This is a unique identifier that will be on each device. It may not be available immediately. If not available leave blank. Please refer to the FDA web site for detailed explanation of UDI.

https://www.fda.gov/MedicalDevices/DeviceRegulationandGuidance/UniqueDeviceIdentification/default.htm?utm_source=Members%E2%80%90Only+Updates&utm_campaign=c7c1e8c870%E2%80%90%20Proposed_Rules_7_5_2012&utm_medium=email

SEQ. #: 3825

Long Name: Previous VAD Explanted During This Admission **Short Name:**

PrevVADExp

Definition: Indicate whether the previously inserted VAD was explanted during this hospitalization.

Intent/Clarification:

This is a unique identifier that will be on each VAD. It may not be available immediately. If not available leave blank.

https://www.fda.gov/MedicalDevices/DeviceRegulationandGuidance/UniqueDeviceIdentification/default.htm?utm_source=Members%E2%80%90Only+Updates&utm

SEQ. #: 3830

Long Name: Previous VAD Explanted During This Admission - Reason

Short Name: PrevVADExpRsn

Definition: Indicate the primary reason the VAD was explanted.

Intent/Clarification:

- Yes, not during this procedure
 - Yes, during this procedure
 - No
-
-

SEQ. #: 3835

Long Name: Previous VAD Explanted During This Admission - Date

Short Name: PrevVADExpDt

Definition: Indicate date of explant.

Intent/Clarification:

Choose the device type from the device list.

SEQ. #: 3840

Long Name: Ventricular Assist Device Implanted During This Hospitalization

Short Name: VADImp

Definition: Indicate whether a VAD was inserted during this hospitalization.

Intent/Clarification:

SEQ. #: 3845

Long Name: VAD-Implant Timing

Short Name: VADImpTmg

Definition: Indicate timing of VAD insertion.

Intent/Clarification:

Indicate the timing of insertion:

- Pre-operative (during the same hospitalization but not the same OR trip as the cardiovascular surgical procedure.
- Stand-alone VAD procedure-this was the only procedure performed.
- In conjunction with the cardiovascular surgical procedure (same trip to the OR)-

planned. In conjunction with a CV surgical procedure and planned before surgery, consent in the chart.

- In conjunction with the cardiovascular surgical procedure (same trip to the OR)-unplanned. In conjunction with a CV surgical procedure and not planned before surgery.
- Post-operative (after the surgical procedure during reoperation)

SEQ. #: 3850

Long Name: VAD-Indication for this VAD

Short Name: VADInd

Definition: Indicate the reason for implanting a Ventricular Assist Device (VAD) during this hospitalization.

Intent/Clarification:

- **Bridge to Transplantation:** Includes those patients who are supported with a VAD until a heart transplant is possible.
- **Bridge to Recovery:** Includes those patients who are expected to have ventricular recovery. (i.e. Myocarditis patients, viral cardiomyopathies, AMI w/revascularization, post-transplant reperfusion injury).
- **Destination:** Includes those patients where a heart transplant is not an option. The VAD is placed for permanent life sustaining support.
- **Post Cardiectomy Ventricular Failure:** Includes those postcardiotomy patients who receive a VAD because of failure to separate from the heart-lung machine. Postcardiotomy refers to those patients with the inability to wean from cardiopulmonary bypass secondary to left, right, or biventricular failure.
- **Device Malfunction:** Includes those patients who are currently VAD supported and are experiencing device failure.
- **End of (device) Life:** Mechanical device pump has reached functional life expectancy and requires replacement.
- **Salvage:** Moribund patients unresponsive to medical interventions.

SEQ. #: 3855

Long Name: VAD-Implant Type

Short Name: VImpTy

Definition: Indicate the first type of VAD implanted during this hospitalization.

Intent/Clarification:

- Right VAD (RVAD) - Right Ventricular Assist Device
 - Left VAD (LVAD) - Left Ventricular Assist Device
 - Biventricular VAD (BiVAD) - Biventricular Assist Device
 - Total Artificial Heart (TAH)
-

SEQ. #: 3860

Long Name: VAD-Device

Short Name: VProdTy

Definition: Indicate the VAD brand name implanted. Implant defined as physical placement of the VAD.

Intent/Clarification:

Choose the device type from the device list.

SEQ. #: 3865

Long Name: VAD-Implant Date

Short Name: VImpDt

Definition: Indicate the date the VAD was implanted.

Intent/Clarification:

Required date format: mm/dd/yyyy

SEQ. #: 3870

Long Name: VAD-Implant Unique Device Identifier (UDI)

Short Name: VImpUDI

Definition: Indicate the device UDI if available, otherwise leave blank.

Intent/Clarification:

This is a unique identifier that will be on each device. It may not be available immediately. If not available leave blank. Please refer to the FDA web site for detailed explanation of UDI.

https://www.fda.gov/MedicalDevices/DeviceRegulationandGuidance/UniqueDeviceIdentification/default.htm?utm_source=Members%E2%80%90Only+Updates&utm_campaign=c7c1e8c870%E2%80%90Proposed_Rules_7_5_2012&utm_medium=email

SEQ. #: 3875

Long Name: VAD-Explant

Short Name: VExp

Definition: Indicate if the VAD was explanted. Explant is defined as physical removal of the VAD.

Intent/Clarification:

- Yes, not during this procedure
- Yes, during this procedure
- No

SEQ. #: 3880

Long Name: VAD-Explant Reason

Short Name: VExpRsn

Definition: Indicate the reason the VAD was explanted.

Intent/Clarification:

- **Cardiac Transplant** -VAD was explanted for cardiac transplant.
- **Recovery** -VAD was removed after cardiac recovery.
- **Device Transfer** -VAD was explanted in order to implant another assist device.
- **Device-Related Infection** - An infection within the pump pocket, driveline, VAD endocarditis, or other infection requiring explantation of the VAD. The body of the VAD has an active infection requiring removal to eliminate the infection. "Device-related infections" are defined as positive culture in the presence of leukocytosis, and/or fever requiring medical or surgical intervention.
- **Device Malfunction** -The VAD pump itself is not functioning properly causing hemodynamic compromise, and/or requiring immediate intervention or VAD replacement.
- **End of (device) Life** -Mechanical device pump has reached functional life expectancy and requires replacement.

Note: Code "No" if the patient expires with the VAD in place; the VAD was not explanted.

SEQ. #: 3885

Long Name: VAD-Explant Date

Short Name: VExpDt

Definition: Indicate the date the VAD was explanted.

Intent/Clarification:

Required date format: mm/dd/yyyy

SEQ. #: 3895

Long Name: VAD-Implant #2

Short Name: VImp2

Definition: Indicate whether a second ventricular assist device was implanted.

Intent/Clarification:

SEQ. #: 3900

Long Name: VAD-Implant Timing #2

Short Name: VADImpTmg2

Definition: Indicate timing of VAD #2 insertion.

Intent/Clarification:

- Pre-operative (during the same hospitalization but not the same OR trip as the cardiovascular surgical procedure.
 - Stand-alone VAD procedure-this was the only procedure performed.
 - In conjunction with the cardiovascular surgical procedure (same trip to the OR)-planned. In conjunction with a CV surgical procedure and planned before surgery, consent in the chart.
 - In conjunction with the cardiovascular surgical procedure (same trip to the OR)-unplanned. In conjunction with a CV surgical procedure and not planned before surgery.
 - Post-operative (after the surgical procedure during reoperation).
-
-

SEQ. #: 3905

Long Name: VAD-Indication for this VAD #2

Short Name: VADInd2

Definition: Indicate the reason for implanting a Ventricular Assist Device (VAD) #2 during this hospitalization.

Intent/Clarification:

- **Bridge to Transplantation:** Includes those patients who are supported with a VAD until a heart transplant is possible.
 - **Bridge to Recovery:** Includes those patients who are expected to have ventricular recovery. (i.e. Myocarditis patients, viral cardiomyopathies, AMI w/revascularization, post-transplant reperfusion injury).
 - **Destination:** Includes those patients where a heart transplant is not an option. The VAD is placed for permanent life sustaining support.
 - **Post Cardiectomy Ventricular Failure:** Includes those postcardiectomy patients who receive a VAD because of failure to separate from the heart-lung machine. Postcardiectomy refers to those patients with the inability to wean from cardiopulmonary bypass secondary to left, right, or biventricular failure.
 - **Device Malfunction:** Includes those patients who are currently VAD supported and are experiencing device failure.
 - **End of (device) Life:** Mechanical device pump has reached functional life expectancy and requires replacement.
 - **Salvage:** Moribund patients unresponsive to medical interventions.
-
-

SEQ. #: 3910

Long Name: VAD-Implant Type #2

Short Name: VImpTy2

Definition: Indicate the second type of ventricular assist device implanted.

Intent/Clarification:

- Right VAD (RVAD) - Right Ventricular Assist Device
 - Left VAD (LVAD) - Left Ventricular Assist Device
 - Biventricular VAD (BiVAD) - Biventricular Assist Device
 - Total Artificial Heart (TAH)
-
-

SEQ. #: 3915

Long Name: VAD-Device #2

Short Name: VProdTy2

Definition: Indicate the specific product #2 implanted. Implant defined as physical placement of the VAD.

Intent/Clarification:

Choose the device type from the device list.

SEQ. #: 3920

Long Name: VAD-Implant Date #2

Short Name: VImpDt2

Definition: Indicate the date the VAD #2 was implanted.

Intent/Clarification:

Required date format: mm/dd/yyyy

SEQ. #: 3925

Long Name: VAD-Implant Unique Device Identifier (UDI) #2

Short Name: VImpUDI2

Definition: Indicate the device UDI if available, otherwise leave blank.

Intent/Clarification:

This is a unique identifier that will be on each device. It may not be available immediately. If not available leave blank. Please refer to the FDA web site for detailed explanation of UDI.

https://www.fda.gov/MedicalDevices/DeviceRegulationandGuidance/UniqueDeviceIdentification/default.htm?utm_source=Members%E2%80%90Only+Updates&utm_campaign=c7c1e8c870%E2%80%90%20Proposed+Rules+7+5+2012&utm_medium=email

SEQ. #: 3930

Long Name: VAD-Explant #2

Short Name: VExp2

Definition: Indicate if the VAD #2 was explanted. Explant is defined as physical removal of the VAD.

Intent/Clarification:

- Yes, not during this procedure
- Yes, during this procedure
- No

SEQ. #: 3935

Long Name: VAD-Explant Reason #2

Short Name: VExpRsn2

Definition: Indicate the reason the VAD #2 was explanted.

Intent/Clarification:

- **Cardiac Transplant** -VAD was explanted for cardiac transplant.
- **Recovery** -VAD was removed after cardiac recovery.
- **Device Transfer** -VAD was explanted in order to implant another assist device.
- **Device-Related Infection** - An infection within the pump pocket, driveline, VAD endocarditis, or other infection requiring explantation of the VAD. The body of the VAD has an active infection requiring removal to eliminate the infection. "Device-related infections" are defined as positive culture in the presence of leukocytosis, and/or fever requiring medical or surgical intervention.
- **Device Malfunction** -The VAD pump itself is not functioning properly causing hemodynamic compromise, and/or requiring immediate intervention or VAD replacement.
- **End of (device) Life** -Mechanical device pump has reached functional life expectancy and requires replacement.

Note: Code "No" if the patient expires with the VAD in place; the VAD was not explanted.

SEQ. #: 3940

Long Name: VAD-Explant Date #2

Short Name: VExpDt2

Definition: Indicate the date the VAD #2 was explanted.

Intent/Clarification:

Required date format: mm/dd/yyyy

SEQ. #: 3950

Long Name: VAD-Implant #3

Short Name: VImp3

Definition: Indicate whether a third ventricular assist device was implanted.

Intent/Clarification:

SEQ. #: 3955

Long Name: VAD-Implant Timing #3

Short Name: VADImpTmg3

Definition: Indicate timing of VAD #3 insertion.

Intent/Clarification:

- Pre-operative (during the same hospitalization but not the same OR trip as the cardiovascular surgical procedure.
 - Stand-alone VAD procedure-this was the only procedure performed.
 - In conjunction with the cardiovascular surgical procedure (same trip to the OR)-planned. In conjunction with a CV surgical procedure and planned before surgery, consent in the chart.
 - In conjunction with the cardiovascular surgical procedure (same trip to the OR)-unplanned. In conjunction with a CV surgical procedure and not planned before surgery.
 - Post-operative (after the surgical procedure during reoperation).
- -----

SEQ. #: 3960

Long Name: VAD-Indication for this VAD #3

Short Name: VADInd3

Definition: Indicate the reason for implanting a Ventricular Assist Device (VAD)#3 during this hospitalization.

Intent/Clarification:

- **Bridge to Transplantation:** Includes those patients who are supported with a VAD until a heart transplant is possible.
- **Bridge to Recovery:** Includes those patients who are expected to have ventricular recovery. (i.e. Myocarditis patients, viral cardiomyopathies, AMI w/revascularization, post-transplant reperfusion injury).
- **Destination:** Includes those patients where a heart transplant is not an option. The VAD is placed for permanent life sustaining support.
- **Post Cardiectomy Ventricular Failure:** Includes those postcardiotomy patients who receive a VAD because of failure to separate from the heart-lung machine. Postcardiotomy refers to those patients with the inability to wean from cardiopulmonary bypass secondary to left, right, or biventricular failure.
- **Device Malfunction:** Includes those patients who are currently VAD supported and are experiencing device failure.

- **End of (device) Life:** Mechanical device pump has reached functional life expectancy and requires replacement.
 - **Salvage:** Moribund patients unresponsive to medical interventions.
-
-

SEQ. #: 3965

Long Name: VAD-Implant Type #3

Short Name: VImpTy3

Definition: Indicate the third type of ventricular assist device implanted.

Intent/Clarification:

- Right VAD (RVAD) - Right Ventricular Assist Device
 - Left VAD (LVAD) - Left Ventricular Assist Device
 - Biventricular VAD (BiVAD) - Biventricular Assist Device
 - Total Artificial Heart (TAH)
-
-

SEQ. #: 3970

Long Name: VAD-Device #3

Short Name: VProdTy3

Definition: Indicate the specific product #3 implanted. Implant defined as physical placement of the VAD.

Intent/Clarification:

Choose the device type from the device list.

SEQ. #: 3975

Long Name: VAD-Implant Date #3 **Short Name:** VImpDt3

Definition: Indicate the date the VAD #3 was implanted.

Intent/Clarification:

Required date format: mm/dd/yyyy

SEQ. #: 3980

Long Name: VAD-Implant Unique Device Identifier (UDI) #3 **Short Name:** VImpUDI3

Definition: Indicate the device UDI if available, otherwise leave blank.

Intent/Clarification:

This is a unique identifier that will be on each device. It may not be available immediately. If not available leave blank. Please refer to the FDA web site for detailed explanation of UDI.

https://www.fda.gov/MedicalDevices/DeviceRegulationandGuidance/UniqueDeviceIdentification/default.htm?utm_source=Members%E2%80%90Only+Updates&utm_campaign=c7c1e8c870%E2%80%90%20Proposed_Rules_7_5_2012&utm_medium=email

SEQ. #: 3985

Long Name: VAD-Explant #3

Short Name: VExp3

Definition: Indicate if the VAD #3 was explanted. Explant is defined as physical removal of the VAD.

Intent/Clarification:

- Yes, not during this procedure
 - Yes, during this procedure
 - No
-
-

SEQ. #: 3990

Long Name: VAD-Explant Reason #3

Short Name: VExpRsn3

Definition: Indicate the reason the VAD #3 was explanted.

Intent/Clarification:

- **Cardiac Transplant** -VAD was explanted for cardiac transplant.
- **Recovery** -VAD was removed after cardiac recovery.
- **Device Transfer** -VAD was explanted in order to implant another assist device.
- **Device-Related Infection** - An infection within the pump pocket, driveline, VAD endocarditis, or other infection requiring explantation of the VAD. The body of the VAD has an active infection requiring removal to eliminate the infection. "Device-related infections" are defined as positive culture in the presence of leukocytosis, and/or fever requiring medical or surgical intervention.
- **Device Malfunction** -The VAD pump itself is not functioning properly causing hemodynamic compromise, and/or requiring immediate intervention or VAD replacement.
- **End of (device) Life** -Mechanical device pump has reached functional life expectancy and requires replacement.

Note: Code "No" if the patient expires with the VAD in place; the VAD was not explanted.

SEQ. #: 3995

Long Name: VAD-Explant Date #3

Short Name: VExpDt3

Definition: Indicate the date the VAD #3 was explanted.

Intent/Clarification:

Required date format: mm/dd/yyyy

Other Cardiac Procedures

SEQ. #: 4030

Long Name: Other Card-ASD Repair - PFO Type

Short Name: OCarASDPFO

Definition: Indicate whether a patent foramen ovale (PFO) was repaired.

Intent/Clarification:

Normally, the opening between the left and right atria closes before birth, but if it does not, the child is born with a hole in this area called patent foramen ovale (PFO). Other types of atrial septal defects occur, most commonly, secundum atrial septal defects, which account for about 70 percent of all ASDs and occur in the middle of the atrial septum.

PFO (Patent Foramen Ovale): Small interatrial communication in the region of the foramen ovale characterized by no deficiency of the septum primum and a normal limbus with no deficiency of the septum secundum.

SEQ. #: 4035

Long Name: Other Card-ASD Repair - Secundum Or Sinus Venosus

Short Name: OCarASDSec

Definition: Indicate whether a secundum or sinus venosus ASD was repaired.

Intent/Clarification:

Atrial Septal Defect (ASD) is closed with/without patch. During normal development of the heart, there is an opening in the atrial septum. ASDs in the upper part of the atrial septum (called sinus venosus) where the superior vena cava and right atrium join and can involve the right upper pulmonary vein.

- **Secundum:** An ASD confined to the region of the fossa ovalis; it's most common etiology is a deficiency of the septum primum, but deficiency of the limbus or septum secundum may also contribute.
- **Sinus Venosus:** An ASD with a vena cava or pulmonary vein (or veins) that overrides the atrial septum or the superior interatrial fold (septum secundum) producing an interatrial or anomalous veno-atrial communication. Although the term sinus venosus atrial septal defect is commonly used; the lesion is more properly termed a sinus venosus communication because, while it functions as an interatrial communication, this lesion is not a defect of the true atrial septum.

When the Mitral Valve procedure is performed via a trans-septal incision the closure of the septum should not be coded as an ASD repair.

SEQ. #: 4040

Long Name: Other Card-AFib Intracardiac Lesions

Short Name: OCarAFibIntraLes

Definition: Indicate whether intracardiac lesions were created for the purpose of AFib ablation.

Intent/Clarification:

Lesions created inside the heart (i.e. Maze procedures; lesions to mitral annulus; etc). Intracardiac procedures carry a higher risk.

SEQ. #: 4045

Long Name: Other Card-AFib Epicardial Lesions

Short Name: OCarAFibEpLes

Definition: Indicate whether epicardial lesions were created for the purpose of AFib ablation.

Intent/Clarification:

Lesions created on the outside surface of the heart (i.e. pulmonary vein isolation with or without connection to the left atrial appendage).

SEQ. #: 4050

Long Name: Other Card-Atrial Appendage Procedure

Short Name: OCarAAProc

Definition: Indicate whether atrial appendage ligation/exclusion was performed.

Intent/Clarification: This should also be coded in the AFib section if done in conjunction with creation of lesions for AFib ablation

- RAA – Right Atrial Appendage
- LAA – Left Atrial Appendage
- Both – Right and Left Atrial Appendage
- No

FAQ September 2017: If a patient has an extensive atrial fibrillation procedure and an LAA clipping how would we include the atrial ligation/exclusion method and device?
Answer: Multiple sequence numbers will be required to code the ligation/exclusion and device.

First capture sequence numbers 2140 and 2145 to open section M and M-1.

Then capture the atrial appendage procedure in sequence 4050; then ligation/exclusion method in sequence 4051 and Model in sequence number 4052 as well as the UDI, if available, in sequence 4053.

Code the extensive ablation procedure in section M-1 as described by your physician including lesion number 7 to complete the LAA.

SEQ. #: 4051

Long Name: Other Card-Atrial Appendage Ligation/Exclusion Method

Short Name: OCarAAMeth

Definition: Indicate the method used to ligate/exclude the atrial appendage

Intent/Clarification:

- Intra-atrial over-sewing
- Epicardial suture ligation
- Amputation with over-sewing
- Stapler (cutting)
- Stapler (noncutting)
- Epicardially applied occlusion device

FAQ September 2017: If a patient has an extensive atrial fibrillation procedure and an LAA clipping how would we include the atrial ligation/exclusion method and device?

Answer: Multiple sequence numbers will be required to code the ligation/exclusion and device.

First capture sequence numbers 2140 and 2145 to open section M and M-1.

Then capture the atrial appendage procedure in sequence 4050; then ligation/exclusion method in sequence 4051 and Model in sequence number 4052 as well as the UDI, if available, in sequence 4053.

Code the extensive ablation procedure in section M-1 as described by your physician including lesion number 7 to complete the LAA.

SEQ. #: 4052

Long Name: Other Card-Atrial Appendage Ligation/Exclusion Model

Short Name: OCarAAModel

Definition: Indicate the epicardial occlusion device model used

Intent/Clarification:

Capture commercially produced exclusion devices:

- AtriClip
- Lariat
- Other

FAQ September 2017: If a patient has an extensive atrial fibrillation procedure and an LAA clipping how would we include the atrial ligation/exclusion method and device?

Answer: Multiple sequence numbers will be required to code the ligation/exclusion and device.

First capture sequence numbers 2140 and 2145 to open section M and M-1.

Then capture the atrial appendage procedure in sequence 4050; then ligation/exclusion method in sequence 4051 and Model in sequence number 4052 as well as the UDI, if available, in sequence 4053.

Code the extensive ablation procedure in section M-1 as described by your physician including lesion number 7 to complete the LAA.

FAQ September 2017: Is there supposed to be a drop down list of atrial ligation devices in this field?

Answer: No, there is no drop down list.

SEQ. #: 4053

Long Name: Other Card-Atrial Appendage Ligation/Exclusion UDI

Short Name: OCarAAUDI

Definition: Indicate the Unique Device Identifier of the epicardial occlusion device

Intent/Clarification:

This is a unique identifier that will be on each device. It may not be available immediately. If not available leave blank. Please refer to the FDA web site for detailed explanation of UDI.

https://www.fda.gov/MedicalDevices/DeviceRegulationandGuidance/UniqueDeviceIdentification/default.htm?utm_source=Members%E2%80%90Only+Updates&utm_campaign=c7c1e8c870%E2%80%90%20Proposed_Rules_7_5_2012&utm_medium=email

FAQ September 2017: If a patient has an extensive atrial fibrillation procedure and an LAA clipping how would we include the atrial ligation/exclusion method and device?

Answer: Multiple sequence numbers will be required to code the ligation/exclusion and device.

First capture sequence numbers 2140 and 2145 to open section M and M-1.

Then capture the atrial appendage procedure in sequence 4050; then ligation/exclusion method in sequence 4051 and Model in sequence number 4052 as well as the UDI, if available, in sequence 4053.

Code the extensive ablation procedure in section M-1 as described by your physician including lesion number 7 to complete the LAA.

SEQ. #: 4055

Long Name: Other Card-Arrhythmia Device Surgery

Short Name: OCarACD

Definition: Indicate which arrhythmia correction device was surgically placed in conjunction with the primary surgical procedure.

Intent/Clarification:

- **Permanent Pacemaker:** An internal electronic generator that controls the heart rate
- **Permanent Pacemaker with Cardiac Resynchronization Technique (CRT-P):** An internal permanent pacemaker that uses biventricular electrical stimulation to synchronize ventricular contraction
- **Implantable Cardioverter Defibrillator (ICD):** An internal device that defibrillates the heart
- **ICD with CRT (CRT-D):** An internal ICD that uses biventricular electrical stimulation to synchronize ventricular contraction
- **Implantable recorder**
- **None**

SEQ. #: 4060

Long Name: Other Card-Lead Insertion

Short Name: OCarLeadInsert

Definition: Indicate whether lead(s) insertion was performed. Do not capture temporary lead placement.

Intent/Clarification:

These include leads for pacemakers, implantable defibrillators or combination devices.

- Yes
- No

Do not capture leads placed for temporary pacemakers.

SEQ. #: 4065

Long Name: Other Card-Arrhythmia Correction Surgery-Lead Extraction

Short Name: OCarACDLE

Definition: Indicate whether procedure included lead extraction for a device intended to treat cardiac arrhythmias.

Intent/Clarification:

- Yes, planned
- Yes, planned due to surgical complication
- Yes, unplanned due to unsuspected disease or anatomy
- No

SEQ. #: 4070

Long Name: Other Card-Congenital

Short Name: OCarCong

Definition: Indicate whether the patient had a congenital defect repair either in conjunction with, or as the primary surgical procedure. Do not include bicuspid Aortic valve or PFO here as these are captured elsewhere.

Intent/Clarification:

Indicate if a congenital procedure was performed.

- Yes
- No

SEQ. #: 4075

Long Name: Other Card-LVA

Short Name: OCarLVA

Definition: Indicate whether the patient had a Left Ventricular Aneurysm Repair either in conjunction with, or as the primary surgical procedure.

Intent/Clarification:

Indicate if a LV aneurysm repair was performed.

- Yes
- No

SEQ. #: 4080

Long Name: Other Card-Myocardial Stem Cell Therapy

Short Name: OCarStemCell

Definition: Indicate whether myocardial stem cell procedure was performed.

Intent/Clarification:

Indicate if regenerative stem cell therapy used for cardiac repair was performed.

- Yes
- No

SEQ. #: 4085

Long Name: Other Card-Pulmonary Thromboembolectomy **Short Name:**

OCPulThromDis

Definition: Indicate whether the patient had surgery for pulmonary thromboembolic disease.

Intent/Clarification:

Indicate if an embolectomy and endarterectomy was performed.

- Yes, Acute
- Yes, Chronic

SEQ. #: 4090

Long Name: Other Card-Subaortic Stenosis Resection

Short Name: OCarSubaStenRes

Definition: Indicate whether resection of subaortic stenosis was performed.

Intent/Clarification:

Subaortic stenosis (or subvalvular aortic stenosis) is a narrowing of the area below the aortic valve. This may vary from a thin layer of extra tissue to large bundles of heart muscle.

This can be performed alone or in conjunction with an aortic valve procedure.

SEQ. #: 4100

Long Name: Other Card-Subaortic Stenosis Resection Type

Short Name: OCarSubaStenResTy

Definition: Indicate the type of subaortic stenosis.

Intent/Clarification:

- Muscle
 - Ring
 - Membrane
 - Web
 - Not reported
- -----

SEQ. #: 4105

Long Name: Other Card-Surgical Ventricular Restoration

Short Name: OCarSVR

Definition: Indicate whether the patient had a Surgical Ventricular Restoration either in conjunction with, or as the primary surgical procedure. Surgical Ventricular Restorations are procedures that restore the geometry of the heart after an anterior MI. They include the Dor procedure or the SAVER procedure. This SVR procedure is distinct from an anterior left ventricular aneurysmectomy (LVA) and from a Batista procedure (left ventricular volume reduction procedure).

Intent/Clarification: Used to treat congestive heart failure caused by myocardial infarction (heart attack). The goal of the SVR is to restore the heart to a more normal size and shape, therefore improving function.

SEQ. #: 4110

Long Name: Other Card-Transmyocardial Laser Revascularization

Short Name: OCarLasr

Definition: Indicate whether the patient underwent the creation of multiple channels in left ventricular myocardium with a laser fiber either in conjunction with, or as the primary surgical procedure.

Intent/Clarification:

A laser is used to make small transmural perforations in the heart. These channels allow for blood to enter the myocardium directly from the ventricle chamber or through communications with the native coronary circulations. Used primarily in areas of the heart where bypass grafting is not feasible, to improve collateralization of circulation.

SEQ. #: 4115

Long Name: Other Card-Tumor

Short Name: OCTumor

Definition: Indicate whether the patient had resection of an intracardiac tumor.

Intent/Clarification:

Cardiac tumors are abnormal growths that can occur in the heart or on the heart valves. The tumors can be malignant or benign. Tumors can begin in the heart or in another part of the body. Tumors cause problems because of their size and location and can embolize:

- Myxoma
 - Fibroelastoma
 - Hypernephroma
 - Sarcoma
 - Other
 - No
-
-

SEQ. #: 4120

Long Name: Other Card-Card Tx

Short Name: OCarCrTx

Definition: Indicate whether the patient had a Heterotopic or Orthotopic heart transplantation either in conjunction with, or as the primary surgical procedure.

Intent/Clarification:

- **Heterotopic Transplant** – The transplant recipient’s heart is not explanted. A donor’s heart is implanted as a “piggy back” to the patient’s native heart. The donor heart acts as an assist pump for the diseased heart. The patient now has two hearts.
 - **Orthotopic** – The patient’s diseased native heart is excised and replaced with a donor heart. The recipient heart is removed completely except for small cuff of right and left atrium.
-
-

SEQ. #: 4125

Long Name: Other Card-Cardiac Trauma

Short Name: OCarTrma

Definition: Indicate whether the patient had a surgical procedure for an injury due to Cardiac Trauma either in conjunction with, or as the primary surgical procedure.

Intent/Clarification: Injury to the heart such as a gunshot wound, stab wound, car accident or other trauma induced injury.

SEQ. #: 4130

Long Name: Other Card-VSD

Short Name: OCarVSD

Definition: Indicate whether the patient had a Ventricular Septal Defect Repair either in conjunction with, or as the primary surgical procedure.

Intent/Clarification: (VSD) Defect of the ventricular septum is closed with/without patch.

- Yes, congenital
 - Yes, acquired
 - No
- -----

SEQ. #: 4135

Long Name: Other Card-Other

Short Name: OCarOthr

Definition: Indicate whether the patient had another cardiac procedure performed either in conjunction with, or as the primary surgical procedure that is not included within this section.

Intent/Clarification: The following is a guideline for assessing which procedures to capture for Other Card - Other:

Code procedures that have a high likelihood of negatively impacting a patient's outcome (survival, quality of life, ability to recover) and/or prolong the patient's length of stay. You do not want to code this if minor procedures were done in conjunction with a CABG or a Valve and lose the patient in the analysis of isolated procedures!

Due to the difficulty of publishing a complete list of procedures to include and not to include in this field, the STS encourages sites to submit the procedure in question as a clinical question. Whether to include or not to include a procedure will be dealt with on a procedure by procedure basis.

Atrial Fibrillation Procedures

SEQ. #: 4191

Long Name: AFib Lesion Location

Short Name: OCarAFibLesLoc

Definition: Indicate the location of the majority of lesions created to treat atrial fibrillation.

Intent/Clarification:

Indicate whether the lesions created were primarily epicardial or primarily intracardiac.

SEQ. #: 4200

Long Name: Atrial Fibrillation Surgical Procedure-Method of Lesion Creation - Radio Frequency

Short Name: OCarAFibMethRad

Definition: Indicate whether the method used to create the lesion(s) for the AFib ablation procedure included radio frequency.

Intent/Clarification: Radiofrequency energy uses an alternating current resulting in thermal injury to disrupt atrial fibrillation pathways. These probes can be applied to either endocardial or epicardial heart surfaces to create transmural linear lesions that block atrial conduction.

SEQ. #: 4205

Long Name: Atrial Fibrillation Surgical Procedure-Method of Lesion Creation - Radio Frequency - Bipolar

Short Name: OCarAFibMethRadBi

Definition: Indicate whether the radiofrequency method used to create the lesion(s) for the AFib ablation was bipolar.

Intent/Clarification: If radiofrequency was used, was it bipolar.

SEQ. #: 4210

Long Name: Atrial Fibrillation Surgical Procedure-Method of Lesion Creation - Cut-And-Sew

Short Name: OCarAFibMethCAS

Definition: Indicate whether the method used to create the lesion(s) for the AFib ablation procedure included cut-and-sew.

Intent/Clarification: A technically difficult procedure where the lesions are created using a scalpel creating surgical incisions in the atrium and sewing them to create scars that inhibit re-entry rhythms.

SEQ. #: 4215

Long Name: Atrial Fibrillation Surgical Procedure-Method of Lesion Creation - Cryo

Short Name: OCarAFibMethCryo

Definition: Indicate whether the method used to create the lesion(s) for the A-Fib ablation procedure included cryoablation.

Intent/Clarification: Cryoablation used to restore normal heart rhythm. It freezes the heart tissue that triggers an irregular heartbeat. Cryoablation is performed with a nitrous oxide cooled probe that, when applied to atrial tissue, produces transmural lesions that block atrial conduction.

SEQ. #: 4240

Long Name: Lesions Documented

Short Name: OCarLesDoc

Definition: Indicate whether the lesions created during the atrial fibrillation surgery are documented.

Intent/Clarification: Indicate whether the lesions were documented in the medical record.

- Yes – there is documentation of lesion lines
- No – documentation is not available for the lesion lines used in the ablation procedure.

SEQ. #: 4250

Long Name: AFib Lesion Location - Bilateral Pulmonary Vein Isolation

Short Name: AFibLes1

Definition: Indicate whether the AFib lesion was pulmonary vein isolation.

Intent/Clarification:

Pulmonary vein ablation is a treatment for atrial fibrillation in which both the left and right pulmonary veins are ablated.

Refer to the pictures and the corresponding numbers graph on the data collection form.

SEQ. #: 4255

Long Name: AFib Lesion Location - Box Lesion Only

Short Name: AFibLes2

Definition: Indicate whether the AFib lesion was a box lesion

Intent/Clarification:

Box is a treatment for atrial fibrillation was performed.

Refer to the pictures and the corresponding numbers graph on the data collection form.

SEQ. #: 4260

Long Name: AFib Lesion Location - Inferior Pulmonary Vein Connecting Lesion

Short Name: AFibLes3a

Definition: Indicate whether the AFib lesion was an Inferior Pulmonary Vein Connecting Lesion

Intent/Clarification:

Refer to the pictures and the corresponding numbers graph on the data collection form.

SEQ. #: 4265

Long Name: AFib Lesion Location - Superior Pulmonary Vein Connecting Lesion

Short Name: AFibLes3b

Definition: Indicate whether the AFib lesion was a Superior Pulmonary Vein Connecting Lesion

Intent/Clarification:

Refer to the pictures and the corresponding numbers graph on the data collection form.

SEQ. #: 4270

Long Name: AFib Lesion Location - Posterior Mitral Annular Line Lesion

Short Name: AFibLes4

Definition: Indicate whether the AFib lesion was a Posterior Mitral Annular Line

Intent/Clarification: Indicate whether the A-fib lesion was a Pulmonary Vein Connecting Lesion to Posterior Mitral Annulus lesion.

Refer to the pictures and the corresponding numbers graph on the data collection form.

SEQ. #: 4275

Long Name: AFib Lesion Location - Pulmonary Vein Connecting Lesion to Anterior Mitral Annulus

Short Name: AFibLes5

Definition: Indicate whether the AFib lesion was a - Pulmonary Vein Connecting Lesion to Anterior Mitral Annulus lesion.

Intent/Clarification: Indicate whether the afib lesion was a - Pulmonary Vein Connecting Lesion to Anterior Mitral Annulus lesion.

Refer to the pictures and the corresponding numbers graph on the data collection form.

SEQ. #: 4280

Long Name: AFib Lesion Location - Mitral Valve Annular Lesion

Short Name: AFibLes6

Definition: Indicate whether the AFib lesion was a Mitral Valve Cryo Lesion

Intent/Clarification:

Refer to the pictures and the corresponding numbers graph on the data collection form.

SEQ. #: 4285

Long Name: AFib Lesion Location - LAA Ligation/Removal/Obliteration

Short Name: AFibLes7

Definition: Indicate whether the left Atrial Appendage was ligated or removed

Intent/Clarification:

Refer to the pictures and the corresponding numbers graph on the data collection form.

FAQ September 2017: If a patient has an extensive atrial fibrillation procedure and an LAA clipping how would we include the atrial ligation/exclusion method and device?

Answer: Multiple sequence numbers will be required to code the ligation/exclusion and device.

First capture sequence numbers 2140 and 2145 to open section M and M-1.

Then capture the atrial appendage procedure in sequence 4050; then ligation/exclusion method in sequence 4051 and Model in sequence number 4052 as well as the UDI, if available, in sequence 4053.

Code the extensive ablation procedure in section M-1 as described by your physician including lesion number 7 to complete the LAA.

SEQ. #: 4290

Long Name: AFib Lesion Location - Pulmonary Vein to LAA Lesion

Short Name: AFibLes8

Definition: Indicate whether the AFib lesion was a Pulmonary Vein to LAA lesion

Intent/Clarification:

Refer to the pictures and the corresponding numbers graph on the data collection form.

SEQ. #: 4295

Long Name: AFib Lesion Location - Intercaval Line to Tricuspid Annulus ('T' lesion)

Short Name: AFibLes9

Definition: Indicate whether the AFib lesion was an Intercaval Line to Tricuspid Annulus ('T' lesion)

Intent/Clarification:

Refer to the pictures and the corresponding numbers graph on the data collection form.

SEQ. #: 4300

Long Name: AFib Lesion Location - Tricuspid Cryo Lesion, Medial (10)

Short Name: AFibLes10

Definition: Indicate whether the AFib lesion was a Tricuspid Cryo Lesion, Medial (10)

Intent/Clarification:

Refer to the pictures and the corresponding numbers graph on the data collection form.

SEQ. #: 4305

Long Name: AFib Lesion Location - Intercaval Line (SVC and IVC)

Short Name: AFibLes11

Definition: Indicate whether the AFib lesion was an Intercaval Line

Intent/Clarification:

Refer to the pictures and the corresponding numbers graph on the data collection form.

SEQ. #: 4310

Long Name: AFib Lesion Location - Tricuspid Annular Line to RAA

Short Name: AFibLes12

Definition: Indicate whether the AFib lesion was a Tricuspid Annular Line to RAA lesion

Intent/Clarification:

Refer to the pictures and the corresponding numbers graph on the data collection form.

SEQ. #: 4315

Long Name: AFib Lesion Location - Tricuspid Cryo Lesion (13)

Short Name: AFibLes13

Definition: Indicate whether the Afib lesion was a Tricuspid Cryo Lesion (13)

Intent/Clarification:

Refer to the pictures and the corresponding numbers graph on the data collection form.

SEQ. #: 4320

Long Name: AFib Lesion Location - RAA Ligation/Removal/Obliteration

Short Name: AFibLes14

Definition: Indicate whether the Right Atrial Appendage was ligated or removed

Intent/Clarification:

Refer to the pictures and the corresponding numbers graph on the data collection form.

SEQ. #: 4325

Long Name: AFib Lesion Location - RAA Lateral Wall (Short)

Short Name: AFibLes15a

Definition: Indicate whether the Afib lesion was a RAA Lateral Wall (Short) lesion

Intent/Clarification:

Refer to the pictures and the corresponding numbers graph on the data collection form.

SEQ. #: 4330

Long Name: AFib Lesion Location - RAA Lateral Wall to 'T' Lesion

Short Name: AFibLes15b

Definition: Indicate whether the Afib lesion was a RAA Lateral Wall to 'T' Lesion

Intent/Clarification:

Refer to the pictures and the corresponding numbers graph on the data collection form.

SEQ. #: 4336

Long Name: AFib Lesion Location - Coronary Sinus Lesion

Short Name: AFitLesCSL

Definition: Indicate whether the Afib lesion was a Coronary Sinus Lesion.

Intent/Clarification:

Refer to the pictures and the corresponding numbers graph on the data collection form.

Aorta And Aortic Root Procedures

Please use the following diagram for reference in section M2

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)
- N. Zone 11 (external iliacs)

SEQ. #: 4500

Long Name: Family History Of Disease Of The Aorta

Short Name: FamHistAorta

Definition: Indicate whether there is a family history of disease of the aorta

Intent/Clarification:

For the purposes of this database (and published guidelines), family history means any alive or dead first-degree relative ('FDR': sibling, parent, child) with either a thoracic

aortic aneurysm (include 'dilated' or 'enlarged' aorta), or aortic dissection/rupture. Abdominal aneurysms should be excluded, as they are typically not familial in nature. Thoracic location is sometimes described as 'near' or 'above' the heart, or 'in the chest'. For this database, in the case of family history of an unexplained death of a first-degree family member select 'unknown'. Patients with a family history of thoracic aneurysm and especially with a history of dissection/rupture who require aortic surgery may have more fragile aortic tissue or require a more extensive procedure, which could affect procedural outcomes.

Excludes isolated abdominal aortic aneurysm/dissections.

SEQ. #: 4505

Long Name: Genetic History

Short Name: PatGenHist

Definition: Indicate the genetic history of the patient

Intent/Clarification:

Indicate whether or not the patient has a history of any of the listed (well-known) genetically triggered thoracic aortic conditions.

- Non-specified' Familial Aneurysm: patients in whom another family member(s) had thoracic aneurysm but no specific gene mutation was identified when tested.
- Other: patient has been told they have a relevant gene mutation related to thoracic aneurysm but did not match any of the choices listed above; these will include known pathogenic mutations familiar to specialists but not associated with a 'named' syndrome like the other choices.
- None: patient has undergone genetic testing with no positive findings.
- Unknown: no known syndromic/genetic diagnosis, but has not specifically been tested for pathogenic mutations.

The diagnosis has been made and is documented in the medical record by clinical or genetic testing.

SEQ. #: 4510

Long Name: Prior Aortic Intervention

Short Name: PriorAorta

Definition: Indicate whether the patient had prior aortic intervention

Intent/Clarification:

Includes both open surgical and/or endovascular (stent) intervention of any part of the aorta.

SEQ. #: 4520

Long Name: Prior Aortic Intervention - Previous Repair - Root

Short Name: PriorRepRoot

Definition: Indicate whether the prior intervention involved the aortic root

Intent/Clarification:

The aortic root is the 'sinus' segment of the aorta that immediately exits the heart, and contains the aortic valve and coronary artery origins. It ends anatomically at the sinotubular junction where the tubular ascending aorta begins.

The region of the aorta designated in zone 0 from the below the sinotubular junction.

SEQ. #: 4521

Long Name: Prior Aortic Intervention - Previous Repair Type - Root

Short Name: PriorRepTyRoot

Definition: Indicate the type of prior root repair

Intent/Clarification:

In this location, surgery would either name aortic root replacement as well as designations of 'mechanical' or biological', also called "Bentall" procedures and also includes 'valve-sparing' root procedures ("David", "Re-implantation", "Yacoub", "Remodeling", "Florida Sleeve).

The only applicable choice for root repair type is open.

SEQ. #: 4522

Long Name: Prior Aortic Intervention - Repair Failure - Root

Short Name: PriorFailRoot

Definition: Indicate whether there is failure of the prior root repair

Intent/Clarification:

Either a secondary 'false' pseudo-aneurysm has developed in or near the previous aortic root repair, or a portion of preserved aortic root tissue (typically the coronary origins or 'buttons') has become aneurysmal.

There are four areas of prior root failure. These include proximal, distal and right coronary and left coronary button suture line issues.

SEQ. #: 4523

Long Name: Prior Aortic Intervention - Disease Progression - Root

Short Name: PriorProgRoot

Definition: Indicate whether there is progression of disease following the prior root repair

Intent/Clarification:

If only a portion of the aortic root (typically the non-coronary sinus) was replaced during the initial root procedure, aneurysmal progression of the left and or right coronary sinuses may have occurred. Also, development of coronary button aneurysms would be considered progression of disease.

SEQ. #: 4525

Long Name: Prior Aortic Intervention - Previous Repair - Ascending

Short Name: PriorRepAsc

Definition: Indicate whether the prior intervention involved the ascending aorta

Intent/Clarification:

The ascending aorta is also called the tubular ascending segment and is the portion above the aortic root ('sinus segment') beginning at the sinotubular junction and extending to the first aortic arch vessel (innominate or brachiocephalic artery).

SEQ. #: 4526

Long Name: Prior Aortic Intervention - Previous Repair Type - Ascending

Short Name: PriorRepTyAsc

Definition: Indicate the type of prior ascending aorta repair

Intent/Clarification:

Most simply classified as ascending aortic replacement with a prosthetic graft, but also includes ascending aortic resection (removal of the aneurysm with end-to-end proximal and distal aortic connection), aortoplasty (reduction of the diameter of the ascending aorta with sutures or by removing a longitudinal segment).

SEQ. #: 4527

Long Name: Prior Aortic Intervention - Repair Failure - Ascending

Short Name: PriorFailAsc

Definition: Indicate whether there is failure of the prior ascending repair

Intent/Clarification:

A situation where there has been a previous replacement or aortoplasty of the (tubular) ascending aortic segment in which the patient has manifested a pseudo-aneurysm and/or further aortic expansion, contained rupture of the proximal or distal suture line.

The region of the aorta designated in zone 0 from the sinotubular junction to distal ascending aorta.

SEQ. #: 4528

Long Name: Prior Aortic Intervention - Disease Progression - Ascending

Short Name: PriorProgAsc

Definition: Indicate whether there is progression of disease following the prior ascending aorta repair/

Intent/Clarification:

A situation could arise where the ascending aorta was previously replaced with a tube graft, but a small segment of the ascending aorta (usually the ascending to proximal arch transition) was not removed, and has subsequently become aneurysmal, now requiring intervention. Another scenario is if an ascending (reduction) aortoplasty was employed as the previous repair, and this segment has become aneurysmal to an extent requiring intervention currently.

SEQ. #: 4530

Long Name: Prior Aortic Intervention - Previous Repair - Arch

Short Name: PriorRepArch

Definition: Indicate whether the prior intervention involved the aortic arch

Intent/Clarification:

The aortic arch is the segment of aorta beyond the tubular ascending segment, and begins at the level of the first branching vessel of the aorta (typically the innominate or brachiocephalic artery), and terminating just after the last branch vessel of the aortic arch (left subclavian artery), before transitioning to the descending thoracic aorta; specifically zones 1, 2 and 3.

The region of the aorta designated in zones 1, 2, & 3 from the distal ascending to the proximal descending thoracic aorta.

SEQ. #: 4531

Long Name: Prior Aortic Intervention - Previous Repair Type - Arch

Short Name: PriorRepTyArch

Definition: Indicate the type of prior arch repair

Intent/Clarification:

Open arch repairs may include 'hemi-arch' repairs, where branch arteries are not re-implanted or bypassed, and extent of arch replacement with a graft includes a significant portion of the lesser curve (non-branched portion) of the aortic arch, as well as 'total' arch replacement (all branch vessels re-implanted or bypassed in addition to graft replacement of the aorta), or 'partial' arch replacement (one or more, but not all arch vessels re-implanted or replaced in addition to graft replacement of a portion the

aortic arch). Additionally, 'hybrid' repairs may combine surgical bypasses to one or more arch vessels with endograft (stent) repair of the aortic arch, and total endovascular arch replacement (rare) includes endovascular perfusion of arch vessels using special techniques.

SEQ. #: 4532

Long Name: Prior Aortic Intervention - Repair Failure - Arch

Short Name: PriorFailArch

Definition: Indicate whether there is failure of the prior arch repair

Intent/Clarification:

Relates to pseudo-aneurysms that have formed as part of arch repair, or failure of an endograft to 'seal' with an endoleak leading to further aortic expansion. Could also indicated a bypassed or re-implanted arch vessel failure that requires a later re-intervention.

An **endoleak** is defined as persistent blood flow in the aneurysm sac through and around the endovascular seal and is the most common complication after endovascular aneurysm repair.

SEQ. #: 4533

Long Name: Prior Aortic Intervention - Disease Progression - Arch

Short Name: PriorProgArch

Definition: Indicate whether there is progression of disease following the prior arch repair

Intent/Clarification:

Cases of partial arch replacement or hemi-arch aortic replacement where the residual aortic arch has become aneurysmal to an extent requiring re-intervention, or a re-implanted branch vessel has become aneurysmal requiring intervention.

SEQ. #: 4535

Long Name: Prior Aortic Intervention - Previous Repair - Descending

Short Name: PriorRepDesc

Definition: Indicate whether the prior intervention involved the descending aorta

Intent/Clarification:

The descending thoracic aorta begins after the aortic arch (beyond the left subclavian artery) and extends to the level of the aortic hiatus at the diaphragm.

The region of the aorta designated in zones 4 & 5 from the distal arch to the celiac arteries.

SEQ. #: 4536

Long Name: Prior Aortic Intervention - Previous Repair Type - Descending

Short Name: PriorRepTyDesc

Definition: Indicate the type of prior descending aorta repair

Intent/Clarification:

The descending thoracic aorta can be replaced with a tube graft (open surgical) or using endovascular (stent) repair. Hybrid repairs include the use of an 'elephant trunk' extension of an aortic arch repair and secondary open surgical or endograft connection to the elephant trunk extension.

SEQ. #: 4537

Long Name: Prior Aortic Intervention - Repair Failure - Descending

Short Name: PriorFailDesc

Definition: Indicate whether there is failure of the prior descending repair

Intent/Clarification:

Formation of pseudo-aneurysm or failure of an endograft repair to 'seal' or an endoleak causing aneurysm expansion at the site of previous treatment.

An **endoleak** is defined as persistent blood flow in the aneurysm sac through and around the endovascular seal and is the most common complication after endovascular aneurysm repair.

SEQ. #: 4538

Long Name: Prior Aortic Intervention - Disease Progression - Descending

Short Name: PriorProgDesc

Definition: Indicate whether there is progression of disease following the prior descending aorta repair

Intent/Clarification:

A situation where a segment of the descending thoracic aorta was previously replaced and an adjacent non-replaced segment has expanded to an extent requiring a new intervention. Also, a pre-emptive elephant trunk extension was created at the time of a previous arch repair, and the descending thoracic aorta has become large enough to complete treatment.

SEQ. #: 4540

Long Name: Prior Aortic Intervention - Previous Repair - Suprarenal Abdominal

Short Name: PriorRepSupraAb

Definition: Indicate whether the prior intervention involved the suprarenal abdominal aorta

Intent/Clarification:

The segment of aorta beginning at the level of the diaphragm and ending just below the renal artery branches. This segment includes major branches to the abdominal organs, including the celiac and superior mesenteric artery, but not the inferior mesenteric artery.

The region of the aorta designated in zone 6 & 7 from the celiac to the renal arteries.

SEQ. #: 4541

Long Name: Prior Aortic Intervention - Previous Repair Type - Suprarenal Abdominal

Short Name: PriorRepTySupraAb

Definition: Indicate the type of prior suprarenal abdominal aorta repair

Intent/Clarification:

Similar to the aortic arch, when this segment is replaced either with open surgery or with endovascular (stent) grafting, its major vessels require either re-implantation or bypass.

SEQ. #: 4542

Long Name: Prior Aortic Intervention - Repair Failure - Suprarenal Abdominal

Short Name: PriorFailSupraAb

Definition: Indicate whether there is failure of the prior suprarenal abdominal repair

Intent/Clarification:

This includes pseudo-aneurysms as well as failure of endograft 'seal' or endoleak causing continued expansion of the aorta requiring another intervention. Additionally, stenosis or occlusion of a bypassed or re-implanted visceral vessel indicates a repair failure that could mandate a re-intervention.

An **endoleak** is defined as persistent blood flow in the aneurysm sac through and around the endovascular seal and is the most common complication after endovascular aneurysm repair.

SEQ. #: 4543

Long Name: Prior Aortic Intervention - Disease Progression - Suprarenal Abdominal

Short Name: PriorProgSupraAb

Definition: Indicate whether there is progression of disease following the prior suprarenal abdominal aorta repair

Intent/Clarification:

If a portion of the supra-renal aorta was not replaced during the initial surgery (most typically proximally, near the diaphragm, during open surgery), aneurysm progression in this location could occur, requiring another intervention. Aneurysm formation of the proximal portions of the visceral vessels themselves could also occur (more likely in genetic aneurysm syndromes) and require re-intervention as well.

SEQ. #: 4545**Long Name:** Prior Aortic Intervention - Previous Repair - Infrarenal Abdominal**Short Name:** PriorReplnfraAb**Definition:** Indicate whether the prior intervention involved the infrarenal abdominal aorta**Intent/Clarification:**

This is the segment of aorta below the renal arteries, and terminating just before the bifurcation of the aorta into the common iliac arteries.

The region of the aorta designated in zone 9 the infrarenal abdominal aorta.

SEQ. #: 4546**Long Name:** Prior Aortic Intervention - Previous Repair Type - Infrarenal Abdominal**Short Name:** PriorRepTyInfraAb**Definition:** Indicate the type of prior infrarenal abdominal aorta repair**Intent/Clarification:**

The infra-renal aorta can be replaced either with open surgery or endovascular (stent) graft repair. The latter usually involves from the kidneys and branching graft into the common iliac arteries.

SEQ. #: 4547**Long Name:** Prior Aortic Intervention - Repair Failure - Infrarenal Abdominal**Short Name:** PriorFailnfraAb**Definition:** Indicate whether there is failure of the prior infrarenal abdominal repair**Intent/Clarification:**

This includes pseudo-aneurysms as well as failure of endograft 'seal' or endoleak causing continued expansion of the aorta requiring another intervention.

An **endoleak** is defined as persistent blood flow in the aneurysm sac through and around the endovascular seal and is the most common complication after endovascular aneurysm repair.

SEQ. #: 4548

Long Name: Prior Aortic Intervention - Disease Progression - Infrarenal Abdominal

Short Name: PriorProgInfraAb

Definition: Indicate whether there is progression of disease following the prior infrarenal abdominal aorta repair

Intent/Clarification:

A situation where a segment of infra-renal aorta was left behind or un-treated during a previous procedure and has now become aneurysmal to an extent requiring re-intervention.

SEQ. #: 4620

Long Name: Endoleak

Short Name: Endoleak

Definition: Indicate whether endoleak is present

Intent/Clarification:

An endoleak is defined as the presence of blood leaking through or around an endograft into the aneurysm sac resulting in perfusion and persistent pressurization of the aneurysm sac, it is the most common complication after endovascular aneurysm repair. In the case of an aortic dissection, an endoleak refers to persistent false lumen perfusion. The intent is to identify the efficacy of the procedure with the optimal therapy resulting in the absence of any endoleak.

SEQ. #: 4625

Long Name: Endoleak - Type I - Leak At Graft Attachment Site

Short Name: EndoleakTypeI

Definition: Indicate whether endoleak is type I

Intent/Clarification:

The intent is to identify the presence of a Type I endoleak. A Type I endoleak is defined as leakage of blood around a graft at the proximal or distal seal zones. This results due to a gap between the aortic wall and the endograft at either the proximal or distal seal zone.

SEQ. #: 4630

Long Name: Endoleak - Type I - Location **Short Name:** EndoleakTyILoc

Definition: Indicate the location of the type I endoleak

Intent/Clarification:

The intent is to identify the location of the Type I endoleak. A Type Ia endoleak is defined as a leak occurring at the proximal seal zone. A Type Ib endoleak is defined as a leak occurring at the distal seal zone. A Type Ic endoleak is defined as a non-occluded iliac artery in patients with an aorto-uni-iliac device with a patent femoral-femoral bypass.

SEQ. #: 4635**Long Name:** Endoleak - Type II - Aneurysm Sac Filling Via Branch Vessel**Short Name:** EndoleakTypeII**Definition:** Indicate whether endoleak is type II**Intent/Clarification:**

The intent is to identify the presence of a Type II endoleak. A Type II endoleak is defined as retrograde filling of the aneurysm sac or false lumen in the case of dissection by aortic branch vessels (e.g. left subclavian artery, intercostal arteries, etc.).

SEQ. #: 4640**Long Name:** Endoleak - Type II - Number Of Vessels**Short Name:** EndoleakVessNum**Definition:** Indicate the number of vessels involved in the type II endoleak**Intent/Clarification:**

The intent is to identify the number of vessels providing retrograde flow into the aneurysm sac or false lumen. A Type IIa endoleak is defined as one branch vessel with retrograde flow causing an endoleak. A Type IIb endoleak is defined as more than one branch vessel with retrograde flow causing an endoleak.

SEQ. #: 4645**Long Name:** Endoleak - Type III - Leak Through Defect In Graft**Short Name:** EndoleakTypeIII**Definition:** Indicate whether endoleak is type III**Intent/Clarification:**

The intent is to identify the presence of a Type III endoleak. A Type III endoleak is defined as leakage of blood into the aneurysm sac, or false lumen in the case of dissection, due to either a gap between separate endograft components, or a defect in the fabric of the graft secondary to graft strut fracture or erosion.

SEQ. #: 4650

Long Name: Endoleak - Type III - Graft Defect Type

Short Name: EndoleakType

Definition: Indicate the graft defect type

Intent/Clarification:

The intent is to identify which type of Type III endograft exists. A Type IIIa defect (junctional separation of modular components) occurs when an endoleak occurs secondary to junctional separation of overlapping endografts. A Type IIIb defect (endograft fracture or holes) occurs when an endoleak occurs secondary to a perforation in the fabric of an endograft secondary to graft strut fracture or erosion.

SEQ. #: 4655

Long Name: Endoleak - Type IV - Leak Through Graft Fabric - Porosity

Short Name: EndoleakTypeIV

Definition: Indicate whether endoleak is type IV

Intent/Clarification:

The intent is to identify the presence of a Type IV endoleak. A Type IV endoleak is defined as the presence of an endoleak secondary to graft porosity. All other types of endoleaks must be definitively ruled out prior to selecting this diagnosis.

SEQ. #: 4660

Long Name: Endoleak - Type V - Endotension-Expansion Aneurysm Sac Without Leak

Short Name: EndoleakTypeV

Definition: Indicate whether endoleak is type V

Intent/Clarification:

The intent is to identify the presence of a Type V endoleak. A Type V endoleak, also known as endotension, is defined as persistent aneurysm expansion in the absence of a confirmed endoleak.

SEQ. #: 4665

Long Name: Aorta Infection

Short Name: Infection

Definition: Indicate whether infection is present

Intent/Clarification:

The intent is to establish the presence of a primary aortic infection (either native aorta or prosthetic graft). This can be prospectively established preoperatively with diagnostic cultures (i.e. perigraft fluid or phlegmon aspiration) or other imaging such (tagged WBC

scan or characteristic MRI or CT changes). The final diagnosis should depend on surgeon report, intraoperative cultures and pathologic data.

SEQ. #: 4670

Long Name: Aorta Infection Type **Short Name:** InfecType

Definition: Indicate the type of aortic infection

Intent/Clarification:

Intent is to establish the type of an infection within the aorta including the sinus of Valsalva and the aortic valve. Infection may involve native tissues or prosthetic graft or prosthetic valve material. Multiples infection types might be described as involving more than one type, i.e. Graft and native aorta.

SEQ. #: 4675

Long Name: Aorta Trauma **Short Name:** Trauma

Definition: Indicate whether there was aortic trauma

Intent/Clarification:

Aortic trauma will include blunt trauma (i.e. blunt aortic injury in motor vehicle accident), penetrating trauma (i.e. gun shot, stabbing, etc.), and iatrogenic trauma (i.e. endovascular catheter induced perforation or dissection). Do not include surgical complications; may include catheter trauma.

SEQ. #: 4680

Long Name: Trauma Location - Root

Short Name: TraumacRoot

Definition: Indicate whether the aortic trauma involved the root

Intent/Clarification:

Includes the sinus of Valsalva, aortic valve leaflets and aortoventricular junction.

SEQ. #: 4685

Long Name: Trauma Location - Ascending

Short Name: TraumaAsc

Definition: Indicate whether the aortic trauma involved the ascending aorta

Intent/Clarification:

Sinotubular junction to the innominate artery

SEQ. #: 4690

Long Name: Trauma Location - Arch

Short Name: TraumaArch

Definition: Indicate whether the aortic trauma involved the arch

Intent/Clarification:

Proximal aspect of the innominate artery to the distal aspect of the left subclavian artery/aortic isthmus

SEQ. #: 4695

Long Name: Trauma Location - Descending

Short Name: TraumaDesc

Definition: Indicate whether the aortic trauma involved the descending aorta

Intent/Clarification:

Aorta distal to the left subclavian to the diaphragmatic hiatus

SEQ. #: 4700

Long Name: Trauma Location - Thoracoabdominal

Short Name: TraumaThorac

Definition: Indicate whether the aortic trauma involved the thoracoabdominal aorta

Intent/Clarification:

Location of trauma includes parts of the descending thoracic aorta and abdominal aorta.

SEQ. #: 4705

Long Name: Trauma Location - Abdominal

Short Name: TraumaAbdom

Definition: Indicate whether the aortic trauma involved the abdominal aorta

Intent/Clarification:

Location of trauma includes parts of the descending thoracic aorta and abdominal aorta.
Trauma isolated to infradiaphragmatic abdominal aorta

SEQ. #: 4710

Long Name: Aorta Presentation

Short Name: Presentation

Definition: Indicate the clinical presentation

Intent/Clarification:

This is intended to define the presenting symptoms that lead to the diagnosis and operative intervention and might include: Pain, CHF, Cardiac Arrest, Syncope, Stroke, limb numbness, Paralysis, Fatigue, Infection, Weakness, Hoarseness (vocal cord dysfunction). However, some patients may be asymptomatic.

There is no specific hierarchy and the primary presentation should be indicated by the surgeon.

SEQ. #: 4715

Long Name: Aorta Primary indication **Short Name:** PrimIndic

Definition: Indicate the primary indication for intervention

Intent/Clarification:

The intent is to identify the condition/diagnosis/pathology for which surgery is being conducted and may include: Aneurysm, Dissection, Valvular Dysfunction, Obstruction, Intramural Hematoma, Infection, Stenosis, and Coarctation.

Intramural hematoma is when there is blood in the wall of the aorta but no dissection flap is visualized.

Aortic coarctation is a narrowing of the aorta and usually a congenital issue

There is no specific hierarchy and the primary presentation should be indicated by the surgeon.

FAQ October 2017: How should the primary indication be coded for the patient with a penetrating ulcer of the descending aorta?

Answer: Code aneurysm.

SEQ. #: 4720

Long Name: Aneurysm - Etiology

Short Name: AnEtiology

Definition: Indicate the aneurysm etiology

Intent/Clarification:

Choices here are Atherosclerosis, Infection, inflammatory, Connective Tissue Disorder, Penetrating Ulcer, Pseudoaneurysm, Mycotic, Traumatic transection, Intercostal visceral patch, Anastomotic site, Unknown.

Mycotic aneurysm refers to a native tissue infection.

Inflammatory refers to an autoimmune disease - Ehlers Danlos

Connective Tissue Disorder refers to Marfans, etc.

Pseudoaneurysm is an outpouching that does not involve all layers of the aortic wall.

If the patient has Bicuspid choose 4505.

FAQ October 2017: If the etiology of the aortic aneurysm is described by the surgeon as systemic hypertension, how is that coded?

Answer: Code unknown.

FAQ October 2017: If the etiology of the aortic aneurysm is described by the surgeon as degenerative, how is that coded?

Answer: Code unknown.

SEQ. #: 4725

Long Name: Aneurysm - Type

Short Name: AnType

Definition: Indicate the aneurysm type

Intent/Clarification:

Saccular aneurysm is a focal dilation of all layers of the aorta.

Fusiform aneurysm is a diffuse dilation of all layers of the aortic wall involving an extended segment.

Most aneurysms tend to be fusiform. Saccular aneurysms would be dictated as such.

SEQ. #: 4730

Long Name: Aneurysm - Rupture

Short Name: AnRupt

Definition: Indicate whether the aneurysm ruptured

Intent/Clarification:

Aneurysm rupture is a complete breakdown in the integrity of the aortic wall and if not "contained" will result in exsanguination.

SEQ. #: 4735

Long Name: Aneurysm - Rupture - Contained

Short Name: AnRuptCon

Definition: Indicate whether the rupture was contained

Intent/Clarification:

Contained rupture is a complete breakdown in the integrity of the aortic wall but is being “contained” by some clot or another structure. It is an unstable situation. When seen on CT scan, it is almost always “contained” as frank rupture is usually fatal.

SEQ. #: 4740

Long Name: Aneurysm - Location

Short Name: AnLoc

Definition: Indicate the location of the aneurysm

Intent/Clarification:

STJ is the Sino-tubular Junction and identifies the boundary between the aortic root and the ascending aorta and is marked as Zone 0 which includes everything from above the aortic root to the innominate artery, i.e.: both the aortic root and ascending aorta.

There is no specific hierarchy and choose the primary zone of maximum diameter indicated by the surgeon.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

FAQ October 2017: If the aneurysm spans more than one zone how should this be coded?

Answer: Code the most proximal zone with largest diameter.

SEQ. #: 4745

Long Name: Dissection - Timing

Short Name: DisTiming

Definition: Indicate the timing of the aortic dissection

Intent/Clarification:

The intent is to define the time interval from occurrence of dissection until presentation of the patient. The best assessment of dissection is the onset of symptoms. Usually found either in the EMS report or history of present illness on the H&P, record the time from first onset of pain until the patient is evaluated for treatment. Report “unknown” ONLY if the patient cannot describe a specific onset of symptoms.

SEQ. #: 4746

Long Name: Dissection Onset Date Known

Short Name: DisOnsetDtKnown

Definition: Indicate whether the date of dissection onset is known

Intent/Clarification:

The intent is to confirm the duration of symptoms preceding the patient’s evaluation for treatment. While dissection timing (seq 4745) describes fairly broad intervals, this sequence refers to the patient’s recall of specific date when symptoms were first felt. Typical symptoms include sudden onset of pain which is usually memorable. Report “no” ONLY for any patient whose dissection is incidentally discovered or if the patient does not recall the onset of pain.

SEQ. #: 4747

Long Name: Dissection Onset Date **Short Name:** DisOnsetDt

Definition: Indicate dissection onset date

Intent/Clarification:

Report the date of symptoms onset if it is known by the patient. If the patient’s recall is non-specific (e.g. “Sometime last week”) leave this item blank. Use 8-digit format (mm/dd/yyyy).

SEQ. #: 4750

Long Name: Dissection - Primry Tear Location **Short Name:** DisTearLoc

Definition: Indicate location of the primary tear

Intent/Clarification:

The intent is to identify the primary entry tear for the dissection. As most dissections include multiple re-entry tears it may be difficult to confirm the primary site and the surgeon MUST be the final arbiter of this definition. This is the site identified by the surgeon at an open operation or judged by the surgeon from imaging as the primary site

to be covered by endovascular stent. If the radiology report names a primary entry point and the surgeon concurs, report this location.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

Zone "0" is subdivided into 3 sections:

- A. Below sinotubular junction
- B. Sinotubular junction to mid-ascending aorta
- C. Mid-ascending to distal ascending (at the innominate artery)

SEQ. #: 4755

Long Name: Dissection - Secondary Tear Location **Short Name:** DisSecLoc

Definition: Indicate location of secondary tear

Intent/Clarification:

The intent is to identify any secondary tear for the dissection. This would be a re-entry site resulting from flow within the false lumen returning to the true lumen. The surgeon MUST be the final arbiter of this definition. This is the site identified by the surgeon at open operation or judged by the surgeon from imaging as a secondary site to be covered by endovascular stent.

Refer to the image showing the zones and note that zone "0" is subdivided into 3 sections:

- A. Below sinotubular junction
- B. Sinotubular junction to mid-ascending aorta

C. Mid-ascending to distal ascending (at the innominate artery)

SEQ. #: 4760

Long Name: Dissection - Retrograde Extension **Short Name:** DisRetExt

Definition: Indicate whether there was retrograde extension

Intent/Clarification:

The intent is to determine whether the dissection propagates proximal (toward the aortic valve) from the primary tear location. Report yes if imaging indicates an extension of the false lumen proximal (toward the aortic valve) to the primary tear location.

SEQ. #: 4765

Long Name: Dissection - Retrograde Location

Short Name: DisRetLoc

Definition: Indicate location of retrograde extension

Intent/Clarification:

The intent is to define how far the retrograde dissection extends toward the aortic valve. This would be the point at which the false lumen comes closest to the aortic valve. The surgeon or radiologist can be the final arbiter of this definition. Refer to the image showing the zones and note that zone "0" is subdivided into 3 sections:

- A. Below sinotubular junction
- B. Sinotubular junction to mid-ascending aorta
- C. Mid-ascending to distal ascending (at the innominate artery)

- | |
|---|
| <ul style="list-style-type: none">A. Below sinotubular junctionB. Sinotubular junction to mid ascendingC. Mid ascending to distal ascendingD. Zone 1 (between innominate and left carotid)E. Zone 2 (between left carotid and left subclavian)F. Zone 3 (first 2 cm. distal to left subclavian)G. Zone 4 (end of zone 3 to mid descending aorta - T6)H. Zone 5 (mid descending aorta to celiac)I. Zone 6 (celiac to superior mesenteric)J. Zone 7 (superior mesenteric to renals)K. Zone 8 (renal to infra-renal abdominal aorta)L. Zone 9 (infrarenal abdominal aorta)M. Zone 10 (common iliac)N. Zone 11 (external iliacs) |
|---|

SEQ. #: 4770

Long Name: Dissection - Post TEVAR

Short Name: DisPosTEVAR

Definition: Indicate whether dissection occurred following TEVAR

Intent/Clarification:

The intent is to identify whether RETROGRADE dissection occurred or extended during TEVAR (Thoracic Endovascular Aortic Repair)

Report yes if:

- A. Retrograde dissection is noted on post TEVAR imaging that was not present on imaging before TEVAR
- OR
- B. Retrograde dissection (false lumen) extends closer to the aortic valve than was noted on pre TEVAR imaging

SEQ. #: 4775

Long Name: Dissection - Distal Extension

Short Name: DistalExt

Definition: Indicate whether there is distal extension

Intent/Clarification:

The intent is to identify where distal (antegrade) dissection occurred or extended.

SEQ. #: 4780

Long Name: Dissection - Distal Extension Location

Short Name: DistalExtLoc

Definition: Indicate location of distal extension

Intent/Clarification:

The intent is to define the how far along the aorta (away from the valve) any new or extended dissection goes. Refer to the image showing the zones and report the most distal (highest # zone) extent of the false lumen.

SEQ. #: 4785

Long Name: Dissection - Malperfusion

Short Name: DisMal

Definition: Indicate whether malperfusion was present

Intent/Clarification:

The intent is to identify whether there is compromised blood flow to any branch vessel as a consequence of the dissection or repair. Radiology report or the surgeon's evaluation may be used to define this. If any vessel has compromised blood flow report "yes". Report "unknown" if the surgeon or radiologist indicate that the imaging is inadequate to confirm the presence or absence of malperfusion.

SEQ. #: 4790

Long Name: Dissection - Malperfusion - Coronary

Short Name: DisMalCor

Definition: Indicate whether coronary malperfusion was present

Intent/Clarification:

The intent is to identify which vessels have compromised flow as a consequence of the dissection or repair. If the answer to Dissection – malperfusion (sequence 4785) is “yes” AND any coronary blood flow is compromised report yes. The surgeon is the final arbiter of this definition.

SEQ. #: 4791

Long Name: Dissection - Malperfusion - Right Subclavian

Short Name: DisMalRtSubclav

Definition: Indicate whether right subclavian malperfusion was present

Intent/Clarification:

The intent is to identify which vessels have compromised flow as a consequence of the dissection or repair. If the answer to Dissection – malperfusion (sequence 4785) is “yes” AND right subclavian blood flow is compromised report yes. The surgeon is the final arbiter of this definition.

SEQ. #: 4792

Long Name: Dissection - Malperfusion - Right Common Carotid

Short Name: DisMalRtComCar

Definition: Indicate whether right common carotid malperfusion was present

Intent/Clarification:

The intent is to identify which vessels have compromised flow as a consequence of the dissection or repair. If the answer to Dissection – malperfusion (sequence 4785) is “yes” AND right common carotid blood flow is compromised report yes. The surgeon is the final arbiter of this definition.

SEQ. #: 4800

Long Name: Dissection - Malperfusion - Left Common Carotid

Short Name: DisMalComL

Definition: Indicate whether left common carotid malperfusion was present

Intent/Clarification:

The intent is to identify which vessels have compromised flow as a consequence of the dissection or repair. If the answer to Dissection – malperfusion (sequence 4785) is “yes” AND left common carotid blood flow is compromised report yes. The surgeon is the final arbiter of this definition.

SEQ. #: 4805

Long Name: Dissection - Malperfusion - Left Subcalvian

Short Name: DisMalSubL

Definition: Indicate whether left subclavian malperfusion was present

Intent/Clarification:

The intent is to identify which vessels have compromised flow as a consequence of the dissection or repair. If the answer to Dissection – malperfusion (sequence 4785) is “yes” AND left subclavian blood flow is compromised report yes. The surgeon is the final arbiter of this definition.

SEQ. #: 4810

Long Name: Dissection - Malperfusion - Celiac

Short Name: DisMalCel

Definition: Indicate whether celiac malperfusion was present

Intent/Clarification:

The intent is to identify which vessels have compromised flow as a consequence of the dissection or repair. If the answer to Dissection – malperfusion (sequence 4785) is “yes” AND celiac blood flow is compromised report yes. The surgeon is the final arbiter of this definition.

SEQ. #: 4815

Long Name: Dissection - Malperfusion - Superior Mesenteric

Short Name: DisMalSup

Definition: Indicate whether superior mesenteric malperfusion was present

Intent/Clarification:

The intent is to identify which vessels have compromised flow as a consequence of the dissection or repair. If the answer to Dissection – malperfusion (sequence 4785) is “yes” AND superior mesenteric blood flow is compromised report yes. The surgeon is the final arbiter of this definition.

SEQ. #: 4820

Long Name: Dissection - Malperfusion - Renal, Left

Short Name: DisMalRenL

Definition: Indicate whether left renal malperfusion was present

Intent/Clarification:

The intent is to identify which vessels have compromised flow as a consequence of the dissection or repair. If the answer to Dissection – malperfusion (sequence 4785) is “yes” AND left renal blood flow is compromised report yes. The surgeon is the final arbiter of this definition.

SEQ. #: 4825

Long Name: Dissection - Malperfusion - Renal, Right

Short Name: DisMalRenR

Definition: Indicate whether right renal malperfusion was present

Intent/Clarification:

The intent is to identify which vessels have compromised flow as a consequence of the dissection or repair. If the answer to Dissection – malperfusion (sequence 4785) is “yes” AND left renal blood flow is compromised report yes. The surgeon is the final arbiter of this definition.

SEQ. #: 4830

Long Name: Dissection - Malperfusion - Iliofemoral

Short Name: DisMalIlio

Definition: Indicate whether iliofemoral malperfusion was present

Intent/Clarification:

The intent is to identify which vessels have compromised flow as a consequence of the dissection or repair. If the answer to Dissection – malperfusion (sequence 4785) is “yes” AND either or both iliofemoral systems blood flow is compromised report yes. The surgeon is the final arbiter of this definition.

SEQ. #: 4835

Long Name: Dissection - Malperfusion - Spinal

Short Name: DisMalSpin

Definition: Indicate whether spinal malperfusion was present

Intent/Clarification:

The intent is to identify which vessels have compromised flow as a consequence of the dissection or repair. If the answer to Dissection – malperfusion (sequence 4785) is “yes” AND any spinal artery blood flow is compromised report yes. The surgeon is the final arbiter of this definition.

SEQ. #: 4836

Long Name: Dissection - lower Extremity Motor Function **Short Name:** DisLowMotFun

Definition: Indicate status of lower extremity motor function

Intent/Clarification:

The intent is to identify if any NEW motor deficit of either lower extremity as a presenting symptom. This is preoperative status and does not include new post-operative paralysis or paraplegia.

This is intended to capture new sensory-motor deficit due to vascular malperfusion and not due to post-operative complication.

SEQ. #: 4837

Long Name: Dissection - Lower Extremity Sensory Deficit

Short Name: DisLowSenDef

Definition: Indicate whether lower extremity sensory deficit is present

Intent/Clarification:

The intent is to identify any NEW sensory deficit of either lower extremity is present following dissection. Report “yes” if any note comments on numbness or insensate areas that were not recorded in the past medical history. Only report “unknown” if there is no comment in the medical record regarding sensation in the lower extremities.

This is preoperative status and does not include post-operative paralysis or paraplegia.

SEQ. #: 4840

Long Name: Dissection - Rupture

Short Name: DisRupt

Definition: Indicate whether dissection ruptured

Intent/Clarification:

Report “yes” if any volume of blood is extravascular (outside the aortic adventitial layer), i.e. beyond the outmost layer of the aortic wall.

SEQ. #: 4845

Long Name: Dissection - Rupture - Contained

Short Name: DisRuptCon

Definition: Indicate whether the rupture was contained.

Intent/Clarification:

Report “yes” if extravascular blood is contained by surrounding structures such that bleeding has stopped.

SEQ. #: 4850

Long Name: Dissection - Rupture Location

Short Name: DisRuptLoc

Definition: Indicate the rupture location

Intent/Clarification:

Intent is to identify where the rupture occurred. This is the site identified by the surgeon at an open operation or judged by the surgeon or radiologist from imaging as the rupture site to be covered by endovascular stent. Refer to the image showing the zones and note that zone “0” is subdivided into 3 sections:

- A. Below sinotubular junction
- B. Sinotubular junction to mid-ascending aorta
- C. Mid-ascending to distal ascending (at the innominate artery)

- | |
|--|
| <ul style="list-style-type: none">A. Below sinotubular junctionB. Sinotubular junction to mid ascendingC. Mid ascending to distal ascendingD. Zone 1 (between innominate and left carotid)E. Zone 2 (between left carotid and left subclavian)F. Zone 3 (first 2 cm. distal to left subclavian)G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)H. Zone 5 (mid descending aorta to celiac)I. Zone 6 (celiac to superior mesenteric)J. Zone 7 (superior mesenteric to renals)K. Zone 8 (renal to infra-renal abdominal aorta)L. Zone 9 (infrarenal abdominal aorta)M. Zone 10 (common iliac)
N. Zone 11 (external iliacs) |
|--|

SEQ. #: 4855

Long Name: Root - Aorto-Annular Ectasia

Short Name: RootAAnnEctasia

Definition: Indicate whether aorto-annular ectasia is present

Intent/Clarification:

Annuloaortic ectasia refers to dilatation of the aortic root involving the annulus and/or the sinuses and/or the sinotubular junction and typically giving rise to aortic insufficiency. The intent of capturing this field is to identify patients with aortic root dilatation specifically that impacts aortic valvular function.

SEQ. #: 4870

Long Name: Root - Asymmetric Root Dilatation

Short Name: RootDilaAsym

Definition: Indicate whether asymmetric root dilatation is present

Intent/Clarification:

Asymmetric root dilatation refers to predominance of dilatation present in one or two sinus segments as opposed to more uniform root dilatation involving all 3 sinus segments (these may often be associated with aortic insufficiency).

The intent of this field is to determine the relative frequency of asymmetric sinus dilatation and its relationship to other clinical manifestations (e.g. aortic insufficiency or aortic dissection) as opposed to more uniform root dilatation.

SEQ. #: 4875

Long Name: Root - Asymmetric Root Dilatation - Location

Short Name: RootDilaAsym

Definition: Indicate location of asymmetric root dilatation

Intent/Clarification:

The intent is to clarify left, right, or non-coronary aortic root dilatation.

SEQ. #: 4880

Long Name: Root - Sinus Of Valsalva Aneurysm

Short Name: RootSinus

Definition: Indicate whether there is a sinus of Valsalva aneurysm

Intent/Clarification:

SOV aneurysm specifically refers to distinct dilatation of a single sinus segment, i.e. does not involve a second sinus segment as would be the case with “asymmetric root dilatation”.

The intent of this field is to identify the frequency of distinct sinus segment aneurysms as opposed to other root pathologies.

SEQ. #: 4881

Long Name: Root - Sinus Of Valsalva Aneurysm - Location

Short Name: RootSinusLoc

Definition: Indicate location of sinus of Valsalva aneurysm

Intent/Clarification: The intent is to clarify left, right, or non-coronary sinus of Valsalva aneurysm.

FAQ September 2017: An aortic dissection case had Sinus of Valsalva aneurysm within the right, left and non-coronary sinuses, it was symmetrical. Since you cannot answer all three, what should be answered?

Answer: Answer yes to sequence number 4880 and leave sequence number 4881 blank.

SEQ. #: 4882

Long Name: Arch Type

Short Name: ArchType

Definition: Indicate arch type

Intent/Clarification:

Right arch: a right arch implies that the aortic arch travels around (anteriorly) to the right mainstem bronchus and right pulmonary artery and then passes posterior to the trachea.

Left arch: a left arch implies that the aortic arch travels anterior to the trachea and then passes over and around the left pulmonary artery.

Grouped Intent/Clarification: The intent of these arch fields is to establish a concise characterization of the arch anatomy and its brachiocephalic branches at the time of arch interventions in order to allow for correlations of specific anatomy with other clinical findings/characterizations.

SEQ. #: 4884

Long Name: Arch - Aberrant Right Subclavian

Short Name: ArchAbRtSub

Definition: Indicate whether the right subclavian is aberrant

Intent/Clarification:

An aberrant right subclavian artery is any artery that does not emanate from the innominate artery (these are typically associated with left arch anatomy).

Grouped Intent/Clarification: The intent of these arch fields is to establish a concise characterization of the arch anatomy and its brachiocephalic branches at the time of arch interventions in order to allow for correlations of specific anatomy with other clinical findings/characterizations.

SEQ. #: 4885

Long Name: Arch - Aberrant Left Subclavian

Short Name: ArchAbLtSub

Definition: Indicate whether the left subclavian is aberrant

Intent/Clarification:

An aberrant left subclavian is any left subclavian that does not emanate from the distal arch as a separate ostium sequential and distal to the takeoff of the left common carotid artery on the greater curvature of the aortic arch.

Grouped Intent/Clarification: The intent of these arch fields is to establish a concise characterization of the arch anatomy and its brachiocephalic branches at the time of arch interventions in order to allow for correlations of specific anatomy with other clinical findings/characterizations.

SEQ. #: 4886

Long Name: Arch - Kommerell

Short Name: ArchKom

Definition: Indicate whether Kommerell arch type is present

Intent/Clarification:

Kommerell's diverticulum: This is not a true diverticulum but a remnant of the left fourth aortic arch and is a bulbous dilatation at the origin of the left subclavian artery. It is often associated with other arch anomalies.

Grouped Intent/Clarification: The intent of these arch fields is to establish a concise characterization of the arch anatomy and its brachiocephalic branches at the time of arch interventions in order to allow for correlations of specific anatomy with other clinical findings/characterizations.

SEQ. #: 4887

Long Name: Arch - Bovine

Short Name: ArchBovine

Definition: Indicate whether bovine arch type is present

Intent/Clarification:

This entity refers to a common origin of both the innominate artery and the left common carotid artery as they emanate from the greater curve of the arch.

Grouped Intent/Clarification: The intent of these arch fields is to establish a concise characterization of the arch anatomy and its brachiocephalic branches at the time of arch interventions in order to allow for correlations of specific anatomy with other clinical findings/characterizations.

SEQ. #: 4888

Long Name: Arch - Variant Vertebral Origin

Short Name: ArchVarVertOr

Definition: Indicate whether there is variant origin of the vertebral

Intent/Clarification:

This refers to any vertebral artery that emanates directly from the aortic arch rather than a branch of either subclavian artery.

Grouped Intent/Clarification: The intent of these arch fields is to establish a concise characterization of the arch anatomy and its brachiocephalic branches at the time of arch interventions in order to allow for correlations of specific anatomy with other clinical findings/characterizations.

SEQ. #: 4889

Long Name: Arch - Patent Internal Mammary Artery Bypass Graft

Short Name: ArchPatIMA

Definition: Indicate whether there is a patent internal mammary bypass graft present

Intent/Clarification:

Patent internal mammary artery bypass graft: this refers specifically to a patient who has undergone prior CABG and has a patent internal mammary graft (either left or right) present.

Grouped Intent/Clarification: The intent of these arch fields is to establish a concise characterization of the arch anatomy and its brachiocephalic branches at the time of arch interventions in order to allow for correlations of specific anatomy with other clinical findings/characterizations.

SEQ. #: 4891

Long Name: Ascending Asymmetric Dilatation

Short Name: AscAsymDil

Definition: Indicate whether there is asymmetric dilatation of the ascending aorta

Intent/Clarification:

Asymmetric dilatation refers to non-uniform dilatation of the aorta distal to the sinotubular junction, as is often noted as a pattern of dilation that affects the greater curvature of the ascending aorta.

SEQ. #: 4892

Long Name: Ascending Proximal Coronary Bypass Grafts

Short Name: AscProxGr

Definition: Indicate whether proximal bypass grafts are present on the aorta

Intent/Clarification:

These refer to any saphenous vein graft, radial artery or free internal mammary artery graft that emanates from the ascending aorta.

The intent is to tabulate the relative frequency of bypass grafts from the ascending aorta during reoperative cases that require aortic reconstruction.

SEQ. #: 4895

Long Name: 3-D Reconstruction Aortic Diameter Measurements Available

Short Name: Diameter3DMeas

Definition: Indicate whether 3-D reconstruction aortic diameter measurements are available

Intent/Clarification:

3-D reconstruction of CT imaging of the aorta, which accounts for the 3-dimensional curvature of the aorta, draws a cross section perpendicular to the centerline direction of blood flow and is more accurate than 2-D axial, coronal, or sagittal images, which may show a cross-section of the aorta diagonal to the centerline direction of blood flow. 3-D reconstruction is most accurate when the CT scan is obtained with ECG-gating, which accounts for cardiac pulsation and motion of the aortic root. The Radiology report should note whether 3-D reconstruction was performed. Alternately, CT images may be sent by the surgeon to an outside laboratory for 3-D reconstruction (e.g., M2S). When measuring the largest diameter of the aorta, this should be done from adventitia to adventitia (i.e., including the wall of the aorta, not just the area with contrast).

- | |
|--|
| <ul style="list-style-type: none">A. Below sinotubular junctionB. Sinotubular junction to mid ascendingC. Mid ascending to distal ascendingD. Zone 1 (between innominate and left carotid)E. Zone 2 (between left carotid and left subclavian)F. Zone 3 (first 2 cm. distal to left subclavian)G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)H. Zone 5 (mid descending aorta to celiac)I. Zone 6 (celiac to superior mesenteric)J. Zone 7 (superior mesenteric to renals)K. Zone 8 (renal to infra-renal abdominal aorta)L. Zone 9 (infrarenal abdominal aorta)M. Zone 10 (common iliac)
N. Zone 11 (external iliacs) |
|--|

FAQ October 2017: If 3D measurements are available in the PACs system, can the data manager use the values to code 3D measurements.

Answer: The data manager should not use the measurements from the PACs unless the measurements are supported by documentation from the surgeon. It is **not** the intention of the STS that the data manager should interpret the measurements.

SEQ. #: 4900

Long Name: Diameter Measurements 3D - Annulus

Short Name: Diam3DAnnulus

Definition: Indicate diameter of the annulus

Intent/Clarification:

The annulus is the region where the left ventricular outflow tract meets the aortic root and lies at the level of a plane defined by the lowest point of the aortic sinuses. It is usually oval-shaped with minimum and maximum diameters. Provide the average of the minimum and maximum diameters in millimeters.

SEQ. #: 4905

Long Name: Diameter Measurements 3D - Sinus Segment

Short Name: Diam3DSinus

Definition: Indicate diameter of the sinus segment

Intent/Clarification:

This is the widest diameter of the aortic root, usually at the mid-sinus level.

SEQ. #: 4910

Long Name: Diameter Measurements 3D - Sinotubular Junction

Short Name: Diam3DSinotubular

Definition: Indicate the diameter of the sinotubular junction

Intent/Clarification:

This is the region where the bulbous aortic root meets the tubular ascending aorta. Provide the largest diameter.

SEQ. #: 4915

Long Name: Diameter Measurements 3D - Mid-ascending

Short Name: Diam3DMidAsc

Definition: Indicate the diameter of the mid-ascending aorta

Intent/Clarification:

The mid ascending aorta is usually measured at the bifurcation of the pulmonary artery. Provide the largest diameter.

SEQ. #: 4920

Long Name: Diameter Measurements 3D - Distal Ascending

Short Name: Diam3DDistalAsc

Definition: Indicate the diameter of the distal ascending aorta

Intent/Clarification:

The distal ascending aorta is usually measured just proximal to the origin of the innominate artery. Provide the largest diameter.

SEQ. #: 4925

Long Name: Diameter Measurements 3D - Zone 1

Short Name: Diam3DZone1

Definition: Indicate the diameter of zone 1

Intent/Clarification:

Zone 1 of the aorta includes the segment of aorta between the innominate artery and left carotid artery as well as the segment of aorta from which the left carotid artery arises (see figure). Provide the largest diameter.

SEQ. #: 4930

Long Name: Diameter Measurements 3D - Zone 2

Short Name: Diam3DZone2

Definition: Indicate the diameter of zone 2

Intent/Clarification:

Zone 2 of the aorta includes the segment of aorta between the left carotid artery and left subclavian artery as well as the segment of aorta from which the left subclavian artery arises (see figure). Provide the largest diameter.

SEQ. #: 4935

Long Name: Diameter Measurements 3D - Zone 3

Short Name: Diam3DZone3

Definition: Indicate the diameter of zone 3

Intent/Clarification:

Zone 3 of the aorta is the 2 cm segment of aorta just beyond the left subclavian artery (see figure). Provide the largest diameter.

SEQ. #: 4940

Long Name: Diameter Measurements 3D - Zone 4

Short Name: Diam3DZone4

Definition: Indicate the diameter of zone 4

Intent/Clarification:

Zone 4 of the aorta extends from 2 cm beyond the left subclavian artery to the mid descending thoracic aorta, which is usually defined by the T6-T7 vertebral bodies (see figure). Provide the largest diameter.

SEQ. #: 4941

Long Name: Diameter Measurements 3D - Zone 5

Short Name: Diam3DZone5

Definition: Indicate the diameter of zone 5

Intent/Clarification:

Zone 5 of the aorta extends from the mid descending thoracic aorta (at T6-T7) to the origin of the celiac artery, but does not include the origin of the celiac artery (see figure). Provide the largest diameter.

SEQ. #: 4942

Long Name: Diameter Measurements 3D - Zone 6

Short Name: Diam3DZone6

Definition: Indicate the diameter of zone 6

Intent/Clarification:

Zone 6 of the aorta extends from the celiac artery to the origin of the superior mesenteric artery, but does not include the origin of the superior mesenteric artery (see figure). Provide the largest diameter.

SEQ. #: 4943

Long Name: Diameter Measurements 3D - Zone 7

Short Name: Diam3DZone7

Definition: Indicate the diameter of zone 7

Intent/Clarification:

Zone 7 of the aorta extends from the superior mesenteric artery to the origin of the first renal artery, but does not include the origin of the first renal artery (see figure). Provide the largest diameter.

SEQ. #: 4944

Long Name: Diameter Measurements 3D - Zone 8

Short Name: Diam3DZone8

Definition: Indicate the diameter of zone 8

Intent/Clarification:

Zone 8 of the aorta is the segment of aorta from which all the renal arteries arise (usually two, but may be more) (see figure). Provide the largest diameter.

SEQ. #: 4945

Long Name: Diameter Measurements 3D - Zone 9

Short Name: Diam3DZone9

Definition: Indicate the diameter of zone 9

Intent/Clarification:

Zone 9 of the aorta is the segment of aorta between the last renal artery take-off and the aortic bifurcation (see figure). Provide the largest diameter.

SEQ. #: 4946

Long Name: Diameter Measurements 3D - Zone 10

Short Name: Diam3DZone10

Definition: Indicate the diameter of zone 10

Intent/Clarification:

Zone 10 is the common iliac arteries (see figure). Provide the largest diameter.

SEQ. #: 4947

Long Name: Diameter Measurements 3D - Zone 11

Short Name: Diam3DZone11

Definition: Indicate the diameter of zone 11

Intent/Clarification:

Zone 11 is the external iliac arteries (see figure). Provide the largest diameter.

SEQ. #: 4948

Long Name: Diameter Measurements Largest - Annulus

Short Name: DiamLgstAnnulus

Definition: Indicate diameter of the annulus.

Intent/Clarification:

The annulus is the region where the left ventricular outflow tract meets the aortic root and is a plane defined by the nadir of the aortic sinuses. It may be best seen on a

coronal or sagittal image. Provide the largest measurement. Use 3D measurements if available.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

FAQ October 2017: If the diameter measurement is not included in the surgeon dictation or the available 2-D or 3-D studies should the values be 0 or should the field be left blank.

Answer: Leave blank.

SEQ. #: 4949

Long Name: Diameter Measurements Largest - Sinus Segment

Short Name: DiamLgstSinus

Definition: Indicate diameter of the sinus segment

Intent/Clarification:

This is the widest diameter of the aortic root at the mid-sinus level. This may be best seen on a coronal or sagittal image.

FAQ October 2017: If the diameter measurement is not included in the surgeon dictation or the available 2-D or 3-D studies should the values be 0 or should the field be left blank.

Answer: Leave blank.

SEQ. #: 4950

Long Name: Diameter Measurements Largest - Sinotubular Junction

Short Name: DiamLgstSinotubular

Definition: Indicate the diameter of the sinotubular junction

Intent/Clarification:

This is the region where the bulbous aortic root meets the tubular ascending aorta. This may be best seen on a coronal image.

FAQ October 2017: If the diameter measurement is not included in the surgeon dictation or the available 2-D or 3-D studies should the values be 0 or should the field be left blank.

Answer: Leave blank.

SEQ. #: 4951

Long Name: Diameter Measurements Largest - Mid-ascending

Short Name: DiamLgstMidAsc

Definition: Indicate the diameter of the mid-ascending aorta

Intent/Clarification:

The mid ascending aorta is usually measured at the bifurcation of the pulmonary artery. This may be best seen on an axial image with the fullest view of the right pulmonary artery or on a coronal image with the largest view of the mid ascending aorta. On the axial image, the largest diameter should be recorded. On the coronal image, the measurement should be taken perpendicular to the direction of blood flow.

FAQ October 2017: If the diameter measurement is not included in the surgeon dictation or the available 2-D or 3-D studies should the values be 0 or should the field be left blank.

Answer: Leave blank.

SEQ. #: 4952

Long Name: Diameter Measurements Largest - Distal Ascending

Short Name: DiamLgstDistalAsc

Definition: Indicate the diameter of the distal ascending aorta

Intent/Clarification:

The distal ascending aorta is usually measured just proximal to the origin of the innominate artery. This may be best seen on a coronal image with the largest view of the aorta. The measurement should be taken perpendicular to the direction of blood flow.

FAQ October 2017: If the diameter measurement is not included in the surgeon dictation or the available 2-D or 3-D studies should the values be 0 or should the field be left blank.

Answer: Leave blank.

SEQ. #: 4953

Long Name: Diameter Measurements Largest - Zone 1

Short Name: DiamLgstZone1

Definition: Indicate the diameter of zone 1

Intent/Clarification:

Zone 1 of the aorta includes the segment of aorta between the innominate artery and left carotid artery as well as the segment of aorta from which the left carotid artery arises (see figure). This may be best seen on a sagittal image. The measurement should be taken perpendicular to the direction of blood flow.

FAQ October 2017: If the diameter measurement is not included in the surgeon dictation or the available 2-D or 3-D studies should the values be 0 or should the field be left blank.

Answer: Leave blank.

SEQ. #: 4954

Long Name: Diameter Measurements Largest - Zone 2

Short Name: DiamLgstZone2

Definition: Indicate the diameter of zone 2

Intent/Clarification:

Zone 2 of the aorta includes the segment of aorta between the left carotid artery and left subclavian artery as well as the segment of aorta from which the left subclavian artery arises (see figure). This may be best seen on a sagittal image. The measurement should be taken perpendicular to the direction of blood flow.

FAQ October 2017: If the diameter measurement is not included in the surgeon dictation or the available 2-D or 3-D studies should the values be 0 or should the field be left blank.

Answer: Leave blank.

SEQ. #: 4955

Long Name: Diameter Measurements Largest - Zone 3

Short Name: DiamLgstZone3

Definition: Indicate the diameter of zone 3

Intent/Clarification:

Zone 3 of the aorta is the 2 cm segment of aorta just beyond the left subclavian artery (see figure). This may be best seen on a sagittal image. The measurement should be taken perpendicular to the direction of blood flow.

FAQ October 2017: If the diameter measurement is not included in the surgeon dictation or the available 2-D or 3-D studies should the values be 0 or should the field be left blank.

Answer: Leave blank.

SEQ. #: 4956

Long Name: Diameter Measurements Largest - Zone 4

Short Name: DiamLgstZone4

Definition: Indicate the diameter of zone 4

Intent/Clarification:

Zone 4 of the aorta extends from 2 cm beyond the left subclavian artery to the mid descending thoracic aorta, which is usually defined by the T6-T7 vertebral bodies (see figure). Provide the largest diameter.

FAQ October 2017: If the diameter measurement is not included in the surgeon dictation or the available 2-D or 3-D studies should the values be 0 or should the field be left blank.

Answer: Leave blank.

SEQ. #: 4957

Long Name: Diameter Measurements Largest - Zone 5

Short Name: DiamLgstZone5

Definition: Indicate the diameter of zone 5

Intent/Clarification:

Zone 5 of the aorta extends from the mid descending thoracic aorta (at T6-T7) to the origin of the celiac artery, but does not include the origin of the celiac artery (see figure). Provide the largest diameter.

FAQ October 2017: If the diameter measurement is not included in the surgeon dictation or the available 2-D or 3-D studies should the values be 0 or should the field be left blank.

Answer: Leave blank.

SEQ. #: 4958

Long Name: Diameter Measurements Largest - Zone 6

Short Name: DiamLgstZone6

Definition: Indicate the diameter of zone 6

Intent/Clarification:

Zone 6 of the aorta extends from the celiac artery to the origin of the superior mesenteric artery, but does not include the origin of the superior mesenteric artery (see figure). Provide the largest diameter.

FAQ October 2017: If the diameter measurement is not included in the surgeon dictation or the available 2-D or 3-D studies should the values be 0 or should the field

be left blank.

Answer: Leave blank.

SEQ. #: 4959

Long Name: Diameter Measurements Largest - Zone 7

Short Name: DiamLgstZone7

Definition: Indicate the diameter of zone 7

Intent/Clarification:

Zone 7 of the aorta extends from the superior mesenteric artery to the origin of the first renal artery, but does not include the origin of the first renal artery (see figure). Provide the largest diameter.

FAQ October 2017: If the diameter measurement is not included in the surgeon dictation or the available 2-D or 3-D studies should the values be 0 or should the field be left blank.

Answer: Leave blank.

SEQ. #: 4960

Long Name: Diameter Measurements Largest - Zone 8

Short Name: DiamLgstZone8

Definition: Indicate the diameter of zone 8

Intent/Clarification:

Zone 8 of the aorta is the segment of aorta from which all the renal arteries arise (usually two, but may be more) (see figure). Provide the largest diameter.

FAQ October 2017: If the diameter measurement is not included in the surgeon dictation or the available 2-D or 3-D studies should the values be 0 or should the field be left blank.

Answer: Leave blank.

SEQ. #: 4961

Long Name: Diameter Measurements Largest - Zone 9

Short Name: DiamLgstZone9

Definition: Indicate the diameter of zone 9

Intent/Clarification:

Zone 9 of the aorta is the segment of aorta between the last renal artery take-off and the aortic bifurcation (see figure). Provide the largest diameter.

FAQ October 2017: If the diameter measurement is not included in the surgeon dictation or the available 2-D or 3-D studies should the values be 0 or should the field be left blank.

Answer: Leave blank.

SEQ. #: 4962

Long Name: Diameter Measurements Largest - Zone 10

Short Name: DiamLgstZone10

Definition: Indicate the diameter of zone 10

Intent/Clarification:

Zone 10 is the common iliac arteries (see figure). Provide the largest diameter.

FAQ October 2017: If the diameter measurement is not included in the surgeon dictation or the available 2-D or 3-D studies should the values be 0 or should the field be left blank.

Answer: Leave blank.

SEQ. #: 4963

Long Name: Diameter Measurements Largest - Zone 11

Short Name: DiamLgstZone11

Definition: Indicate the diameter of zone 11

Intent/Clarification:

Zone 11 is the external iliac arteries (see figure). Provide the largest diameter.

FAQ October 2017: If the diameter measurement is not included in the surgeon dictation or the available 2-D or 3-D studies should the values be 0 or should the field be left blank.

Answer: Leave blank.

SEQ. #: 4970

Long Name: Planned Staged Hybrid

Short Name: PlanStagHybrid

Definition: Indicate whether the procedure was a planned staged hybrid

Intent/Clarification:

The intent is to identify procedures that will involve a combination of open and endovascular procedures or devices. In particular, the combination of an open approach with stent grafts which can be deployed open on endovascularly. Staged procedure

means that this will be done in more than one setting. For instance, two trips to the operating room or hybrid room.

FAQ October 2017: The patient was initially scheduled for TEVAR but after a series of CT scans and manifestations of signs and symptoms, the surgeon opted to repair the ascending aortic flap first and then, 3 days later, he took the patient back to the OR for the TEVAR. Should this be captured as post-operative event aortic reintervention?

Answer: No, code as a planned staged hybrid. Then capture the TEVAR accordingly in Seq. No. 5095.

SEQ. #: 4975

Long Name: Open Arch Procedure

Short Name: ArchProc

Definition: Indicate whether there was an open arch procedure

Intent/Clarification:

The intent is to identify procedures with replacement of or connection to the arch of the aorta. Anything from the base of the innominate through the subclavian takeoff would be included.

SEQ. #: 4980

Long Name: Open Arch Procedure - Distal Technique

Short Name: ArchDisTech

Definition: Indicate the distal technique for the arch procedure

Intent/Clarification:

The intent is to define that the distal anastomosis was done with or without a clamp. Many arch procedures are done with the clamp removed, sewing to the aorta looking down the barrel of the vessel. This of course requires circulatory arrest. The clamp means that the aorta is clamped with an instrument and the anastomosis is completed proximal (close to the heart) to that part of the aorta.

SEQ. #: 4985

Long Name: Open Arch Procedure - Distal Site

Short Name: ArchDiscSite

Definition: Indicate the distal site

Intent/Clarification:

The intent of this is to define the level of the distal (far from the heart) anastomosis. Ascending aorta implies the ascending was resected with a clamp on the distal

ascending aorta. Hemiarch means a single anastomosis was done somewhere in the ascending or proximal arch without separate grafts to the head vessels. Zone 1 means the innominate was reconnected with a graft between the innominate and left common carotid takeoffs. Zone 2 means the innominate and carotid were reconnected with a graft sewn to between the left common carotid and the left subclavian takeoffs. Zone three means the innominate, carotid and the left subclavian were reconnected with the graft being sewn beyond the left subclavian takeoff. Zone 4 means the graft was sewn to the mid descending thoracic aorta.

SEQ. #: 4990

Long Name: Open Arch Procedure - Distal Extention

Short Name: ArchDisExt

Definition: Indicate distal extension type

Intent/Clarification:

The intent of the question is to define whether graft was left that extended (distally) beyond the arch anastomosis. An elephant trunk is a soft graft, while a frozen elephant trunk means a stent was placed distally.

SEQ. #: 4995

Long Name: Open Arch Procedure - Arch Branch Reimplantation

Short Name: ArchBranReimp

Definition: Indicate whether arch branch reimplantation was performed

Intent/Clarification:

The intent of this is to define the end branches that were sewn to the graft.

SEQ. #: 5000

Long Name: Open Arch Procedure - Arch Branch Reimplantation - Innominate

Short Name: ArchBranInnom

Definition: Indicate whether arch branch reimplantation included the innominate artery

Intent/Clarification:

The intent is to determine whether the innominate artery was reattached to the graft.

SEQ. #: 5001

Long Name: Open Arch Procedure - Arch Branch Reimplantation - Right Subclavian

Short Name: ArchBranRSub

Definition: Indicate whether arch branch reimplantation included the right subclavian artery

Intent/Clarification:

The intent is to determine whether the right subclavian artery was reattached to the graft. This means the right subclavian was sewn to directly, not from the trunk or bifurcation of the innominate.

SEQ. #: 5002

Long Name: Open Arch Procedure - Arch Branch Reimplantation - Right Common Carotid

Short Name: ArchBranRComm

Definition: Indicate whether arch branch reimplantation included the right common carotid artery

Intent/Clarification:

The intent is to determine whether the right carotid artery was reattached to the graft. This means the right subclavian was sewn to directly, not from the trunk or bifurcation of the innominate.

SEQ. #: 5005

Long Name: Open Arch Procedure - Arch Branch Reimplantation - Left Common Carotid

Short Name: ArchBranLComm

Definition: Indicate whether arch branch reimplantation included the left common carotid artery

Intent/Clarification:

The intent is to determine whether the left common carotid artery was reattached to the graft.

SEQ. #: 5010

Long Name: Open Arch Procedure - Arch Branch Reimplantation - Left Subclavian

Short Name: ArchBranLSub

Definition: Indicate whether arch branch reimplantation included the left subclavian artery

Intent/Clarification:

The intent is to determine whether the left subclavian artery was reattached to the graft.

SEQ. #: 5011

Long Name: Open Arch Procedure - Arch Branch Reimplantation - Left Vertebral **Short Name:** ArchBranLVert

Definition: Indicate whether arch branch reimplantation included the left vertebral artery

Intent/Clarification:

The intent is to determine whether the left vertebral artery was reattached to the graft. This means a separate graft or anastomosis was created for the vertebral, not when it remains attached to the left subclavian artery.

SEQ. #: 5012

Long Name: Open Arch Procedure - Arch Branch Reimplantation - Other

Short Name: ArchBranOth

Definition: Indicate whether arch branch reimplantation included any other artery

Intent/Clarification:

The intent is to determine whether the left vertebral artery was reattached to the graft. This means a separate graft or anastomosis was created for the vertebral, not when it remains attached to the left subclavian artery.

SEQ. #: 5015

Long Name: Open Descending Thoracic Aorta or Thoracoabdominal Procedure

Short Name: DescAortaProc

Definition: Indicate whether there was an open procedure of the descending thoracic or thoracoabdominal aorta

Intent/Clarification:

The intent of this is to define procedures involving the descending thoracic aorta or the thoraco-abdominal aorta, usually through the left chest.

SEQ. #: 5020

Long Name: Proximal Location

Short Name: DescAortaLoc

Definition: Indicate the proximal location of the descending aorta procedure

Intent/Clarification:

The intent of this procedure is to define the proximal extent or coverage of the arch as defined by the zones defined on the collection form or with an open anastomosis to the mid to distal arch, without branch anastomosis, known as a hemiarch. Zones imply the zone branches are taken or revascularized.

SEQ. #: 5030

Long Name: Intercostal Reimplantation

Short Name: AortaInterReimp

Definition: Indicate whether intercostal vessels were reimplanted

Intent/Clarification:

The intent of this is to define procedures where either an island of intracostals is sewn to the graft or a separate branch is used to sew them to the graft.

SEQ. #: 5035

Long Name: Distal Location

Short Name: AortaDisZone

Definition: Indicate the distal location of the descending/thoracoabdominal procedure

Intent/Clarification:

The intent of this is to define the distal extant of the aortic intervention as defined by the zones defined on the collection form.

SEQ. #: 5045

Long Name: Visceral Vessel Intervention

Short Name: AortaVisceral

Definition: Indicate whether there was visceral vessel intervention

Intent/Clarification:

The intent of this is to define whether the celiac artery was revascularized by sewing it to the graft, reimplantation, or using a branch graft.

SEQ. #: 5050

Long Name: Visceral Vessel Intervention - Celiac

Short Name: AortaViscCel

Definition: Indicate whether the visceral vessel intervention involved the celiac artery

Intent/Clarification:

The intent of this is to define whether the celiac artery was revascularized by sewing it to the graft, reimplantation, or using a branch graft.

SEQ. #: 5055

Long Name: Visceral Vessel Intervention - Superior Mesenteric

Short Name: AortaViscSup

Definition: Indicate whether the visceral vessel intervention involved the superior mesenteric artery

Intent/Clarification:

The intent of this is to define whether the superior mesenteric artery was revascularized by sewing it to the graft, reimplantation, or using a branch graft.

SEQ. #: 5060

Long Name: Visceral Vessel Intervention - Right Renal

Short Name: AortaViscRenR

Definition: Indicate whether the visceral vessel intervention involved the right renal artery

Intent/Clarification:

The intent of this is to define whether the right renal artery was revascularized by sewing it to the graft, reimplantation, or using a branch graft.

SEQ. #: 5065

Long Name: Visceral Vessel Intervention - Left Renal

Short Name: AortaViscRenL

Definition: Indicate whether the visceral vessel intervention involved the left renal artery

Intent/Clarification:

The intent of this is to define whether the left renal artery was revascularized by sewing it to the graft, reimplantation, or using a branch graft.

SEQ. #: 5066

Long Name: Endovascular Procedures

Short Name: EndovasProc

Definition: Indicate whether there was an endovascular procedure

Intent/Clarification:

The intent is to capture catheter based procedures where a stent is implanted into the aorta.

SEQ. #: 5067

Long Name: Endovascular Procedures - Access

Short Name: EndovasAccess

Definition: Indicate the access used for the endovascular procedure

Intent/Clarification:

Please label the blood vessel through which the stent graft was delivered.

SEQ. #: 5068

Long Name: Endovascular Procedures - Percutaneous Access

Short Name: EndovasPercAcc

Definition: Indicate whether access was percutaneous

Intent/Clarification:

The intent is to capture needle access; no incision is required; a stab wound may be required for sheath placement.

SEQ. #: 5070

Long Name: Endovascular Procedures - Proximal Landing Zone

Short Name: EndoProxZone

Definition: Indicate the proximal landing zone

Intent/Clarification:

The proximal landing zone is the area of the area of the aorta closest to the heart where the graft is located. If two or more stent grafts were used, please label the proximal landing zone according to the site where the most proximal stent graft has its most proximal location.

- | |
|--|
| <ul style="list-style-type: none">A. Below sinotubular junctionB. Sinotubular junction to mid ascendingC. Mid ascending to distal ascendingD. Zone 1 (between innominate and left carotid)E. Zone 2 (between left carotid and left subclavian)F. Zone 3 (first 2 cm. distal to left subclavian)G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)H. Zone 5 (mid descending aorta to celiac)I. Zone 6 (celiac to superior mesenteric)J. Zone 7 (superior mesenteric to renals)K. Zone 8 (renal to infra-renal abdominal aorta)L. Zone 9 (infrarenal abdominal aorta)M. Zone 10 (common iliac)
N. Zone 11 (external iliacs) |
|--|

SEQ. #: 5080

Long Name: Endovascular Procedures - Distal Landing Zone

Short Name: EndoDistalZone

Definition: Indicate the distal landing zone

Intent/Clarification:

The distal landing zone defines the closet to the iliac bifurcation (furthest from the heart). If two or more stent grafts were used, please label the distal landing zone according to the site where the most distal stent graft has its most distal location.

SEQ. #: 5090

Long Name: Endovascular Procedures - TAVR

Short Name: EndovasTAVR

Definition: Indicate whether there was a transcatheter aortic valve procedure component

Intent/Clarification:

Indicate whether TAVR was performed in conjunction with this endovascular procedure.

SEQ. #: 5095

Long Name: Endovascular Procedures - Ascending TEVAR

Short Name: EndovasTEVAR

Definition: Indicate whether an ascending TEVAR was performed

Intent/Clarification:

Intent is to identify whether a stent graft placed in zone 0, a region spanning from the STJ to the innominate artery.

FAQ October 2017: The patient was initially scheduled for TEVAR but after a series of CT scans and manifestations of signs and symptoms, the surgeon opted to repair the ascending aortic flap first and then, 3 days later, he took the patient back to the OR for the TEVAR. Should this be captured as post operative event aortic reintervention?

Answer: No, code as a planned staged hybrid. Then capture the TEVAR accordingly in Seq. No. 5095.

SEQ. #: 5100

Long Name: Arch Vessel Management - Innominate

Short Name: Innominate

Definition: Indicate the management of the innominate artery

Intent/Clarification:

The innominate artery originates in the aortic arch as the first branch of the arch and divides into the right common carotid and right subclavian arteries.

Intent is to understand how the innominate artery received its blood flow following an endovascular procedure. Options included **native flow** where no direct endo-intervention on the vessel was performed or **extra-anatomic bypasses** which may be performed in a staged fashion (i.e. commonly an additional operation previous to the endovascular one, typically done during the same admission).

SEQ. #: 5105

Long Name: Innominate - Extra-Anatomic Bypass - Aorta-Innominate

Short Name: InAortaInnom

Definition: Indicate whether the extra-anatomic bypass was an aorta to innominate bypass

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy

Intent is to describe one option of how the arch branch vessel was managed as part of a hybrid strategy (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission) to provide blood flow following or as part of that strategy.

Aorta to innominate bypass means a graft was created from the native aorta or surgically replaced aorta to the innominate artery.

SEQ. #: 5110

Long Name: Innominate - Extra-Anatomic Bypass - Aorta-Right Carotid

Short Name: InAortaCarotid

Definition: Indicate whether the extra-anatomic bypass was an aorta to right carotid bypass

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

Intent is to describe specifically how the arch branch vessel was managed as part of a hybrid strategy (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission) to provide blood flow following or as part of that strategy.

Aorta to Right carotid bypass means a graft was created from the native aorta or surgically replaced aorta to the right carotid artery. This bypass is done beyond the innominate and often for aneurysm of the innominate and includes bypass of the right subclavian as well.

SEQ. #: 5115

Long Name: Innominate - Extra-Anatomic Bypass - Aorta-Right Subclavian

Short Name: InAortaSubclav

Definition: Indicate whether the extra-anatomic bypass was an aorta to right subclavian bypass

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

Intent is to describe specifically how the arch branch vessel was managed as part of a hybrid strategy (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission) to provide blood flow following or as part of that strategy.

Aorta to Right subclavian bypass means a graft was created from the native aorta or surgically replaced aorta to the right subclavian artery. This bypass is done beyond the innominate and often for aneurysm of the innominate and includes bypass of the right carotid as well.

SEQ. #: 5125

Long Name: Innominate - Extra-Anatomic Bypass - Right Carotid - Right Subclavian

Short Name: InCaroSubclav

Definition: Indicate whether the extra-anatomic bypass was a right carotid to right subclavian bypass

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

Intent is to describe specifically how the arch branch vessel was managed as part of a hybrid strategy (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission).

This means that either a bypass (i.e. use of a separate graft) or direct transposition was performed to create a communication between the right carotid and subclavian vessels.

SEQ. #: 5135

Long Name: Innominate - Extra-Anatomic Bypass - Other

Short Name: InOther

Definition: Indicate whether any other extra-anatomic innominate bypass was performed

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

Intent is to describe specifically how the arch branch vessel was managed as part of a hybrid strategy (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission).

SEQ. #: 5140

Long Name: Arch Vessel Management - Left Carotid

Short Name: LeftCarotid

Definition: Indicate the management of the left carotid artery

Intent/Clarification:

Intent is to describe specifically how the arch branch vessel was managed as part of a hybrid strategy (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission). Options included native flow where no direct endo-intervention on the vessel was performed or extra-anatomic bypasses which may be performed in a staged fashion (i.e. commonly an additional operation previous to the endovascular one).

The left carotid artery arises from the aortic arch.

SEQ. #: 5150

Long Name: Left Carotid - Extra-Anatomic Bypass - Aorta-Left Carotid

Short Name: LTCaroAortaCaro

Definition: Indicate whether the extra-anatomic bypass was an aorta to left carotid bypass

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

Intent is to describe specifically how the arch branch vessel was managed as part of a hybrid strategy (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission).

SEQ. #: 5160

Long Name: Left Carotid - Extra-Anatomic Bypass - Innominate-Left Carotid

Short Name: LTCarolInnomCaro

Definition: Indicate whether the extra-anatomic bypass was an innominate to left carotid bypass

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy. Intent is to describe specifically how the arch branch vessel was managed as part of a hybrid strategy (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission).

SEQ. #: 5170

Long Name: Left Carotid - Extra-Anatomic Bypass - Right Carotid - Left Carotid

Short Name: LTCaroCarotid

Definition: Indicate whether the extra-anatomic bypass was a right carotid to left carotid bypass

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

Intent is to describe specifically how the arch branch vessel was managed as part of a (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission). This bypass crossed the midline and may go either in front of or behind the esophagus.

SEQ. #: 5175

Long Name: Left Carotid - Extra-Anatomic Bypass - Other

Short Name: LTCaroOther

Definition: Indicate whether any other extra-anatomic left carotid bypass was performed

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

Intent is to describe specifically how the arch branch vessel was managed as part of a (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission).

SEQ. #: 5180

Long Name: Arch Vessel Management - Left Subclavian

Short Name: LeftSubclavian

Definition: Indicate the management of the left subclavian artery

Intent/Clarification:

Intent is to describe specifically how the arch branch vessel was managed as part of a hybrid strategy (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission). Options included native flow where no direct endo-intervention on the vessel was performed or extra-anatomic bypasses which may be performed in a staged fashion (i.e. commonly an additional operation previous to the endovascular one).

The left subclavian artery arises from the distal aortic arch.

SEQ. #: 5195

Long Name: Left Subclavian - Extra-Anatomic Bypass - Aorta-Left Subclavian

Short Name: LTSubAortaSub

Definition: Indicate whether the extra-anatomic bypass was an aorta to left subclavian bypass

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

Intent is to describe specifically how the arch branch vessel was managed as part of a hybrid strategy (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission).

SEQ. #: 5205

Long Name: Left Subclavian - Extra-Anatomic Bypass - Left Carotid-Left Subclavian

Short Name: LTSubCarotidSub

Definition: Indicate whether the extra-anatomic bypass was a left carotid to left subclavian bypass

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

Intent is to describe specifically how the arch branch vessel was managed as part of a hybrid strategy (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission).

This means that either a bypass (i.e. use of a separate graft) or direct transposition was performed to create a communication between the left carotid and subclavian vessels.

SEQ. #: 5213

Long Name: Left Subclavian - Extra-Anatomic Bypass - Other

Short Name: LTSubOther

Definition: Indicate whether any other extra-anatomic left subclavian bypass was performed

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

Intent is to describe specifically how the arch branch vessel was managed as part of a hybrid strategy (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission). Although most strategies are included in the list above this may be checked if another procedure is performed such as a transposition.

SEQ. #: 5214

Long Name: Arch Vessel Management - Other Arch Vessels Extra-Anatomic Bypass

Short Name: OthArchVes

Definition: Indicate whether other arch vessel extra-anatomic bypass was performed

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

Intent is to describe specifically how the arch branch vessel was managed as part of a hybrid strategy (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission) when something other than the vessels described above are included in the description.

SEQ. #: 5215

Long Name: Other - Extra-Anatomic Bypass - Innominate - Carotid

Short Name: OthInnomCaro

Definition: Indicate whether the extra-anatomic bypass was innominate to carotid

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

Intent is to describe specifically how the arch branch vessel was managed as part of a hybrid strategy (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission) when a bypass was created from the innominate to one of the carotid arteries.

This field may be redundant if the bypass was to the left carotid because it is described in a field above, but may checked if an innominate to right carotid bypass was performed but that would be a very rare event.

SEQ. #: 5216**Long Name:** Other - Extra-Anatomic Bypass - Innominate - Subclavian**Short Name:** OthInnomSub**Definition:** Indicate whether the extra-anatomic bypass was innominate to subclavian**Intent/Clarification:**

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

Intent is to describe specifically how the arch branch vessel was managed as part of a hybrid strategy (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission) when a bypass was created from the innominate artery to one of the subclavian arteries.

SEQ. #: 5217**Long Name:** Other - Extra-Anatomic Bypass - Subclavian - Subclavian**Short Name:** OthSubSub**Definition:** Indicate whether the extra-anatomic bypass was subclavian to subclavian**Intent/Clarification:**

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

Intent is to describe specifically how the arch branch vessel was managed as part of a hybrid strategy (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission) when a bypass was created between the two subclavian arteries.

SEQ. #: 5218

Long Name: Other - Extra-Anatomic Bypass - Other

Short Name: OthOther

Definition: Indicate whether any other extra-anatomic arch vessel bypass was performed

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

Intent is to describe specifically how the arch branch vessel was managed as part of a hybrid strategy (i.e. separate surgical procedure either done concurrently or at a previous time, typically done during the same admission). Many patients have their left vertebral artery originating from the aorta directly and this could be bypassed or reconstructed by transposition to the carotid artery or other means.

SEQ. #: 5220

Long Name: Visceral Vessel Management - Celiac

Short Name: Celiac

Definition: Indicate management of the celiac artery

Intent/Clarification:

The intent is to clarify whether the celiac axis/artery was revascularized during an endovascular repair of the thoracic or thoracoabdominal aorta.

If the celiac artery was left intact and continues to receive blood flow from the native aorta in the normal manner after the endovascular procedure then “**Native Flow**” should be selected.

If the celiac artery is instrumented with an endovascular branch graft extending from the main body of an aortic endograft and landing distally within the celiac as would be performed in an endovascular thoracoabdominal aortic repair then “**Endovascular Branch Graft**” should be selected.

If the celiac artery is instrumented with a separate endovascular graft that does not extend from an aortic endograft but rather courses parallel to an aortic endograft and lands distally within the celiac with flow through the celiac endovascular graft from the native aorta as would be performed with a “chimney” or “periscope” technique for endovascular thoracoabdominal aortic repair then “**Endovascular Parallel Graft**” should be selected.

If the celiac artery is fed by a surgical bypass graft (e.g. iliac artery to celiac bypass, infrarenal aorta to celiac bypass – see below) with subsequent endovascular coverage

of the celiac by the aortic endograft typically as would be performed in a “hybrid” thoracoabdominal aortic repair then “**Extra-anatomic Bypass**” should be selected.

If the celiac artery is covered by an aortic endograft with a fenestration/opening in the endograft corresponding to the location of the celiac and allowing continued antegrade flow into the celiac via this fenestration despite endograft coverage as would be performed in an endovascular juxtarenal or thoracoabdominal aortic repair then “**Fenestrated**” should be selected.

This may include coverage of a graft.

SEQ. #: 5225

Long Name: Celiac - Extra-Anatomic Bypass - Aorta-Celiac

Short Name: CeliacAortaCeli

Definition: Indicate whether the extra-anatomic bypass was aorta to celiac

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

The bypass may originate from any segment of the aorta (e.g. ascending, descending, abdominal).

SEQ. #: 5245

Long Name: Celiac - Extra-Anatomic Bypass - Iliac-Celiac

Short Name: CeliacIliacCeliac

Definition: Indicate whether the extra-anatomic bypass was iliac to celiac

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

The bypass may originate from any segment of the iliac artery (e.g. common, external, or internal).

SEQ. #: 5265

Long Name: Celiac - Extra-Anatomic Bypass - Other

Short Name: CeliacOther

Definition: Indicate whether another extra-anatomic celiac bypass was performed

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

This includes a bypass from any other vessel to the celiac or one of its branches (e.g. hepatic, splenic). Examples would include hepatorenal or splenorenal bypass.

SEQ. #: 5270

Long Name: Visceral Vessel Management - Superior Mesenteric

Short Name: SupMesenteric

Definition: Indicate management of the superior mesenteric artery

Intent/Clarification:

The intent is to clarify whether the superior mesenteric artery (SMA) was revascularized during an endovascular repair of the thoracic or thoracoabdominal aorta.

If the SMA was left intact and continues to receive blood flow from the native aorta in the normal manner after the endovascular procedure then “**Native Flow**” should be selected.

If the SMA is instrumented with an endovascular branch graft extending from the main body of an aortic endograft and landing distally within the SMA as would be performed in an endovascular thoracoabdominal aortic repair then “**Endovascular Branch Graft**” should be selected.

If the SMA is instrumented with a separate endovascular graft that does not extend from an aortic endograft but rather courses parallel to an aortic endograft and lands distally within the SMA with flow through the SMA endovascular graft from the native aorta as would be performed with a “chimney” or “periscope” technique for endovascular thoracoabdominal aortic repair then “**Endovascular Parallel Graft**” should be selected.

If the SMA is fed by a surgical bypass graft (e.g. iliac artery to SMA bypass, infrarenal aorta to SMA bypass – see below) with subsequent endovascular coverage of the SMA by the aortic endograft typically as would be performed in a “hybrid” thoracoabdominal aortic repair then “**Extra-anatomic Bypass**” should be selected.

If the SMA is covered by an aortic endograft with a fenestration/opening in the endograft corresponding to the location of the SMA and allowing continued antegrade flow into the SMA via this fenestration despite endograft coverage as would be performed in an endovascular juxtarenal or thoracoabdominal aortic repair then “**Fenestrated**” should be selected.

SEQ. #: 5280

Long Name: Superior Mesenteric - Extra-Anatomic Bypass - Aorta-Superior Mesenteric

Short Name: SupMesAortaSuMe

Definition: Indicate whether the extra-anatomic bypass was aorta to superior mesenteric

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

The bypass may originate from any segment of the aorta (e.g. ascending, descending, abdominal).

SEQ. #: 5300

Long Name: Superior Mesenteric - Extra-Anatomic Bypass - Iliac-Superior Mesenteric

Short Name: SupMesIliacSupMe

Definition: Indicate whether the extra-anatomic bypass was iliac to superior mesenteric

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

The bypass may originate from any segment of the iliac artery (e.g. common, external, or internal).

SEQ. #: 5315

Long Name: Superior Mesenteric - Extra-Anatomic Bypass - Other

Short Name: SupMesOther

Definition: Indicate whether another extra-anatomic superior mesenteric bypass was performed

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

This includes a bypass from any other vessel to the SMA.

SEQ. #: 5320

Long Name: Visceral Vessel Management - Right Renal

Short Name: RightRenal

Definition: Indicate management of the right renal artery

Intent/Clarification:

The intent is to clarify whether the right renal artery was revascularized during an endovascular repair of the thoracic or thoracoabdominal aorta.

If the right renal artery was left intact and continues to receive blood flow from the native aorta in the normal manner after the endovascular procedure then **“Native Flow”** should be selected.

If the right renal artery is instrumented with an endovascular branch graft extending from the main body of an aortic endograft and landing distally within the right renal artery as would be performed in an endovascular thoracoabdominal aortic repair then **“Endovascular Branch Graft”** should be selected.

If the right renal artery is instrumented with a separate endovascular graft that does not extend from an aortic endograft but rather courses parallel to an aortic endograft and lands distally within the right renal artery with flow through the right renal artery endovascular graft from the native aorta as would be performed with a “chimney” or “periscope” technique for endovascular thoracoabdominal aortic repair then **“Endovascular Parallel Graft”** should be selected.

If the right renal artery is fed by a surgical bypass graft (e.g. iliac artery to right renal artery bypass, infrarenal aorta to right renal artery bypass – see below) with subsequent endovascular coverage of the right renal artery by the aortic endograft typically as would be performed in a “hybrid” thoracoabdominal aortic repair then **“Extra-anatomic Bypass”** should be selected.

If the right renal artery is covered by an aortic endograft with a fenestration/opening in the endograft corresponding to the location of the right renal artery and allowing continued antegrade flow into the right renal artery via this fenestration despite endograft coverage as would be performed in an endovascular juxtarenal or thoracoabdominal aortic repair then **“Fenestrated”** should be selected.

SEQ. #: 5335

Long Name: Right Renal - Extra-Anatomic Bypass - Aorta-Right Renal

Short Name: RtRenAortaRtRe

Definition: Indicate whether the extra-anatomic bypass was aorta to right renal

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

The bypass may originate from any segment of the aorta (e.g. ascending, descending, abdominal).

SEQ. #: 5355

Long Name: Right Renal - Extra-Anatomic Bypass - Iliac-Right Renal

Short Name: RtRenIliacRtRen

Definition: Indicate whether the extra-anatomic bypass was iliac to right renal.

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

The bypass may originate from any segment of the aorta (e.g. ascending, descending, abdominal).

SEQ. #: 5365

Long Name: Right Renal - Extra-Anatomic Bypass - Other

Short Name: RtRenOther

Definition: Indicate whether another extra-anatomic right renal bypass was performed

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

This includes a bypass from any other vessel to the right renal artery. An example would include hepatorenal bypass.

SEQ. #: 5370

Long Name: Visceral Vessel Management - Left Renal

Short Name: LeftRenal

Definition: Indicate management of the left renal artery

Intent/Clarification:

The intent is to clarify whether the left renal artery was revascularized during an endovascular repair of the thoracic or thoracoabdominal aorta.

If the left renal artery was left intact and continues to receive blood flow from the native aorta in the normal manner after the endovascular procedure then “**Native Flow**” should be selected.

If the left renal artery is instrumented with an endovascular branch graft extending from the main body of an aortic endograft and landing distally within the left renal artery as would be performed in an endovascular thoracoabdominal aortic repair then **“Endovascular Branch Graft”** should be selected.

If the left renal artery is instrumented with a separate endovascular graft that does not extend from an aortic endograft but rather courses parallel to an aortic endograft and lands distally within the left renal artery with flow through the left renal artery endovascular graft from the native aorta as would be performed with a “chimney” or “periscope” technique for endovascular thoracoabdominal aortic repair then **“Endovascular Parallel Graft”** should be selected.

If the left renal artery is fed by a surgical bypass graft (e.g. iliac artery to left renal artery bypass, infrarenal aorta to left renal artery bypass – see below) with subsequent endovascular coverage of the left renal artery by the aortic endograft typically as would be performed in a “hybrid” thoracoabdominal aortic repair then **“Extra-anatomic Bypass”** should be selected.

If the left renal artery is covered by an aortic endograft with a fenestration/opening in the endograft corresponding to the location of the left renal artery and allowing continued antegrade flow into the left renal artery via this fenestration despite endograft coverage as would be performed in an endovascular juxtarenal or thoracoabdominal aortic repair then **“Fenestrated”** should be selected.

SEQ. #: 5375

Long Name: Left Renal - Extra-Anatomic Bypass - Aorta-Left Renal

Short Name: LtRenAortaLtRe

Definition: Indicate whether the extra-anatomic bypass was aorta to left renal

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

The bypass may originate from any segment of the aorta (e.g. ascending, descending, abdominal).

SEQ. #: 5380

Long Name: Left Renal - Extra-Anatomic Bypass - Iliac-Left Renal

Short Name: LtRenIliacLtRen

Definition: Indicate whether the extra-anatomic bypass was iliac to left renal

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

The bypass may originate from any segment of the iliac artery (e.g. common, external, or internal).

SEQ. #: 5385

Long Name: Left Renal - Extra-Anatomic Bypass - Other

Short Name: LtRenOther

Definition: Indicate whether another extra-anatomic left renal bypass was performed

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

This includes a bypass from any other vessel to the left renal artery. An example would include splenorenal bypass.

SEQ. #: 5390

Long Name: Visceral Vessel Management - Right Iliac

Short Name: RightIliac

Definition: Indicate management of the right iliac artery

Intent/Clarification:

The intent is to clarify whether the right iliac artery was revascularized during an endovascular repair of the thoracic or thoracoabdominal aorta.

If the right iliac artery was left intact and continues to receive blood flow from the native aorta in the normal manner after the endovascular procedure then “**Native Flow**” should be selected.

If the right iliac artery is instrumented with an iliac limb extending from the main body of an abdominal aortic endograft (either bifurcated or aorto uni-iliac) and landing distally within the right iliac artery then “**Bifurcated Graft**” should be selected.

If the right iliac artery is fed by a surgical bypass graft and not an endovascular device or native antegrade flow from the aorta then “**Extra-anatomic Bypass**” should be selected.

SEQ. #: 5391

Long Name: Visceral Vessel Management - Right Iliac - Femoral-Femoral

Short Name: RtIliacFemFem

Definition: Indicate whether the extra-anatomic bypass was femoral to femoral

Intent/Clarification:

This would typically be a bypass from the left common femoral artery to the right common femoral artery.

SEQ. #: 5392

Long Name: Visceral Vessel Management - Right Iliac - Other

Short Name: RtIliacOther

Definition: Indicate whether another right iliac extra-anatomic bypass was performed

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

This includes a bypass from any other vessel to the right iliac artery. An example would include aorto-iliac bypass.

SEQ. #: 5393

Long Name: Visceral Vessel Management - Left Iliac

Short Name: LeftIliac

Definition: Indicate management of the left iliac artery

Intent/Clarification:

The intent is to clarify whether the left iliac artery was revascularized during an endovascular repair of the thoracic or thoracoabdominal aorta.

If the left iliac artery was left intact and continues to receive blood flow from the native aorta in the normal manner after the endovascular procedure then “**Native Flow**” should be selected.

If the left iliac artery is instrumented with an iliac limb extending from the main body of an abdominal aortic endograft (either bifurcated or aorto uni-iliac) and landing distally within the left iliac artery then “**Bifurcated Graft**” should be selected.

If the left iliac artery is fed by a surgical bypass graft and not an endovascular device or native antegrade flow from the aorta then “**Extra-anatomic Bypass**” should be selected.

SEQ. #: 5394

Long Name: Visceral Vessel Management - Left Iliac - Femoral-Femoral

Short Name: LtIliacFemFem

Definition: Indicate whether the extra-anatomic bypass was femoral to femoral

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

This would typically be a bypass from the right common femoral artery to the left common femoral artery.

SEQ. #: 5395

Long Name: Visceral Vessel Management - Left Iliac - Other

Short Name: LtIliacOther

Definition: Indicate whether another left iliac extra-anatomic bypass was performed

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

This includes a bypass from any other vessel to the left iliac artery. An example would include aorto-iliac bypass.

SEQ. #: 5396

Long Name: Visceral Vessel Management - Internal Iliac Preserved

Short Name: IntIliacPres

Definition: Indicate whether the internal iliac was preserved

Intent/Clarification:

The intent is to clarify whether native antegrade flow is maintained within the internal iliac arteries during an endovascular repair of the thoracoabdominal aorta.

SEQ. #: 5397

Long Name: Visceral Vessel Management - Other visceral Vessels Extra-Anatomic Bypass

Short Name: OthVisVes

Definition: Indicate whether extra-anatomic bypass of other visceral vessels was performed

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

This includes a bypass to any branch of the major visceral vessels such as the hepatic or splenic branches of the celiac axis, a bypass to the inferior mesenteric artery or accessory renal artery, or a bypass to another named visceral vessel other than the celiac, SMA, left or right renal arteries.

SEQ. #: 5398

Long Name: Visceral Vessel Management - Other visceral Vessels Extra-Anatomic Bypass - Aorta-Other

Short Name: OthVisAortOth

Definition: Indicate whether other extra-anatomic bypass included an aorta to other bypass

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

The bypass may originate from any segment of the aorta (e.g. ascending, descending, abdominal).

SEQ. #: 5399

Long Name: Visceral Vessel Management - Other visceral Vessels Extra-Anatomic Bypass - Iliac-Other

Short Name: OthVisIliacOth

Definition: Indicate whether other extra-anatomic bypass included an iliac to other bypass

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

The bypass may originate from any segment of the iliac artery (e.g. common, external, or internal).

SEQ. #: 5400

Long Name: Visceral Vessel Management - Other visceral Vessels Extra-Anatomic Bypass - Other

Short Name: OthVisOther

Definition: Indicate whether any other visceral vessel extra-anatomic bypass was performed

Intent/Clarification:

An extra-anatomic bypass refers to any bypass graft that is placed outside of the normal anatomic vascular pathway to provide blood flow following or as part of that strategy.

This includes a bypass from any vessel other than the aorta or iliac artery to another named visceral vessel other than the celiac, SMA, left or right renal arteries.

SEQ. #: 5401

Long Name: Dissection Proximal Entry Tear Covered

Short Name: DisProxTearCov

Definition: Indicate whether the proximal entry tear was covered.

Intent/Clarification:

If the proximal entry tear (so-called primary tear) of an aortic dissection is fully covered by an aortic endograft then "Yes" should be selected.

SEQ. #: 5402

Long Name: Endoleak At End Of Procedure

Short Name: EndoEndProc

Definition: Indicate whether there was endoleak present at the end of the procedure

Intent/Clarification:

The intent is to define whether an endoleak is noted at the completion of an endovascular repair. This would typically be determined by the surgeon's assessment of the intraoperative completion angiogram.

An **endoleak** is defined as persistent blood flow in the aneurysm sac through and around the endovascular seal and is the most common complication after endovascular aneurysm repair.

SEQ. #: 5403

Long Name: Endoleak At End Of Procedure - Type

Short Name: EndoEndProcTy

Definition: Indicate the type of endoleak present

Intent/Clarification:

If an endoleak is noted **at the completion** of an endovascular repair before exiting the operating room, the intent is to define the type of endoleak present. This would typically be determined by the surgeon’s assessment of the intraoperative completion angiogram.

A **Type Ia endoleak** is defined as a leak occurring at the proximal seal zone.

A **Type Ib endoleak** is defined as a leak occurring at the distal seal zone.

A **Type II endoleak** is defined as retrograde filling of the aneurysm sac or false lumen in the case of dissection by aortic branch vessels (e.g. left subclavian artery, intercostal arteries, etc.).

A **Type III endoleak** is defined as leakage of blood into the aneurysm sac, or false lumen in the case of dissection, due to either a gap between separate endograft components, or a defect in the fabric of the graft secondary to graft strut fracture or erosion.

A **Type IV endoleak** is defined as the presence of an endoleak secondary to graft porosity. All other types of endoleaks must be definitively ruled out prior to selecting this diagnosis.

A **Type V endoleak**, also known as endotension, is defined as persistent aneurysm expansion in the absence of a confirmed endoleak.

SEQ. #: 5404

Long Name: Conversion To Open

Short Name: ConvToOpen

Definition: Indicate whether there was an unplanned conversion to an open procedure

Intent/Clarification:

This includes any conversion to open surgery not pre-specified as part of the operative plan.

SEQ. #: 5405

Long Name: Conversion To Open - Reason

Short Name: ConvToOpenRes

Definition: Indicate the reason for conversion to open procedure

Intent/Clarification:

If the reason for open conversion was failure of an endograft to deploy, either partially or fully, such that the planned endovascular treatment could not be completed then **“Deployment failure”** should be selected. If the reason for open conversion is a

persistent endoleak noted on completion angiogram then “**Endoleak**” should be selected. If the aorta or a branch vessel ruptures intraoperatively requiring open conversion then “**Rupture**” should be selected. If partial or complete (occlusion) loss of antegrade flow within a branch vessel occurs and requires open conversion to restore flow then “**Occlusion/loss of branch**” should be selected.

SEQ. #: 5406

Long Name: Intraop Dissection Extension

Short Name: IntDisExten

Definition: Indicate whether there was intraoperative dissection extension

Intent/Clarification:

If a pre-existing aortic dissection is made to propagate either proximally or distally beyond its preoperative extent during the operation then extension of dissection has occurred. If the pre-existing dissection extends proximally (i.e. back towards the aortic arch or ascending aorta) beyond the original extent then “**Retrograde**” should be selected. If the pre-existing dissection extends distally (i.e. downstream towards the descending or abdominal aorta) beyond the original extent then “**Antegrade**” should be selected. If the pre-existing dissection extends both proximally and distally then “**Both**” should be selected.

SEQ. #: 5407

Long Name: Unintentional Rupture Of Dissection Septum

Short Name: UnintRup

Definition: Indicate whether there was unintentional rupture of the dissection septum

Intent/Clarification:

The intent is to capture those instances where the dissection membrane/septum is unintentionally ruptured during an endovascular repair of an aortic dissection. This is typically due to the septum being fractured by a balloon or endograft, and the result is the creation of a new fenestration/connection between the true and false lumens of the dissection (so-called stent graft induced new entry tear (SINE)). This would typically be determined by the surgeon’s assessment of the intraoperative completion angiogram, intravascular ultrasound, and/or transesophageal echocardiography.

It should be noted that in certain instances the surgeon may elect to intentionally fracture/rupture the dissection septum/membrane, typically using a balloon, and these cases should not be coded as unintentional rupture.

SEQ. #: 5408

Long Name: Unintentional Rupture Of Dissection Septum - Location

Short Name: UnintRupLoc

Definition: Indicate the location of the unintentional rupture of the dissection septum

Intent/Clarification:

The exact aortic segment in which the unintentional rupture of the dissection septum occurred should be specified. This would typically be determined by the surgeon's assessment of the intraoperative completion angiogram, intravascular ultrasound, and/or transesophageal echocardiography.

Zone 0 definitions:

A. If the unintentional rupture of the dissection septum occurred in the aortic root (sinus of Valsalva segment just above the aortic valve) then "Below STJ" should be selected.

B. If the unintentional rupture of the dissection septum occurred in the segment of the ascending aorta between the sinotubular junction (defined as the junction between the aortic root and tubular ascending aorta) and the mid-point of the ascending aorta (i.e. proximal tubular ascending aorta) then "STJ-midascending" should be selected.

C. If the unintentional rupture of the dissection septum occurred in the segment of the ascending aorta between the mid-point of the ascending aorta and the origin of the innominate artery or first branch vessel off the aortic arch then "Midascending-distal ascending" should be selected.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

SEQ. #: 5420

Long Name: Spinal Drain

Short Name: SpinalDrain

Definition: Indicate when/if a spinal drain was placed

Intent/Clarification:

Indicate whether a cerebrospinal fluid drain was used for the thoracic aortic intervention. Cerebrospinal fluid (CSF) drainage is an adjunct to protect against paraplegia during aortic repairs. CSF pressure may increase during the perioperative period of aortic

repair leading to paraplegia. High CSF pressure may reduce spinal cord blood perfusion. CSF drainage reduces CSF pressure promoting spinal cord blood perfusion, reducing the risk of paraplegia. This field will capture the use of the cerebrospinal fluid drain during aortic repair. This is most often placed by anesthesia. This will also include any failed attempt (maldeployed).

SEQ. #: 5425

Long Name: IntraOp Motor Evoked Potential

Short Name: MotorEvoke

Definition: Indicate whether motor evoked potential was measured intraoperatively

Intent/Clarification:

Motor evoked potentials are used to monitor spinal cord function (motor cortex) during aortic intervention. Monitoring involves direct monitoring of electrical impulses used to stimulate the motor cortex and measurement of the response along the spinal cord. Changes in function during aortic intervention may indicate spinal cord injury. Adjunctive measures may be beneficial in restoring MEPs implying improvement in spinal cord function. It requires trained personnel, e.g. neurophysiologist, for monitoring. This will include unsuccessfully attempts at use of MEPs.

SEQ. #: 5426

Long Name: IntraOp Motor Evoked Potential - Documented MEP Abnormality

Short Name: MotorEvokeAb

Definition: Indicate whether any abnormality of motor evoked potential was documented

Intent/Clarification:

Motor evoked potentials are used to monitor spinal cord function (motor cortex) during aortic intervention. Monitoring involves direct monitoring of electrical impulses used to stimulate the motor cortex and measurement of the response along the spinal cord. Changes in function during aortic intervention may indicate spinal cord injury. Adjunctive measures may be beneficial in restoring MEPs implying improvement in spinal cord function. It requires trained personnel, e.g. neurophysiologist, for monitoring. This will include unsuccessfully attempts at use of MEPs.

SEQ. #: 5430

Long Name: IntraOp Somatosensory Evoked Potential

Short Name: SomatEvoke

Definition: indicate whether somatosensory evoked potential was measured intraoperatively

Intent/Clarification:

Somatosensory evoked potentials are used to monitor spinal cord function (sensory function) during aortic intervention. Monitoring involves direct monitoring of electrical impulses used to stimulate the sensory ventral tracts and measurement of the sensory response along the spinal cord. Changes in function during aortic intervention may indicate spinal cord injury. Adjunctive measures may be beneficial in restoring SSEPs implying improvement in spinal cord function. SSEPs may be less sensitive than MEPs for spinal cord dysfunction. It requires trained personnel, e.g. neurophysiologist, for monitoring. This will include unsuccessfully attempts at use of SSEPs.

SEQ. #: 5431**Long Name:** IntraOp Somatosensory Evoked Potential - Documented SEP Abnormality**Short Name:** SomatEvokeAb**Definition:** Indicate whether any abnormality of somatosensory evoked potential was documented**Intent/Clarification:**

Somatosensory evoked potentials are used to monitor spinal cord function (sensory function) during aortic intervention. Monitoring involves direct monitoring of electrical impulses used to stimulate the sensory ventral tracts and measurement of the sensory response along the spinal cord. Changes in function during aortic intervention may indicate spinal cord injury. Adjunctive measures may be beneficial in restoring SSEPs implying improvement in spinal cord function. SSEPs may be less sensitive than MEPs for spinal cord dysfunction. It requires trained personnel, e.g. neurophysiologist, for monitoring. This will include unsuccessfully attempts at use of SSEPs.

SEQ. #: 5432**Long Name:** IntraOp EEG**Short Name:** IntraOpEEG**Definition:** Indicate whether EEG was monitored intraoperatively**Intent/Clarification:**

Intraoperative electroencephalography may be used to monitor overall brain function during thoracic aortic procedures. Like MEPs and SSEPs, it requires trained personnel, e.g. neurophysiologist, for monitoring.

This will include unsuccessfully attempts at use of IntraOp/EEG.

FAQ October 2017: Can intraoperative Bispectral Index monitoring (BIS) be used as an acceptable substitute for EEG, since the measurement is a processed EEG parameter?

Answer: No, BIS does not provide the same information.

SEQ. #: 5433

Long Name: IntraOp EEG - Documented EEG Abnormality

Short Name: IntraOpEEGAb

Definition: Indicate whether any abnormality of intraoperative EEG was documented

Intent/Clarification:

Intraoperative electroencephalography may be used to monitor overall brain function during thoracic aortic procedures. Like MEPs and SSEPs, it requires trained personnel, e.g. neurophysiologist, for monitoring.

This will include unsuccessfully attempts at use of IntraOp/EEG.

SEQ. #: 5434

Long Name: IntraOp Intravascular Ultrasound (IVUS)

Short Name: IntraOpIVUS

Definition: indicate whether intravascular ultrasound was used intraoperatively

Intent/Clarification:

The unique point-of-view picture, generated in real time, yielding information that goes beyond what is possible with routine imaging methods, such as coronary angiography, performed in the Cath lab, or even non-invasive multi-slice CT scans.

SEQ. #: 5435

Long Name: IntraOp Transcutaneous Doppler

Short Name: TransDoppler

Definition: Indicate whether a transcutaneous Doppler was used intraoperatively

Intent/Clarification:

Transcutaneous Doppler enables the surgeon to alter his or her approach depending on the size and the location of aortic atheromatous burden, and provides an opportunity for intervention guidance during aortic cannulation, cross clamping and aortotomy.

SEQ. #: 5436

Long Name: IntraOp Angiogram

Short Name: IntraOpAng

Definition: Indicate whether an intraoperative angiogram was performed

Intent/Clarification:

An intraoperative angiography allows the surgeon to inspect the anatomic results of the surgical procedure.

SEQ. #: 5437

Long Name: IntraOp Angiogram - Volume Of Contrast

Short Name: IntraOpAngVol

Definition: Indicate the total volume of contrast given intraoperatively.

Intent/Clarification:

The volume of contrast used during the intraoperative angiogram documented in the perioperative record, the operative dictation, or the Cath Lab event log.

SEQ. #: 5438

Long Name: IntraOp Angiogram - Fluoroscopy Time In Minutes

Short Name: IntraOpAngFITm

Definition: Indicate the total intraoperative fluoroscopy time in minutes.

Intent/Clarification:

The total number of minute's intraoperative fluoroscopy documented in the perioperative record, the operative dictation, or the Cath Lab event log.

SEQ. #: 5440

Long Name: Aorta Device Inserted

Short Name: ADevIns

Definition: Indicate whether one or more devices were inserted into the aorta.

Intent/Clarification:

This will include all synthetic prosthetics inserted. This may include Dacron, PTFE, homografts, autografts, stents, and stentgrafts. Some aortic interventions may not require prosthetic materials or device implants such as primary repair of a pseudoaneurysm. This will be indicated as "No."

SEQ. #: 5450

Long Name: Aorta Device - Location #01

Short Name: ADevLoc01

Definition: Indicate the location within the aorta where device #01 was inserted.

Intent/Clarification:

Zone 0 is the Ascending Aorta and includes letter A-C. Verify exact location with CV Surgeon. Aortic Root (letter A) is below sinotubular junction.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

SEQ. #: 5455

Long Name: Aorta Device - Delivery Method #01

Short Name: ADevDelMeth01

Definition: Indicate the delivery method used to insert device #01 within the aorta.

Intent/Clarification:

For each device the method of implant should be specified as either “open” or “endovascular.” If a device was attempted to be implanted endo-vascularly but eventually implanted by open techniques, then this is designated “open.”

SEQ. #: 5460

Long Name: Aorta Device - Outcome #01

Short Name: ADevOut01

Definition: Indicate the outcome of the attempt to insert device #01.

Intent/Clarification:

Maldeployed: This indicates that a device implantation was attempted by endovascular means but was not successfully deployed in the intended position. An example would be attempted deployment of a fenestrated or branched device for transverse arch intervention and mal-alignment of the portal did not allow successful branch vessel bypass. If a device was mal-deployed but later removed this would be classified as “DelivMeth/deploy/remove.”

Deployed and Removed: This indicates that a device was attempted to be implanted by endovascular means but was not successfully deployed. An example would be open repair of descending thoracic aortic repair after failed thoracic endovascular aortic repair from rupture or persistent endoleak. This would be classified as TEVAR as “DelivMeth/deploy /remove.”

Successfully Deployed: This indicates that the device was successfully deployed by endovascular means.

SEQ. #: 5465

Long Name: Aorta Device - Model Number #01

Short Name: ADevModel01

Definition: Indicate the model number of aorta device #01.

Intent/Clarification:

This is the model number from the manufacturer related to the type of device implanted.

FAQ September 2017: Is there supposed to be a drop down list of model numbers for aorta devices in this field?

Answer: No, there is no drop down list.

SEQ. #: 5470

Long Name: Aorta Device - Unique Device Identifier #01

Short Name: ADevUDI01

Definition: Indicate the Unique Device Identifier (UDI) of aorta device #01 if available, otherwise leave blank. Note that the UDI is not the same as the serial number.

Intent/Clarification:

This is the number supplied from the manufacturer to identify the specific to the exact device inserted. This number is used to link the specific patient to the specific device implanted.

SEQ. #: 5475

Long Name: Aorta Device - Location #02

Short Name: ADevLoc02

Definition: Indicate the location within the aorta where device #02 was inserted, or indicate that no additional devices were inserted.

Intent/Clarification:

Zone 0 is the Ascending Aorta and includes letter A-C. Verify exact location with CV Surgeon. Aortic Root (letter A) is below sinotubular junction.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

SEQ. #: 5480

Long Name: Aorta Device - Delivery Method #02

Short Name: ADevDelMeth02

Definition: Indicate the delivery method used to insert device #02 within the aorta.

Intent/Clarification:

For each device the method of implant should be specified as either “open” or “endovascular.” If a device was attempted to be implanted endo-vascularly but eventually implanted by open techniques, then this is designated “open.”

SEQ. #: 5485

Long Name: Aorta Device - Outcome #02

Short Name: ADevOut02

Definition: Indicate the outcome of the attempt to insert device #02.

Intent/Clarification:

Maldeployed: This indicates that a device implantation was attempted by endovascular means but was not successfully deployed in the intended position. An example would be attempted deployment of a fenestrated or branched device for transverse arch intervention and mal-alignment of the portal did not allow successful branch vessel bypass. If a device was mal-deployed but later removed this would be classified as “DelivMeth/deploy/remove.”

Deployed and Removed: This indicates that a device was attempted to be implanted by endovascular means but was not successfully deployed. An example would be open repair of descending thoracic aortic repair after failed thoracic endovascular aortic repair

from rupture or persistent endoleak. This would be classified as TEVAR as “DelivMeth/deploy /remove.”

Successfully Deployed: This indicates that the device was successfully deployed by endovascular means.

SEQ. #: 5490

Long Name: Aorta Device - Model Number #02

Short Name: ADevModel02

Definition: Indicate the model number of aorta device #02.

Intent/Clarification:

This is the model number from the manufacturer related to the type of device implanted.

FAQ September 2017: Is there supposed to be a drop down list of model numbers for aorta devices in this field?

Answer: No, there is no drop down list.

SEQ. #: 5495

Long Name: Aorta Device - Unique Device Identifier #02

Short Name: ADevUDI02

Definition: Indicate the Unique Device Identifier (UDI) of aorta device #02 if available, otherwise leave blank. Note that the UDI is not the same as the serial number.

Intent/Clarification:

This is the number supplied from the manufacturer to identify the specific to the exact device inserted. This number is used to link the specific patient to the specific device implanted.

SEQ. #: 5500

Long Name: Aorta Device - Location #03

Short Name: ADevLoc03

Definition: Indicate the location within the aorta where device #03 was inserted, or indicate that no additional devices were inserted.

Intent/Clarification:

Zone 0 is the Ascending Aorta and includes letter A-C. Verify exact location with CV Surgeon. Aortic Root (letter A) is below sinotubular junction.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

SEQ. #: 5505

Long Name: Aorta Device - Delivery Method #03

Short Name: ADevDelMeth03

Definition: Indicate the delivery method used to insert device #03 within the aorta.

Intent/Clarification:

For each device the method of implant should be specified as either “open” or “endovascular.” If a device was attempted to be implanted endovascularly but eventually implanted by open techniques, then this is designated “open.”

SEQ. #: 5510

Long Name: Aorta Device - Outcome #03

Short Name: ADevOut03

Definition: Indicate the outcome of the attempt to insert device #03.

Intent/Clarification:

Maldeployed: This indicates that a device implantation was attempted by endovascular means but was not successfully deployed in the intended position. An example would be attempted deployment of a fenestrated or branched device for transverse arch intervention and mal-alignment of the portal did not allow successful branch vessel bypass. If a device was mal-deployed but later removed this would be classified as “DelivMeth/deploy/remove.”

Deployed and Removed: This indicates that a device was attempted to be implanted by endovascular means but was not successfully deployed. An example would be open repair of descending thoracic aortic repair after failed thoracic endovascular aortic repair from rupture or persistent endoleak. This would be classified as TEVAR as “DelivMeth/deploy /remove.”

Successfully Deployed: This indicates that the device was successfully deployed by endovascular means.

SEQ. #: 5515

Long Name: Aorta Device - Model Number #03

Short Name: ADevModel03

Definition: Indicate the model number of aorta device #03.

Intent/Clarification:

This is the model number from the manufacturer related to the type of device implanted.

FAQ September 2017: Is there supposed to be a drop down list of model numbers for aorta devices in this field?

Answer: No, there is no drop down list.

SEQ. #: 5520

Long Name: Aorta Device - Unique Device Identifier #03 **Short Name:** ADevUDI03

Definition: Indicate the Unique Device Identifier (UDI) of aorta device #03 if available, otherwise leave blank. Note that the UDI is not the same as the serial number.

Intent/Clarification:

This is the number supplied from the manufacturer to identify the specific to the exact device inserted. This number is used to link the specific patient to the specific device implanted.

SEQ. #: 5525

Long Name: Aorta Device - Location #04

Short Name: ADevLoc04

Definition: Indicate the location within the aorta where device #04 was inserted, or indicate that no additional devices were inserted.

Intent/Clarification:

Zone 0 is the Ascending Aorta and includes letter A-C. Verify exact location with CV Surgeon. Aortic Root (letter A) is below sinotubular junction.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

SEQ. #: 5530

Long Name: Aorta Device - Delivery Method #04

Short Name: ADevDelMeth04

Definition: Indicate the delivery method used to insert device #04 within the aorta.

Intent/Clarification:

For each device the method of implant should be specified as either “open” or “endovascular.” If a device was attempted to be implanted endo-vascularly but eventually implanted by open techniques, then this is designated “open.”

SEQ. #: 5535

Long Name: Aorta Device - Outcome #04

Short Name: ADevOut04

Definition: Indicate the outcome of the attempt to insert device #04.

Intent/Clarification:

Maldeployed: This indicates that a device implantation was attempted by endovascular means but was not successfully deployed in the intended position. An example would be attempted deployment of a fenestrated or branched device for transverse arch intervention and mal-alignment of the portal did not allow successful branch vessel bypass. If a device was mal-deployed but later removed this would be classified as “DelivMeth/deploy/remove.”

Deployed and Removed: This indicates that a device was attempted to be implanted by endovascular means but was not successfully deployed. An example would be open repair of descending thoracic aortic repair after failed thoracic endovascular aortic repair from rupture or persistent endoleak. This would be classified as TEVAR as “DelivMeth/deploy /remove.”

Successfully Deployed: This indicates that the device was successfully deployed by endovascular means.

SEQ. #: 5540

Long Name: Aorta Device - Model Number #04

Short Name: ADevModel04

Definition: Indicate the model number of aorta device #04.

Intent/Clarification:

This is the model number from the manufacturer related to the type of device implanted.

FAQ September 2017: Is there supposed to be a drop down list of model numbers for aorta devices in this field?

Answer: No, there is no drop down list.

SEQ. #: 5545

Long Name: Aorta Device - Unique Device Identifier #04

Short Name: ADevUDI04

Definition: Indicate the Unique Device Identifier (UDI) of aorta device #04 if available, otherwise leave blank. Note that the UDI is not the same as the serial number.

Intent/Clarification:

This is the number supplied from the manufacturer to identify the specific to the exact device inserted. This number is used to link the specific patient to the specific device implanted.

SEQ. #: 5550

Long Name: Aorta Device - Location #05 **Short Name:** ADevLoc05

Definition: Indicate the location within the aorta where device #05 was inserted, or indicate that no additional devices were inserted.

Intent/Clarification:

Zone 0 is the Ascending Aorta and includes letter A-C. Verify exact location with CV Surgeon. Aortic Root (letter A) is below sinotubular junction.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

SEQ. #: 5555

Long Name: Aorta Device - Delivery Method #05

Short Name: ADevDelMeth05

Definition: Indicate the delivery method used to insert device #05 within the aorta.

Intent/Clarification:

For each device the method of implant should be specified as either “open” or “endovascular.” If a device was attempted to be implanted endo-vascularly but eventually implanted by open techniques, then this is designated “open.”

SEQ. #: 5560

Long Name: Aorta Device - Outcome #05

Short Name: ADevOut05

Definition: Indicate the outcome of the attempt to insert device #05.

Intent/Clarification:

Maldeployed: This indicates that a device implantation was attempted by endovascular means but was not successfully deployed in the intended position. An example would be attempted deployment of a fenestrated or branched device for transverse arch intervention and mal-alignment of the portal did not allow successful branch vessel bypass. If a device was mal-deployed but later removed this would be classified as “DelivMeth/deploy/remove.”

Deployed and Removed: This indicates that a device was attempted to be implanted by endovascular means but was not successfully deployed. An example would be open repair of descending thoracic aortic repair after failed thoracic endovascular aortic repair from rupture or persistent endoleak. This would be classified as TEVAR as “DelivMeth/deploy /remove.”

Successfully Deployed: This indicates that the device was successfully deployed by endovascular means.

SEQ. #: 5565

Long Name: Aorta Device - Model Number #05

Short Name: ADevModel05

Definition: Indicate the model number of aorta device #05.

Intent/Clarification:

This is the model number from the manufacturer related to the type of device implanted.

FAQ September 2017: Is there supposed to be a drop down list of model numbers for aorta devices in this field?

Answer: No, there is no drop down list.

SEQ. #: 5570

Long Name: Aorta Device - Unique Device Identifier #05

Short Name: ADevUDI05

Definition: Indicate the Unique Device Identifier (UDI) of aorta device #05 if available, otherwise leave blank. Note that the UDI is not the same as the serial number.

Intent/Clarification:

This is the number supplied from the manufacturer to identify the specific to the exact device inserted. This number is used to link the specific patient to the specific device implanted.

SEQ. #: 5575

Long Name: Aorta Device - Location #06

Short Name: ADevLoc06

Definition: Indicate the location within the aorta where device #06 was inserted, or indicate that no additional devices were inserted.

Intent/Clarification:

Zone 0 is the Ascending Aorta and includes letter A-C. Verify exact location with CV Surgeon. Aortic Root (letter A) is below sinotubular junction.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

SEQ. #: 5580

Long Name: Aorta Device - Delivery Method #06 **Short Name:** ADevDelMeth06

Definition: Indicate the delivery method used to insert device #06 within the aorta.

Intent/Clarification:

For each device the method of implant should be specified as either “open” or “endovascular.” If a device was attempted to be implanted endo-vascularly but eventually implanted by open techniques, then this is designated “open.”

SEQ. #: 5585

Long Name: Aorta Device - Outcome #06

Short Name: ADevOut06

Definition: Indicate the outcome of the attempt to insert device #06.

Intent/Clarification:

Maldeployed: This indicates that a device implantation was attempted by endovascular means but was not successfully deployed in the intended position. An example would be attempted deployment of a fenestrated or branched device for transverse arch intervention and mal-alignment of the portal did not allow successful branch vessel bypass. If a device was mal-deployed but later removed this would be classified as “DelivMeth/deploy/remove.”

Deployed and Removed: This indicates that a device was attempted to be implanted by endovascular means but was not successfully deployed. An example would be open repair of descending thoracic aortic repair after failed thoracic endovascular aortic repair from rupture or persistent endoleak. This would be classified as TEVAR as “DelivMeth/deploy /remove.”

Successfully Deployed: This indicates that the device was successfully deployed by endovascular means.

SEQ. #: 5590

Long Name: Aorta Device - Model Number #06

Short Name: ADevModel06

Definition: Indicate the model number of aorta device #06.

Intent/Clarification:

This is the model number from the manufacturer related to the type of device implanted.
FAQ September 2017: Is there supposed to be a drop down list of model numbers for aorta devices in this field?

Answer: No, there is no drop down list.

SEQ. #: 5595

Long Name: Aorta Device - Unique Device Identifier #06

Short Name: ADevUDI06

Definition: Indicate the Unique Device Identifier (UDI) of aorta device #06 if available, otherwise leave blank. Note that the UDI is not the same as the serial number.

Intent/Clarification:

This is the number supplied from the manufacturer to identify the specific to the exact device inserted. This number is used to link the specific patient to the specific device implanted.

SEQ. #: 5600

Long Name: Aorta Device - Location #07

Short Name: ADevLoc07

Definition: Indicate the location within the aorta where device #07 was inserted, or indicate that no additional devices were inserted.

Intent/Clarification:

Zone 0 is the Ascending Aorta and includes letter A-C. Verify exact location with CV Surgeon. Aortic Root (letter A) is below sinotubular junction.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

SEQ. #: 5605

Long Name: Aorta Device - Delivery Method #07

Short Name: ADevDelMeth07

Definition: Indicate the delivery method used to insert device #07 within the aorta.

Intent/Clarification:

For each device the method of implant should be specified as either “open” or “endovascular.” If a device was attempted to be implanted endo-vascularly but eventually implanted by open techniques, then this is designated “open.”

SEQ. #: 5610

Long Name: Aorta Device - Outcome #07

Short Name: ADevOut07

Definition: Indicate the outcome of the attempt to insert device #07.

Intent/Clarification:

Maldeployed: This indicates that a device implantation was attempted by endovascular means but was not successfully deployed in the intended position. An example would be attempted deployment of a fenestrated or branched device for transverse arch intervention and mal-alignment of the portal did not allow successful branch vessel bypass. If a device was mal-deployed but later removed this would be classified as “DelivMeth/deploy/remove.”

Deployed and Removed: This indicates that a device was attempted to be implanted by endovascular means but was not successfully deployed. An example would be open repair of descending thoracic aortic repair after failed thoracic endovascular aortic repair from rupture or persistent endoleak. This would be classified as TEVAR as “DelivMeth/deploy /remove.”

Successfully Deployed: This indicates that the device was successfully deployed by endovascular means.

SEQ. #: 5615

Long Name: Aorta Device - Model Number #07

Short Name: ADevModel07

Definition: Indicate the model number of aorta device #07.

Intent/Clarification:

This is the model number from the manufacturer related to the type of device implanted.

FAQ September 2017: Is there supposed to be a drop down list of model numbers for aorta devices in this field?

Answer: No, there is no drop down list.

SEQ. #: 5620

Long Name: Aorta Device - Unique Device Identifier #07

Short Name: ADevUDI07

Definition: Indicate the Unique Device Identifier (UDI) of aorta device #07 if available, otherwise leave blank. Note that the UDI is not the same as the serial number.

Intent/Clarification:

This is the number supplied from the manufacturer to identify the specific to the exact device inserted. This number is used to link the specific patient to the specific device implanted.

SEQ. #: 5625

Long Name: Aorta Device - Location #08

Short Name: ADevLoc08

Definition: Indicate the location within the aorta where device #08 was inserted, or indicate that no additional devices were inserted.

Intent/Clarification:

Zone 0 is the Ascending Aorta and includes letter A-C. Verify exact location with CV Surgeon. Aortic Root (letter A) is below sinotubular junction.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

SEQ. #: 5630

Long Name: Aorta Device - Delivery Method #08

Short Name: ADevDelMeth08

Definition: Indicate the delivery method used to insert device #08 within the aorta.

Intent/Clarification:

For each device the method of implant should be specified as either “open” or “endovascular.” If a device was attempted to be implanted endo-vascularly but eventually implanted by open techniques, then this is designated “open.”

SEQ. #: 5635

Long Name: Aorta Device - Outcome #08

Short Name: ADevOut08

Definition: Indicate the outcome of the attempt to insert device #08.

Intent/Clarification:

Maldeployed: This indicates that a device implantation was attempted by endovascular means but was not successfully deployed in the intended position. An example would be attempted deployment of a fenestrated or branched device for transverse arch intervention and mal-alignment of the portal did not allow successful branch vessel bypass. If a device was mal-deployed but later removed this would be classified as “DelivMeth/deploy/remove.”

Deployed and Removed: This indicates that a device was attempted to be implanted by endovascular means but was not successfully deployed. An example would be open repair of descending thoracic aortic repair after failed thoracic endovascular aortic repair from rupture or persistent endoleak. This would be classified as TEVAR as “DelivMeth/deploy /remove.”

Successfully Deployed: This indicates that the device was successfully deployed by endovascular means.

SEQ. #: 5640

Long Name: Aorta Device - Model Number #08

Short Name: ADevModel08

Definition: Indicate the model number of aorta device #08.

Intent/Clarification:

This is the model number from the manufacturer related to the type of device implanted.

FAQ September 2017: Is there supposed to be a drop down list of model numbers for aorta devices in this field?

Answer: No, there is no drop down list.

SEQ. #: 5645

Long Name: Aorta Device - Unique Device Identifier #08

Short Name: ADevUDI08

Definition: Indicate the Unique Device Identifier (UDI) of aorta device #08 if available, otherwise leave blank. Note that the UDI is not the same as the serial number.

Intent/Clarification:

This is the number supplied from the manufacturer to identify the specific to the exact device inserted. This number is used to link the specific patient to the specific device implanted.

SEQ. #: 5650

Long Name: Aorta Device - Location #09

Short Name: ADevLoc09

Definition: Indicate the location within the aorta where device #09 was inserted, or indicate that no additional devices were inserted.

Intent/Clarification:

Zone 0 is the Ascending Aorta and includes letter A-C. Verify exact location with CV Surgeon. Aortic Root (letter A) is below sinotubular junction.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

SEQ. #: 5655

Long Name: Aorta Device - Delivery Method #09

Short Name: ADevDelMeth09

Definition: Indicate the delivery method used to insert device #09 within the aorta.

Intent/Clarification:

For each device the method of implant should be specified as either “open” or “endovascular.” If a device was attempted to be implanted endo-vascularly but eventually implanted by open techniques, then this is designated “open.”

SEQ. #: 5660

Long Name: Aorta Device - Outcome #09

Short Name: ADevOut09

Definition: Indicate the outcome of the attempt to insert device #09.

Intent/Clarification:

Maldeployed: This indicates that a device implantation was attempted by endovascular means but was not successfully deployed in the intended position. An example would be attempted deployment of a fenestrated or branched device for transverse arch intervention and mal-alignment of the portal did not allow successful branch vessel bypass. If a device was mal-deployed but later removed this would be classified as “DelivMeth/deploy/remove.”

Deployed and Removed: This indicates that a device was attempted to be implanted by endovascular means but was not successfully deployed. An example would be open repair of descending thoracic aortic repair after failed thoracic endovascular aortic repair from rupture or persistent endoleak. This would be classified as TEVAR as “DelivMeth/deploy /remove.”

Successfully Deployed: This indicates that the device was successfully deployed by endovascular means.

SEQ. #: 5665

Long Name: Aorta Device - Model Number #09

Short Name: ADevModel09

Definition: Indicate the model number of aorta device #09.

Intent/Clarification:

This is the model number from the manufacturer related to the type of device implanted.

FAQ September 2017: Is there supposed to be a drop down list of model numbers for aorta devices in this field?

Answer: No, there is no drop down list.

SEQ. #: 5670

Long Name: Aorta Device - Unique Device Identifier #09

Short Name: ADevUDI09

Definition: Indicate the Unique Device Identifier (UDI) of aorta device #09 if available, otherwise leave blank. Note that the UDI is not the same as the serial number.

Intent/Clarification:

This is the number supplied from the manufacturer to identify the specific to the exact device inserted. This number is used to link the specific patient to the specific device implanted.

SEQ. #: 5675

Long Name: Aorta Device - Location #10

Short Name: ADevLoc10

Definition: Indicate the location within the aorta where device #10 was inserted, or indicate that no additional devices were inserted.

Intent/Clarification:

Zone 0 is the Ascending Aorta and includes letter A-C. Verify exact location with CV Surgeon. Aortic Root (letter A) is below sinotubular junction.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

SEQ. #: 5680

Long Name: Aorta Device - Delivery Method #10

Short Name: ADevDelMeth10

Definition: Indicate the delivery method used to insert device #10 within the aorta.

Intent/Clarification:

For each device the method of implant should be specified as either “open” or “endovascular.” If a device was attempted to be implanted endovascularly but eventually implanted by open techniques, then this is designated “open.”

SEQ. #: 5685

Long Name: Aorta Device - Outcome #10

Short Name: ADevOut10

Definition: Indicate the outcome of the attempt to insert device #10.

Intent/Clarification:

Maldeployed: This indicates that a device implantation was attempted by endovascular means but was not successfully deployed in the intended position. An example would be attempted deployment of a fenestrated or branched device for transverse arch intervention and mal-alignment of the portal did not allow successful branch vessel bypass. If a device was mal-deployed but later removed this would be classified as “DelivMeth/deploy/remove.”

Deployed and Removed: This indicates that a device was attempted to be implanted by endovascular means but was not successfully deployed. An example would be open repair of descending thoracic aortic repair after failed thoracic endovascular aortic repair from rupture or persistent endoleak. This would be classified as TEVAR as “DelivMeth/deploy /remove.”

Successfully Deployed: This indicates that the device was successfully deployed by endovascular means.

SEQ. #: 5690

Long Name: Aorta Device - Model Number #10

Short Name: ADevModel10

Definition: Indicate the model number of aorta device #10.

Intent/Clarification:

This is the model number from the manufacturer related to the type of device implanted.

FAQ September 2017: Is there supposed to be a drop down list of model numbers for aorta devices in this field?

Answer: No, there is no drop down list.

SEQ. #: 5695

Long Name: Aorta Device - Unique Device Identifier #10

Short Name: ADevUDI10

Definition: Indicate the Unique Device Identifier (UDI) of aorta device #10 if available, otherwise leave blank. Note that the UDI is not the same as the serial number.

Intent/Clarification:

This is the number supplied from the manufacturer to identify the specific to the exact device inserted. This number is used to link the specific patient to the specific device implanted.

SEQ. #: 5700

Long Name: Aorta Device - Location #11

Short Name: ADevLoc11

Definition: Indicate the location within the aorta where device #11 was inserted, or indicate that no additional devices were inserted.

Intent/Clarification:

Zone 0 is the Ascending Aorta and includes letter A-C. Verify exact location with CV Surgeon. Aortic Root (letter A) is below sinotubular junction.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

SEQ. #: 5705

Long Name: Aorta Device - Delivery Method #11

Short Name: ADevDelMeth11

Definition: Indicate the delivery method used to insert device #11 within the aorta.

Intent/Clarification:

For each device the method of implant should be specified as either “open” or “endovascular.” If a device was attempted to be implanted endovascularly but eventually implanted by open techniques, then this is designated “open.”

SEQ. #: 5710

Long Name: Aorta Device - Outcome #11

Short Name: ADevOut11

Definition: Indicate the outcome of the attempt to insert device #11.

Intent/Clarification:

Maldeployed: This indicates that a device implantation was attempted by endovascular means but was not successfully deployed in the intended position. An example would be attempted deployment of a fenestrated or branched device for transverse arch intervention and mal-alignment of the portal did not allow successful branch vessel bypass. If a device was mal-deployed but later removed this would be classified as “DelivMeth/deploy/remove.”

Deployed and Removed: This indicates that a device was attempted to be implanted by endovascular means but was not successfully deployed. An example would be open repair of descending thoracic aortic repair after failed thoracic endovascular aortic repair from rupture or persistent endoleak. This would be classified as TEVAR as “DelivMeth/deploy /remove.”

Successfully Deployed: This indicates that the device was successfully deployed by endovascular means.

SEQ. #: 5715

Long Name: Aorta Device - Model Number #11

Short Name: ADevModel11

Definition: Indicate the model number of aorta device #11.

Intent/Clarification:

This is the model number from the manufacturer related to the type of device implanted.

FAQ September 2017: Is there supposed to be a drop down list of model numbers for aorta devices in this field?

Answer: No, there is no drop down list.

SEQ. #: 5720

Long Name: Aorta Device - Unique Device Identifier #11

Short Name: ADevUDI11

Definition: Indicate the Unique Device Identifier (UDI) of aorta device #11 if available, otherwise leave blank. Note that the UDI is not the same as the serial number.

Intent/Clarification:

This is the number supplied from the manufacturer to identify the specific to the exact device inserted. This number is used to link the specific patient to the specific device implanted.

SEQ. #: 5725

Long Name: Aorta Device - Location #12

Short Name: ADevLoc12

Definition: Indicate the location within the aorta where device #12 was inserted, or indicate that no additional devices were inserted.

Intent/Clarification:

Zone 0 is the Ascending Aorta and includes letter A-C. Verify exact location with CV Surgeon. Aortic Root (letter A) is below sinotubular junction.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

SEQ. #: 5730

Long Name: Aorta Device - Delivery Method #12

Short Name: ADevDelMeth12

Definition: Indicate the delivery method used to insert device #12 within the aorta.

Intent/Clarification:

For each device the method of implant should be specified as either “open” or “endovascular.” If a device was attempted to be implanted endovascularly but eventually implanted by open techniques, then this is designated “open.”

SEQ. #: 5735

Long Name: Aorta Device - Outcome #12

Short Name: ADevOut12

Definition: Indicate the outcome of the attempt to insert device #12.

Intent/Clarification:

Maldeployed: This indicates that a device implantation was attempted by endovascular means but was not successfully deployed in the intended position. An example would be attempted deployment of a fenestrated or branched device for transverse arch intervention and mal-alignment of the portal did not allow successful branch vessel bypass. If a device was mal-deployed but later removed this would be classified as “DelivMeth/deploy/remove.”

Deployed and Removed: This indicates that a device was attempted to be implanted by endovascular means but was not successfully deployed. An example would be open repair of descending thoracic aortic repair after failed thoracic endovascular aortic repair from rupture or persistent endoleak. This would be classified as TEVAR as “DelivMeth/deploy /remove.”

Successfully Deployed: This indicates that the device was successfully deployed by endovascular means.

SEQ. #: 5740

Long Name: Aorta Device - Model Number #12

Short Name: ADevModel12

Definition: Indicate the model number of aorta device #12.

Intent/Clarification:

This is the model number from the manufacturer related to the type of device implanted.

FAQ September 2017: Is there supposed to be a drop down list of model numbers for aorta devices in this field?

Answer: No, there is no drop down list.

SEQ. #: 5745

Long Name: Aorta Device - Unique Device Identifier #12

Short Name: ADevUDI12

Definition: Indicate the Unique Device Identifier (UDI) of aorta device #12 if available, otherwise leave blank. Note that the UDI is not the same as the serial number.

Intent/Clarification:

This is the number supplied from the manufacturer to identify the specific to the exact device inserted. This number is used to link the specific patient to the specific device implanted.

SEQ. #: 5750

Long Name: Aorta Device - Location #13

Short Name: ADevLoc13

Definition: Indicate the location within the aorta where device #13 was inserted, or indicate that no additional devices were inserted.

Intent/Clarification:

Zone 0 is the Ascending Aorta and includes letter A-C. Verify exact location with CV Surgeon. Aortic Root (letter A) is below sinotubular junction.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

SEQ. #: 5755

Long Name: Aorta Device - Delivery Method #13

Short Name: ADevDelMeth13

Definition: Indicate the delivery method used to insert device #13 within the aorta.

Intent/Clarification:

For each device the method of implant should be specified as either “open” or “endovascular.” If a device was attempted to be implanted endovascularly but eventually implanted by open techniques, then this is designated “open.”

SEQ. #: 5760

Long Name: Aorta Device - Outcome #13

Short Name: ADevOut13

Definition: Indicate the outcome of the attempt to insert device #13.

Intent/Clarification:

Maldeployed: This indicates that a device implantation was attempted by endovascular means but was not successfully deployed in the intended position. An example would be attempted deployment of a fenestrated or branched device for transverse arch intervention and mal-alignment of the portal did not allow successful branch vessel bypass. If a device was mal-deployed but later removed this would be classified as “DelivMeth/deploy/remove.”

Deployed and Removed: This indicates that a device was attempted to be implanted by endovascular means but was not successfully deployed. An example would be open repair of descending thoracic aortic repair after failed thoracic endovascular aortic repair from rupture or persistent endoleak. This would be classified as TEVAR as “DelivMeth/deploy /remove.”

Successfully Deployed: This indicates that the device was successfully deployed by endovascular means.

SEQ. #: 5765

Long Name: Aorta Device - Model Number #13

Short Name: ADevModel13

Definition: Indicate the model number of aorta device #13.

Intent/Clarification:

This is the model number from the manufacturer related to the type of device implanted.

FAQ September 2017: Is there supposed to be a drop down list of model numbers for aorta devices in this field?

Answer: No, there is no drop down list.

SEQ. #: 5770

Long Name: Aorta Device - Unique Device Identifier #13

Short Name: ADevUDI13

Definition: Indicate the Unique Device Identifier (UDI) of aorta device #13 if available, otherwise leave blank. Note that the UDI is not the same as the serial number.

Intent/Clarification:

This is the number supplied from the manufacturer to identify the specific to the exact device inserted. This number is used to link the specific patient to the specific device implanted.

SEQ. #: 5775

Long Name: Aorta Device - Location #14

Short Name: ADevLoc14

Definition: Indicate the location within the aorta where device #14 was inserted, or indicate that no additional devices were inserted.

Intent/Clarification:

Zone 0 is the Ascending Aorta and includes letter A-C. Verify exact location with CV Surgeon. Aortic Root (letter A) is below sinotubular junction.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

SEQ. #: 5780

Long Name: Aorta Device - Delivery Method #14

Short Name: ADevDelMeth14

Definition: Indicate the delivery method used to insert device #14 within the aorta.

Intent/Clarification:

For each device the method of implant should be specified as either “open” or “endovascular.” If a device was attempted to be implanted endovascularly but eventually implanted by open techniques, then this is designated “open.”

SEQ. #: 5785

Long Name: Aorta Device - Outcome #14

Short Name: ADevOut14

Definition: Indicate the outcome of the attempt to insert device #14.

Intent/Clarification:

Maldeployed: This indicates that a device implantation was attempted by endovascular means but was not successfully deployed in the intended position. An example would be attempted deployment of a fenestrated or branched device for transverse arch intervention and mal-alignment of the portal did not allow successful branch vessel bypass. If a device was mal-deployed but later removed this would be classified as “DelivMeth/deploy/remove.”

Deployed and Removed: This indicates that a device was attempted to be implanted by endovascular means but was not successfully deployed. An example would be open repair of descending thoracic aortic repair after failed thoracic endovascular aortic repair from rupture or persistent endoleak. This would be classified as TEVAR as “DelivMeth/deploy /remove.”

Successfully Deployed: This indicates that the device was successfully deployed by endovascular means.

SEQ. #: 5790

Long Name: Aorta Device - Model Number #14

Short Name: ADevModel14

Definition: Indicate the model number of aorta device #14.

Intent/Clarification:

This is the model number from the manufacturer related to the type of device implanted.

FAQ September 2017: Is there supposed to be a drop down list of model numbers for aorta devices in this field?

Answer: No, there is no drop down list.

SEQ. #: 5795

Long Name: Aorta Device - Unique Device Identifier #14

Short Name: ADevUDI14

Definition: Indicate the Unique Device Identifier (UDI) of aorta device #14 if available, otherwise leave blank. Note that the UDI is not the same as the serial number.

Intent/Clarification:

This is the number supplied from the manufacturer to identify the specific to the exact device inserted. This number is used to link the specific patient to the specific device implanted.

SEQ. #: 5800

Long Name: Aorta Device - Location #15

Short Name: ADevLoc15

Definition: Indicate the location within the aorta where device #15 was inserted, or indicate that no additional devices were inserted.

Intent/Clarification:

Zone 0 is the Ascending Aorta and includes letter A-C. Verify exact location with CV Surgeon. Aortic Root (letter A) is below sinotubular junction.

- A. Below sinotubular junction
- B. Sinotubular junction to mid ascending
- C. Mid ascending to distal ascending
- D. Zone 1 (between innominate and left carotid)
- E. Zone 2 (between left carotid and left subclavian)
- F. Zone 3 (first 2 cm. distal to left subclavian)
- G. Zone 4 (end of zone 3 to mid descending aorta ~ T6)
- H. Zone 5 (mid descending aorta to celiac)
- I. Zone 6 (celiac to superior mesenteric)
- J. Zone 7 (superior mesenteric to renals)
- K. Zone 8 (renal to infra-renal abdominal aorta)
- L. Zone 9 (infrarenal abdominal aorta)
- M. Zone 10 (common iliac)

- N. Zone 11 (external iliacs)

SEQ. #: 5805

Long Name: Aorta Device - Delivery Method #15

Short Name: ADevDelMeth15

Definition: Indicate the delivery method used to insert device #15 within the aorta.

Intent/Clarification:

For each device the method of implant should be specified as either “open” or “endovascular.” If a device was attempted to be implanted endovascularly but eventually implanted by open techniques, then this is designated “open.”

SEQ. #: 5810

Long Name: Aorta Device - Outcome #15

Short Name: ADevOut15

Definition: Indicate the outcome of the attempt to insert device #15.

Intent/Clarification:

Maldeployed: This indicates that a device implantation was attempted by endovascular means but was not successfully deployed in the intended position. An example would be attempted deployment of a fenestrated or branched device for transverse arch intervention and mal-alignment of the portal did not allow successful branch vessel bypass. If a device was mal-deployed but later removed this would be classified as “DelivMeth/deploy/remove.”

Deployed and Removed: This indicates that a device was attempted to be implanted by endovascular means but was not successfully deployed. An example would be open repair of descending thoracic aortic repair after failed thoracic endovascular aortic repair from rupture or persistent endoleak. This would be classified as TEVAR as “DelivMeth/deploy /remove.”

Successfully Deployed: This indicates that the device was successfully deployed by endovascular means.

SEQ. #: 5815

Long Name: Aorta Device - Model Number #15

Short Name: ADevModel15

Definition: Indicate the model number of aorta device #15.

Intent/Clarification:

This is the model number from the manufacturer related to the type of device implanted.

FAQ September 2017: Is there supposed to be a drop down list of model numbers for aorta devices in this field?

Answer: No, there is no drop down list.

SEQ. #: 5820

Long Name: Aorta Device - Unique Device Identifier #15

Short Name: ADevUDI15

Definition: Indicate the Unique Device Identifier (UDI) of aorta device #15 if available, otherwise leave blank. Note that the UDI is not the same as the serial number.

Intent/Clarification:

This is the number supplied from the manufacturer to identify the specific to the exact device inserted. This number is used to link the specific patient to the specific device implanted.

Congenital Defect Repair

SEQ. #: 6500

Long Name: Other Card-Congenital Diagnosis 1

Short Name: OCarCongDiag1

Definition: Indicate the first of the three most significant congenital diagnoses.

Intent/Clarification:

A comprehensive list of diagnoses is available at (under V2.9 Congenital Diagnoses and Procedure List):

<http://www.sts.org/sts-national-database/database-managers/adult-cardiac-surgery-database/data-collection>

SEQ. #: 6505

Long Name: Other Card-Congenital Diagnosis 2

Short Name: OCarCongDiag2

Definition: Indicate the second of the three most significant congenital diagnoses.

Intent/Clarification:

A comprehensive list of diagnoses is available at (under V2.9 Congenital Diagnoses and Procedure List):

<http://www.sts.org/sts-national-database/database-managers/adult-cardiac-surgery-database/data-collection>

SEQ. #: 6510

Long Name: Other Card-Congenital Diagnosis 3

Short Name: OCarCongDiag3

Definition: Indicate the third of the three most significant congenital diagnoses.

Intent/Clarification:

A comprehensive list of diagnoses is available at (under V2.9 Congenital Diagnoses and Procedure List):

<http://www.sts.org/sts-national-database/database-managers/adult-cardiac-surgery-database/data-collection>

SEQ. #: 6515

Long Name: Other Card-Congenital Procedure 1

Short Name: OCarCongProc1

Definition: Indicate the first of the three most significant congenital procedures.

Intent/Clarification:

A comprehensive list of diagnoses is available at (under V2.9 Congenital Diagnoses and Procedure List):

<http://www.sts.org/sts-national-database/database-managers/adult-cardiac-surgery-database/data-collection>

SEQ. #: 6520

Long Name: Other Card-Congenital Procedure 2

Short Name: OCarCongProc2

Definition: Indicate the second of the three most significant congenital procedures.

Intent/Clarification:

A comprehensive list of diagnoses is available at (under V2.9 Congenital Diagnoses and Procedure List):

<http://www.sts.org/sts-national-database/database-managers/adult-cardiac-surgery-database/data-collection>

SEQ. #: 6525

Long Name: Other Card-Congenital Procedure 3

Short Name: OCarCongProc3

Definition: Indicate the third of the three most significant congenital procedures.

Intent/Clarification:

A comprehensive list of diagnoses is available at (under V2.9 Congenital Diagnoses and Procedure List):

<http://www.sts.org/sts-national-database/database-managers/adult-cardiac-surgery-database/data-collection>

Other Non-Cardiac Procedures

SEQ. #: 6530

Long Name: Other Non Card-Caro Endart

Short Name: ONCCarEn

Definition: Indicate whether the patient underwent surgical removal of stenotic atheromatous plaque or percutaneous/surgical placement of carotid stent in conjunction with the primary surgical procedure.

Intent/Clarification: Right and/or left carotid arteries are branches of the arch of the aorta that transverse the neck and supply blood flow to the brain.

- Yes, planned
 - Yes, unplanned due to surgical complication
 - Yes, unplanned due to unsuspected disease or anatomy
 - No
-
-

SEQ. #: 6535

Long Name: Other Non Card-Other Vasc

Short Name: ONCOVasc

Definition: Indicate whether patient had procedures treating peripheral vascular disease or condition in conjunction with the primary surgical procedure.

Intent/Clarification: May include bypass of superior vena cava syndrome, renal artery bypass, or lower extremity bypass.

- Yes, planned
 - Yes, unplanned due to surgical complication
 - Yes, unplanned due to unsuspected disease or anatomy
 - No
-
-

SEQ. #: 6540

Long Name: Other Non Card-Other Thor

Short Name: ONCOThor

Definition: Indicate whether patient underwent procedures involving Thorax/Pleura in conjunction with the primary surgical procedure. This includes but is not limited to open lung biopsy, lung resection, mediastinal mass and/or lung dissection.

Intent/Clarification: This includes, but is not limited to, lung resection, mediastinal mass and/or lung dissection. Do not code minor thoracic procedures, such as a biopsy. Only capture procedures that increase the risk of morbidity or mortality when done in conjunction with the index procedure. For procedures considered “major” in the Thoracic Database, review the data collection form:

http://www.sts.org/sites/default/files/documents/STSThoracicDCF_V2_3_MajorProc_Annotated.pdf

- Yes, planned
 - Yes, unplanned due to surgical complication
 - Yes, unplanned due to unsuspected disease or anatomy
 - No
-
-

SEQ. #: 6545

Long Name: Other Non Card-Other

Short Name: ONCOther

Definition: Indicate whether the patient had any other non-cardiac procedure performed in conjunction with the primary surgical procedure that is not included within this section.

Intent/Clarification: The goal is to keep as many procedures as possible in the “isolated” category. Only code “yes” for procedures that high likelihood of negatively impacting a patient's outcome (survival, quality of life, ability to recover) and/or prolong the patient's length of stay.

- Yes, planned
- Yes, unplanned due to surgical complication
- Yes, unplanned due to unsuspected disease or anatomy
- No

Postoperative

SEQ. #: 6550

Long Name: Postoperative Peak Glucose

Short Name: PostOpPeakGlu

Definition: Indicate the postoperative peak glucose measured within 18-24 hours of anesthesia end time.

Intent/Clarification: Hyperglycemia has been associated with increased in-hospital morbidity and mortality in patients undergoing surgery. The risk of infection was significantly higher for patients undergoing CABG if blood glucose levels were elevated. Hyperglycemia in the immediate postoperative phase increases infection in both diabetic and nondiabetic patients and the higher the level of hyperglycemia the higher the potential for infection in both populations.

Cardiac surgery patients must have controlled postoperative blood glucose (less than or equal to 180 mg/dL) in the timeframe of 18 to 24 hours after Anesthesia End Time.

(Van den Berghe, 2001) (Zerr, et al 1997) (Latham, et al, 2001)

Code the highest postoperative glucose **18 – 24 hours after Anesthesia End Time**. Can be serum or POC (point of care).

Inclusion Guidelines for Abstraction (SCIP)

- Blood glucose level
- Blood sugar
- Fasting glucose
- Finger stick glucose
- Glucometer results
- Glucose
- Non-fasting glucose
- Random glucose
- Serum glucose

SEQ. #: 6555

Long Name: Postoperative Creatinine Level

Short Name: PostCreat

Definition: Indicate the postoperative Creatinine level. If more than one level is obtained, code the highest level.

Intent/Clarification: The postoperative creatinine will be used to evaluate renal function according to the RIFLE criteria. The Acute Dialysis Quality Initiative, a multidisciplinary

collaboration, defined a range of acute renal dysfunction called the RIFLE Classification system. It is used to define grades of severity based on objective measurements. STS will use the underlined serum creatinine values to analyze post op renal function. GFR and urine output will not be included at this time. Renal Failure criteria are highlighted. Classifications of Loss and End-stage disease are beyond the current scope of follow-up.

- Risk (R) - Increase in serum creatinine level X 1.5 or decrease in GFR by 25%, or UO <0.5 mL/kg/h for 6 hours
- Injury (I) - Increase in serum creatinine level X 2.0 or decrease in GFR by 50%, or UO <0.5 mL/kg/h for 12 hours
- **Failure (F): Increase in serum creatinine level X 3.0, or serum creatinine level \geq 4.0 mg/dL; acute rise must be \geq 0.5 mg/dL or decrease in GFR by 75%;, UO , 0.3 mL/kg/hr. X 24 hours, or anuria for 12 hours.**
- Loss (L) - Persistent ARF, complete loss of kidney function >4 weeks
- End-stage kidney disease (E) - Loss of kidney function >3 months

Code the highest creatinine level from first postoperative lab to discharge.

Reference: <https://ccforum.biomedcentral.com/articles/10.1186/cc2872>

The data specifications allow only one decimal place for this value. ~~This may require standard rounding for values reported in two decimal places. For example, for creatinine reported 1.75 code 1.8; for creatinine reported as 1.52 code 1.5.~~ **Correction: DCRI has determined that you should not round the values and can only code creatinine to one decimal place.**

SEQ. #: 6556

Long Name: Postoperative Hemoglobin

Short Name: PostopHemoglobin

Definition: Indicate the postoperative hemoglobin closest to discharge

Intent/Clarification: The hemoglobin (Hgb) from the laboratory report closest to the time of discharge should be accessed first when coding this variable. If this is unavailable, then point of care testing results may be used.

SEQ. #: 6557

Long Name: Postoperative Hematocrit

Short Name: PostopHct

Definition: Indicate the postoperative hematocrit closest to discharge

Intent/Clarification: The hematocrit (Hct) from laboratory report closest to the time of discharge should be accessed first when coding this variable. If this is unavailable, then point of care testing results may be used.

SEQ. #: 6560

Long Name: Blood Prod

Short Name: BldProd

Definition: Indicate whether blood products were transfused any time postoperatively. Postoperatively is defined as any blood started after the initial surgery. Include blood transfused after the initial surgery, including any blood transfused during a re-operative surgery.

Intent/Clarification: The intent is to track postoperative blood utilization. Blood products refer to, RBC (includes whole blood), FFP, Cryoprecipitate, and Platelets.

Do NOT include:

- Pre-donated autologous blood
 - Cell saver blood
 - Pump residual blood
 - Chest tube re-circulated blood
-
-

SEQ. #: 6565

Long Name: Blood Prod - RBC Units

Short Name: BdRBCU

Definition: Indicate the number of units of packed red blood cells that were transfused any time postoperatively.

Do not include autologous, cell-saver or chest tube recirculated blood.

Intent/Clarification: The intent is to track postoperative blood utilization

SEQ. #: 6570

Long Name: Blood Prod - FFP Units

Short Name: BdFFPU

Definition: Indicate the number of units of fresh frozen plasma that were transfused any time postoperatively.

Intent/Clarification: The intent is to track postoperative blood utilization

SEQ. #: 6575

Long Name: Blood Prod - Cryo Units

Short Name: BdCryoU

Definition: Indicate the number of units of cryoprecipitate that were transfused postoperatively. One bag of cryo = one unit.
The number of units is not volume dependent.

Intent/Clarification: The intent is to track postoperative blood utilization

SEQ. #: 6580

Long Name: Blood Prod - Platelet Units

Short Name: BdPlatU

Definition: Indicate the number of units of platelets that were transfused postoperatively. Count the dose pack as one unit. A dose pack may consist of 4, 6, 8, 10, or any number of donor platelets obtained. The number of units coded is not volume dependent.

Intent/Clarification: The intent is to track postoperative blood utilization

SEQ. #: 6585

Long Name: Extubated In OR

Short Name: ExtubOR

Definition: Indicate whether the patient was extubated prior to leaving the operating room during the initial surgery.
If patient expires in the operating room during the initial surgery, answer "Yes".

Intent/Clarification:

- Yes: if the patient is extubated in the OR during the initial surgery
- No: if patient extubated after leaving the operating room

FAQ October 2017: If a patient is extubated in the OR, will this patient be counted in the <6 hours metric.

Answer: Yes, they will now be included in the <6 metric and their vent times will calculate to zero hours by DCRI.

SEQ. #: 6591

Long Name: Postop Intubation/Reintubation During Hospital Stay

Short Name: PostopIntub

Definition: Indicate whether the patient was intubated for the first time after leaving the OR from the initial procedure, or re-intubated during the hospital stay after the initial extubation.

Intent/Clarification: Do not include reintubation for surgical procedures when the patient is extubated prior to leaving the operating room.

SEQ. #: 6595

Long Name: Additional Hours Ventilated

Short Name: VentHrsA

Definition: Indicate how many additional hours the patient was on ventilator after initial extubation.

Intent/Clarification: If the patient was reintubated during the current hospital stay, this value is used in the calculation to determine prolonged ventilation.

Ventilator hours are calculated with a decimal point so that minutes can be included. Divide the number of minutes by 60.

Examples:

0.1 = 6 minutes

0.3 = 15 minutes

0.5 = 30 minutes

0.8 = 45 minutes etc.

SEQ. #: 6600

Long Name: Total Postoperative Ventilation Hours

Short Name: VentHrsTot

Definition: Calculated variable measuring OR exit time to extubation time plus any additional hours due to reintubation.

Intent/Clarification: This will be system calculated by the software by adding initial post-op vent hours plus additional postop vent hours to determine total post op vent time. Anything greater than 24 hours is considered prolonged postop vent time. Total hours ventilated is rounded in the calculation.

SEQ. #: 6605

Long Name: ICU Visit

Short Name: ICUVisit

Definition: Indicate whether the patient received ICU level of care immediately following the initial surgery. Include ICU unit, post-anesthesia recovery, and other similar critical care environments.

Intent/Clarification: Indicate whether the patient received ICU level of care immediately following the initial surgery. Include ICU units and other similar critical care environments. Do not include PACU if only used for Phase I recovery, but do include PACU if used as a critical care unit when an ICU bed was not available.

SEQ. #: 6610

Long Name: Initial ICU hours **Short Name:** ICUInHrs

Definition: Indicate the number of hours the patient received ICU level of care immediately following the initial surgery until the time of actual transfer out of ICU. Include ICU unit, post-anesthesia recovery, and other similar critical care environments. For those sites providing postop ICU level of care in one single stay unit (admission to ICU to hospital discharge), document the number of hours immediately following the initial surgery until a physician order is written to change the level of care provided.

Intent/Clarification: ICU hours begin when the patient arrives in the ICU or your institutions equivalent to an ICU and ends when they physically leave the ICU. For those sites with single stay units (admission to ICU to hospital discharge), document the number of hours immediately following the initial surgery until a *physician order is written* to change the level of care provided. If the patient expires, use the date/time on the death certificate (time pronounced dead).

Time frame is OR Exit Date and Time until the patient leaves the ICU.

The only way to objectively count ICU time is to count the actual time the patient physically leaves the ICU. Using the time of transfer orders misrepresents actual ICU time.

SEQ. #: 6615

Long Name: Readmission to ICU

Short Name: ICUReadm

Definition: Indicate whether the patient spent time in an ICU after having been transferred to a step-down unit (lower level care). Specific situations are described below:

- OR → ICU → OR → ICU = No
- OR → ICU → STEP DOWN →→ ICU = Yes
- OR → STEP DOWN → ICU = Yes

Single care unit:

Code ICU readmission when the level of care increases and is noted in the physician order.

Intent/Clarification: The intent is to capture episodes of patient deterioration necessitating a higher level of care. For single stay units, this is indicated by a physician order.

SEQ. #: 6620

Long Name: Additional ICU Hours

Short Name: ICUAdHrs

Definition: Indicate the number of additional hours spent in the ICU, or at the equivalent higher level of care in single stay units.

Intent/Clarification: This will be used, along with initial ICU hours, to determine total post op ICU hours, an indication of resource utilization.

For single stay units, time should be calculated by the time stamp on a physician's order to elevate the patient's level of care to Intensive Care until the level of care is deescalated. If the patient expires, time should be counted from order to elevate the level of care to the time of death as noted in the medical record.

SEQ. #: 6625

Long Name: Postop Echo

Short Name: POpTTEch

Definition: Indicate whether an echo was performed postoperatively to evaluate valvular function prior to discharge.

Intent/Clarification: Capture echocardiograms performed after the patient leaves the operating room but prior to hospital discharge. Code the exam closest to discharge.

SEQ. #: 6630

Long Name: Postop Echo Aortic Insufficiency

Short Name: POpTTAR

Definition: Indicate the level of aortic insufficiency/regurgitation found on post op echo closest to discharge. Mild-to-moderate should be coded as moderate; moderate to severe should be coded as severe.

Intent/Clarification: Capture echocardiograms performed after the patient leaves the operating room but prior to hospital discharge. If the report for an echocardiogram does not address valve disease, code "not reported". Use the following to categorize the level of insufficiency/regurgitation:

- None
- Trace/trivial
- Mild
- Moderate
- Severe

- Not Reported

SEQ. #: 6631

Long Name: Postop Echo Aortic Paravalvular Leak

Short Name: POpAortParaLk

Definition: Indicate the level of aortic paravalvular leak found on post op echo closest to discharge. Mild-to-moderate should be coded as moderate; moderate to severe should be coded as severe.

Intent/Clarification: To identify a paravalvular leak in a prosthetic aortic valve.

SEQ. #: 6635

Long Name: Postop Echo Mitral Insufficiency

Short Name: POpTTMR

Definition: Indicate the level of mitral insufficiency/regurgitation found on post op echo closest to discharge. Mild-to-moderate should be coded as moderate; moderate to severe should be coded as severe.

Intent/Clarification: Capture echocardiograms performed after the patient leaves the operating room but prior to hospital discharge. If the report for an echo does not address valve disease, code “not reported”. Use the following to categorize the level of insufficiency/regurgitation:

- None
 - Trace/trivial
 - Mild
 - Moderate
 - Severe
 - Not Reported
-

SEQ. #: 6636

Long Name: Postop Echo Mitral Paravalvular leak

Short Name: POpMitParaLk

Definition: Indicate the level of mitral paravalvular leak found on post op echo closest to discharge. Mild-to-moderate should be coded as moderate; moderate to severe should be coded as severe.

Intent/Clarification: To identify a paravalvular leak in a prosthetic mitral valve.

SEQ. #: 6640

Long Name: Postop Echo Tricuspid Insufficiency

Short Name: POpTTTR

Definition: Indicate the level of tricuspid insufficiency/ regurgitation found on post op echo closest to discharge. Mild-to-moderate should be coded as moderate; moderate to severe should be coded as severe.

Intent/Clarification: Capture echocardiograms performed after the patient leaves the operating room but prior to hospital discharge. If the report for an echo does not address valve disease, code “not reported”. Use the following to categorize the level of insufficiency/regurgitation:

- None
 - Trace/trivial
 - Mild
 - Moderate
 - Severe
 - Not Reported
-
-

SEQ. #: 6645

Long Name: Postop Echo Pulmonic Insufficiency

Short Name: POpTTPu

Definition: Indicate the level of pulmonic insufficiency/ regurgitation found on post op echo closest to discharge. Mild-to-moderate should be coded as moderate; moderate to severe should be coded as severe.

Intent/Clarification: Capture echocardiograms performed after the patient leaves the operating room but prior to hospital discharge. If the report for an echo does not address valve disease, code “not reported”. Use the following to categorize the level of insufficiency/regurgitation:

- None
 - Trace/trivial
 - Mild
 - Moderate
 - Severe
 - Not Reported
-
-

SEQ. #: 6650

Long Name: Postop EF Done

Short Name: POpEFD

Definition: Indicate whether the Ejection Fraction was measured postoperatively.

Intent/Clarification:

- Yes

- No

SEQ. #: 6655

Long Name: Postop EF

Short Name: POpEF

Definition: Indicate the percentage of the blood emptied from the left ventricle at the end of the contraction measured postoperatively.

Intent/Clarification: Enter a range of 1-99. If a percentage range is reported, report a whole number using the “mean” (i.e., 50- 55% is reported as 53%). If a qualitative description is reported, code the mean value for that range; i.e., normal (50-70%) is coded as 60%.

- Hyperdynamic: >70% (code 71%)
- Normal: 50%–70% (midpoint 60%)
- Mild dysfunction: 40%–49% (midpoint 45%)
- Moderate dysfunction: 30%–39% (midpoint 35%)
- Severe dysfunction: <30% (code 29%)

Note: If no diagnostic report is in the medical record, a value documented in the medical record is acceptable.

SEQ. #: 6660

Long Name: Postop Cardiac Enzymes Drawn

Short Name: POpEnzDrawn

Definition: Indicate whether Cardiac Enzymes (biomarkers) were drawn post procedure.

Intent/Clarification: Capture cardiac enzymes that were drawn after surgery, prior to discharge. This does not imply that enzymes should be drawn on all patients; the intent is to capture the values if they were drawn. Include one-time draws if serial enzymes were not drawn.

SEQ. #: 6665

Long Name: Postop Peak CKMB

Short Name: POpPkCKMB

Definition: Indicate the peak CKMB (highest level post procedure).

Intent/Clarification: CKMB is the fraction of the enzyme directly related to myocardial tissue. Record the highest level post procedure if multiple were drawn.

SEQ. #: 6670

Long Name: Postop Peak Troponin I

Short Name: POpPkTrI

Definition: Indicate the peak Troponin I (highest level post procedure).

Intent/Clarification: Troponin I is a very sensitive and specific indicator of damage to the heart muscle (myocardium). It is used in conjunction with other diagnostic criteria to diagnose myocardial infarction. Record the highest level if multiple were drawn.

SEQ. #: 6675

Long Name: Postop Peak Troponin T

Short Name: POpPkTrT

Definition: Indicate the peak Troponin T (highest level post procedure).

Intent/Clarification: Troponin T is a very sensitive and specific indicator of damage to the heart muscle (myocardium). It is used in conjunction with other diagnostic criteria to diagnose myocardial infarction. Record the highest level if multiple were drawn.

SEQ. #: 6680

Long Name: Postop 12 Lead EKG

Short Name: POpEKG

Definition: Indicate the post procedure 12 lead EKG findings, if performed.

Intent/Clarification: This does not imply 12 leads are standard procedures for all post op patients. If more than one 12 lead EKG is done following surgery, capture the last one done prior to discharge.

- Not Performed
 - No ischemic changes
 - New ST changes (does not include LBBB or ST elevation)
 - New Pathologic Q Wave or LBBB
 - New STEMI
 - Other
 - N/A (No pre-op EKG for comparison, transplant) Arrhythmias are not captured here.
- -----

Postoperative Events

SEQ. #: 6690

Long Name: Post-Op-Surgical Site Infection

Short Name: SurSInf

Definition: Indicate whether a surgical site infection (SSI) was diagnosed within 30 days of the procedure or any time during the hospitalization for surgery. Refer to the most current CDC definition for SSI which can be found in the training manual.

Intent/Clarification:

- **Superficial Incisional SSI:** Must meet the following criteria:
 - Infection occurs \leq 30 days, **and** involves only skin/subcutaneous tissue of the incision, **and** patient has \geq one of the following:
 - Purulent drainage from the superficial incision.
 - Organisms isolated from an aseptically-obtained culture of fluid or tissue from the superficial incision.
 - Superficial incision that is deliberately opened by a surgeon, attending physician or other designee and is culture positive or not cultured **and** patient has \geq one of the following:
 - pain or tenderness
 - localized swelling
 - redness
 - heat
 - A culture with negative findings does not meet this criterion.
 - Diagnosis of a superficial incisional SSI by the surgeon or attending physician or other designee.
 - There are two specific types of superficial incisional SSIs:
 - Superficial Incisional Primary (SIP) – a superficial incisional SSI that is identified in the primary incision in a patient that has had an operation with one or more incisions (chest incision for CABG)
 - Superficial Incisional Secondary (SIS) – a superficial incisional SSI that is identified in the secondary incision in a patient that has had an operation with more than one incision (e.g., donor site incision for CABG)

Do not include:

- A stitch abscess alone (minimal inflammation and discharge confined to the points of suture penetration)
- A localized stab wound or pin site infection.
- Diagnosis of “cellulitis” by itself
- **Deep incisional SSI:** Must meet the following criteria
 - Infection occurs within 30 days after the operative procedure, **and** involves deep soft tissues of the incision (e.g., fascial and muscle layers) **and** patient has at least one of the following:
 - Purulent drainage from the deep incision.
 - A deep incision that spontaneously dehisces or is deliberately opened by a surgeon, attending physician or other designee and is culture-positive or not cultured, **and** patient has at least one of the following signs or symptoms:
 - Fever ($>38^{\circ}\text{C}$)
 - Localized pain or tenderness
 - An abscess or other evidence of infection involving the deep incision that is detected on direct examination, during invasive procedure, or by histopathologic examination or imaging test.
 - A culture with negative findings does not meet this criterion.
 - There are two specific types of deep incisional SSIs:

- Deep Incisional Primary (DIP) – a deep incisional SSI that is identified in a primary incision in a patient that has had an operation with one or more incisions (e.g., chest incision for CABG)
- Deep Incisional Secondary (DIS) – a deep incisional SSI that is identified in the secondary incision in a patient that has had an operation with more than one incision (e.g., donor site incision for CABG)
- **Organ/Space SSI:** Must meet the following criteria
 - Infection occurs within 30 days after the operative procedure, and infection involves any part of the body, excluding the skin incision, fascia, or muscle layers, that is opened or manipulated during the operative procedure, **and** patient has at least one of the following:
 - Purulent drainage from a drain that is placed into the organ/space
 - Organisms isolated from an aseptically-obtained culture of fluid or tissue in the organ/space
 - An abscess or other evidence of infection involving the organ/space that is detected on direct examination, during invasive procedure, or by histopathologic examination or imaging test, **and** meets at least one criterion for a specific organ/space infection of mediastinitis below:
- **MED-Mediastinitis:** Must meet the following criteria
 - Mediastinitis must meet at least 1 of the following criteria:
 - Patient has organisms cultured from mediastinal tissue or fluid obtained during an invasive procedure.
 - Patient has evidence of mediastinitis seen during an invasive procedure or histopathologic examination.
 - Patient has **at least 1** of the following signs or symptoms:
 - Fever (>38°C)
 - Chest pain*
 - Sternal instability*
 - and at least 1** of the following:
 - Purulent discharge from mediastinal area
 - Organisms cultured from blood or discharge from mediastinal area
 - Mediastinal widening on imaging test.

* With no other recognized cause

Report mediastinitis following cardiac surgery that is accompanied by osteomyelitis as SSI-MED rather than SSI-BONE

SEQ. #: 6695

Long Name: Post-Op-Sternal-Superficial Wound Infection

Short Name: CSternalSupInf

Definition: Indicate whether a superficial sternal wound infection was diagnosed within 30 days of the procedure or any time during the hospitalization for surgery.

Intent/Clarification: See above definition for superficial site infection.

SEQ. #: 6700

Long Name: Post-Op-Deep Sternal Infection / Mediastinitis

Short Name: DeepSternInf

Definition: Indicate whether a deep sternal wound infection or mediastinitis was diagnosed within 30 days of the procedure or any time during the hospitalization for surgery.

Intent/Clarification: See above definition for Deep Sternal Infection/Mediastinitis. The STS Composite scores weighs deep sternal wound infection and mediastinitis the same.

SEQ. #: 6705

Long Name: Post-Op-Deep Sternal Infection / Mediastinitis - Date

Short Name: DeepSternInfDt

Definition: Indicate the first date that deep sternal wound infection or mediastinitis was documented.

Intent/Clarification: Required date format: mm/dd/yyyy

SEQ. #: 6710

Long Name: Post-Op-Infect-Thoracotomy

Short Name: CITHor

Definition: Indicate whether a surgical site infection involving a thoracotomy or parasternal site was diagnosed within 30 days of the procedure or any time during the hospitalization for surgery.

Intent/Clarification: Time frame is from OR Exit time to 30 days post procedure or discharge from initial hospital visit if admitted for greater than 30 days.

SEQ. #: 6715

Long Name: Post-Op-Conduit Harvest

Short Name: ConduitHarv

Definition: Indicate whether a surgical site infection involving a conduit harvest site was diagnosed within 30 days of the procedure or any time during the hospitalization for surgery.

Intent/Clarification: Capture infections at the site of an endovascular harvest site or an open harvest site, arm or leg.

Time frame is from OR Exit time to 30 days post hospital visit if admitted for greater than 30 days.

SEQ. #: 6720

Long Name: Post-Op-Cannulation Site

Short Name: CanSite

Definition: Indicate whether a surgical site infection involving a cannulation site was diagnosed within 30 days of the procedure or any time during the hospitalization for surgery.

Intent/Clarification:

Capture infections of cannulation sites. These are considered secondary surgical site infections since they do not involve the primary surgical incision. Follow CDC criteria above.

Time frame is from OR Exit time to 30 days post procedure or discharge from initial hospital visit if admitted for greater than 30 days.

SEQ. #: 6725

Long Name: Post-Op-Wound Intervention / Procedure

Short Name: WoundInter

Definition: Indicate whether a wound intervention or procedure was performed.

Intent/Clarification: The intent is to capture treatment strategies employed to treat the surgical site infection(s). Indicate below whether treatment was applied to the primary incision, secondary incision or both.

SEQ. #: 6730

Long Name: Post-Op-Wound Intervention - Open With Packing / Irrigation

Short Name: WoundIntOpen

Definition: Indicate whether wound intervention(s) involved opening the wound and packing and/or irrigation.

Intent/Clarification: The intent is to capture treatment strategies employed to treat the surgical site infection within 30 days following procedure included leaving the incision open with packing/irrigation.

SEQ. #: 6735

Long Name: Post-Op-Wound Intervention - Wound Vac

Short Name: WoundIntVac

Definition: Indicate whether wound intervention(s) included application of a wound vac.

Intent/Clarification: A wound-vac may also be called negative pressure wound therapy. A wound-vac is a device is used to facilitate wound healing by converting an open wound to a closed wound. The application of negative pressure causes removal of excess fluids, increased blood flow and decreased bacterial colonization; promotes granulation tissue formation and wound closure.

SEQ. #: 6740

Long Name: Post-Op-Wound Intervention - Secondary Procedure Muscle Flap

Short Name: WoundIntMuscle

Definition: Indicate whether wound intervention(s) included a secondary procedure involving a muscle flap.

Intent/Clarification: Refer to operative dictations. Muscle flaps are typically performed by a CV Surgeon or Plastic Surgeon.

SEQ. #: 6745

Long Name: Post-Op-Wound Intervention - Secondary Procedure Omental Flap

Short Name: WoundIntOmental

Definition: Indicate whether wound intervention(s) included a secondary procedure involving an Omental flap.

Intent/Clarification: Refer to operative dictations.

SEQ. #: 6750

Long Name: In Hospital Post-Op Events

Short Name: Complics

Definition: Indicate whether a postoperative event occurred during the hospitalization for surgery. This includes the entire postoperative period up to discharge, even if over 30 days.

Intent/Clarification:

The intent is to document those events/complications that:

- Pose either a life threatening situation or create a potential long-term deficit
- Require pharmacological, surgical or medical intervention to prevent further clinical deterioration
- Increase length of stay and/or resource utilization.

If the patient expires in the operative room, the complications section does not need to be completed. There would not have been a post-operative period for the patient, therefore, no post-operative complications. Code the Complications data fields "No".

SEQ. #: 6755

Long Name: Post-Op-ReOp Bleed

Short Name: COpReBld

Definition: Indicate whether the patient was re-explored for mediastinal bleeding with or without tamponade either in the ICU or returned to the operating room.

Intent/Clarification: Do not capture reopening of the chest or situations of excessive bleeding that occur prior to the patient leaving the operating room at the time of the primary procedure. Tamponade is a situation which occurs when there is compression or restriction placed on the heart within the chest that creates hemodynamic instability or a hypoperfused state. Do not include medically (non-operatively) treated excessive post-operative bleeding/tamponade events.

Include patients that return to an OR suite or equivalent OR environment (i.e., ICU setting) as identified by your institution, that require surgical re-intervention to investigate/correct bleeding with or without tamponade. Include only those interventions that pertain to the mediastinum or thoracic cavity.

SEQ. #: 6760

Long Name: Post-Op-ReOp Bleed Timing

Short Name: COpReBldTim

Definition: Indicate when reoperation for bleeding took place.

Intent/Clarification:

- Acute* - Within 24 hours of the end of the case
- Late - more than 24 hours after case ends

*Code exactly 24 hours as Acute

SEQ. #: 6765

Long Name: Post-Op-ReOp Vlv Dys

Short Name: COpReVlv

Definition: Indicate whether the patient returned to the operating room for prosthetic or native valve dysfunction. Dysfunction may be structural and/or non-structural failure. Dysfunction may be of prosthesis, a progressive native disease process, or an acute event process that disrupts valve function and creates either clinical compromising insufficiency/regurgitation or valve orifice narrowing.

Intent/Clarification:

- Yes, surgical
 - Yes, transcatheter
 - No
-
-

SEQ. #: 6771**Long Name:** Post-Op-Reintervention-Myocardial Ischemia**Short Name:** CReintMI**Definition:** Indicate whether the patient required postoperative reintervention for Myocardial Ischemia.**Intent/Clarification:**

Only capture surgical or Cath lab interventions that occur during the hospitalization prior to discharge.

- Yes, surgical
- Yes, PCI
- No

FAQ September 2017: Patient was 4 days post op CAB developed chest pain with elevated troponin. Patient was taken to the Cath lab and found to have a freshly occluded graft. After opening the graft with "mechanical thrombolysis" the graft was open, but noticed an AV fistula between the graft and the LV. The physician then ballooned the area to re-occlude the graft and seal the fistula. This was successful. How does this get captured?

Answer: Code Yes, PCI.

SEQ. #: 6772**Long Name:** Post-Op-Reintervention-Myocardial Ischemia-Vessel**Short Name:** CReintMIVes**Definition:** Indicate the type of vessels that required postoperative reintervention for Myocardial Ischemia.**Intent/Clarification:** Reintervention may involve native coronary arteries, coronary artery bypass grafts or both.

- Native Coronary
- Graft
- Both

FAQ September 2017: Patient was 4 days post op CAB developed chest pain with elevated troponin. Patient was taken to the Cath lab and found to have a freshly occluded graft. After opening the graft with "mechanical thrombolysis" the graft was open, but noticed an AV fistula between the graft and the LV. The physician then

ballooned the area to re-occlude the graft and seal the fistula. This was successful. How does this get captured?

Answer: Code Graft.

SEQ. #: 6773

Long Name: Post-Op-Reintervention-Myocardial Ischemia - Intervention Type

Short Name: CReintMIIntTy

Definition: Indicate the type of intervention used postoperatively for Myocardial Ischemia.

Intent/Clarification: Reintervention may include surgery, PCI or both.

FAQ September 2017: Patient was 4 days post op CAB developed chest pain with elevated troponin. Patient was taken to the Cath lab and found to have a freshly occluded graft. After opening the graft with "mechanical thrombolysis" the graft was open, but noticed an AV fistula between the graft and the LV. The physician then ballooned the area to re-occlude the graft and seal the fistula. This was successful. How does this get captured?

Answer: Code PCI.

SEQ. #: 6774

Long Name: Post-Op-Aortic Reintervention

Short Name: CAortReint

Definition: Indicate whether the patient underwent postoperative aortic reintervention.

Intent/Clarification:

- Yes
- No

FAQ October 2017: The patient was initially scheduled for TEVAR but after a series of CT scans and manifestations of signs and symptoms, the surgeon opted to repair the ascending aortic flap first and then, 3 days later, he took the patient back to the OR for the TEVAR. Should this be captured as post-operative event aortic reintervention?

Answer: No, code as a planned staged hybrid. Then capture the TEVAR accordingly in Seq. No. 5095.

SEQ. #: 6775

Long Name: Post-Op-Aortic Reintervention-Type

Short Name: CAortReintTy

Definition: Indicate the type of aortic intervention the patient received.

Intent/Clarification: Reintervention may be open or endovascular.

SEQ. #: 6778

Long Name: Post-Op-ReOp Other Card

Short Name: COpReOth

Definition: Indicate whether the patient returned to the operating room for other cardiac reasons.

Intent/Clarification: Capture any other cardiac reasons for reoperation.

- Yes
 - No
-
-

SEQ. #: 6780

Long Name: Post-Op-Return To OR For Other Non-cardiac Reason

Short Name: COpReNon

Definition: Indicate whether the patient returned to the operating room for other non-cardiac reasons.

This includes procedures requiring a return to the operating room such as tracheostomy, general surgery procedures.

This does not include procedures performed outside the operating room such as GI Lab for peg tube, shunts for dialysis, etc.

Intent/Clarification:

Non-cardiac events include, but are not limited to, events as described in Section N.

Code only those non-cardiac events that require a return to the surgical suite.

This includes procedures requiring a return to the operating room, such as a tracheostomy, hematoma evacuation, etc.

This does not include procedures performed outside the operating room, such as GI lab for peg tubes, shunts for dialysis, etc. Due to practice pattern(s) determined by institutional culture or practice driven patterns, some sites may have included in this section cases and/or events that other sites may not. Capture those events that may pose a clinically or resource utilization impact on the patient AND necessitate a return to the OR.

For planned procedures, (i.e. a patient who is scheduled for lower extremity vascular surgery requiring a CAB prior to the scheduled vascular procedure), Code "No," as this is not a complication, coding it as a complication misrepresents the outcome of the surgery.

Events captured here are not included in the reoperative measure of the composite score.

FAQ September 2017: Should we code a GI event for an abdominal compartment syndrome leading to a decompressive laparotomy?

Answer: Yes, code GI event for abdominal compartment syndrome in addition to the reoperation other non-cardiac for the decompression laparotomy.

FAQ October 2017: If the patient has a percutaneous tracheostomy done at the bedside, is this considered a reoperation other non-cardiac.

Answer: Yes, code 'Reoperation, other non-cardiac' for all tracheostomies regardless of the location where the tracheostomy was performed.

SEQ. #: 6785

Long Name: Post-Op-Open Chest With Planned Delayed Sternal Closure

Short Name: COpPIndDelay

Definition: Indicate whether the chest was left open with planned delayed sternal closure.

Intent/Clarification: This allows capture of patients who have the chest left open with a planned delayed sternal closure.

SEQ. #: 6790

Long Name: Post-Op-Sternotomy Issue

Short Name: CSternal

Definition: Indicate presence of a post-operative sternotomy issue.

Intent/Clarification: Indicate presence of a post-operative sternotomy issue prior to discharge. Any condition requiring operative intervention involving the sternotomy should be coded YES.

SEQ. #: 6795

Long Name: Post-Op Sternal instability/dehiscence (sterile)

Short Name: CSternalDehis

Definition: The code indicates sterile dehiscence of the sternal edges without evidence of infection but which requires surgical intervention. Skin and subcutaneous tissue may remain intact.

Intent/Clarification: Wound dehiscence (sterile) is defined as separation of the layers of a surgical wound. This separation can either be superficial or deep and can include the sternum in the case of a median sternotomy incision. The code "Sternal instability (sterile)" should be used to record the complication when the superficial and deep layers of the incision remain intact but non-union of the sternal edges is present. Causes of

wound dehiscence can include tissue ischemia, nutritional deficiencies, use of corticosteroids, vitamin C deficiency, and others.

Wound dehiscence due to wound infection should be recorded as a wound infection.

SEQ. #: 6800

Long Name: Post-Op-Sepsis

Short Name: CSepsis

Definition: Sepsis is defined as evidence of serious infection accompanied by a deleterious systemic response. In the time period of the first 48 postoperative or post procedural hours, the diagnosis of sepsis requires the presence of a Systemic Inflammatory Response Syndrome (SIRS) resulting from a proven infection (such as bacteremia, fungemia or urinary tract infection). In the time period after the first 48 postoperative or post procedural hours, sepsis may be diagnosed by the presence of a SIRS resulting from suspected or proven infection. During the first 48 hours, a SIRS may result from the stress associated with surgery and/or cardiopulmonary bypass. Thus, the clinical criteria for sepsis during this time period should be more stringent.

A systemic inflammatory response syndrome (SIRS) is present when at least two of the following criteria are present: hypo- or hyperthermia (>38.5 or <36.0), tachycardia or bradycardia, tachypnea, leukocytosis or leukopenia, or thrombocytopenia.

Intent/Clarification: Indicate whether sepsis was diagnosed during initial hospitalization. Sepsis is defined as having 2 or more of the SIRS (systemic inflammatory response syndrome) criteria and a known or suspected infection, typically occurring within 6 hours of each other. SIRS may occur unrelated to infection, as in the case of cardiac surgery, and not indicative of Sepsis.

- Within the first 48 hours of cardiac surgery a patient **MUST** meet 2 SIRS criteria typically within 6 hours of each other and have a **PROVEN** infection (not suspected). Clinical criteria of sepsis must be more stringent during the first 48 hours following surgery because the stress surgery produces results in SIRS criteria and is not typically related to an infection.
- After 48 hours, the patient must have 2 or more SIRS criteria typically within 6 hours of each other and a known or suspected infection.

SIRS criteria included:

- HR > 90 (acute and not a chronic condition)
 - Temp >38.5 <36.0
 - Resp >20 bpm or PaCO₂ <32 mmHg
 - WBC <4000 or >12000 or >10% Bands
-
-

SEQ. #: 6805

Long Name: Post-Op-Sepsis-Positive Blood Cultures

Short Name: CSepsisPBC

Definition: Indicate whether a recognized pathogen is cultured from 1 or more blood cultures and is not related to an infection at another site.

Intent/Clarification: Indicate if blood cultures obtained in the post-operative period were positive for infectious pathogen. Staph epidermis (*S. epidermis*) is a common contaminate of blood cultures. Refer to the ordering physician's interpretation of results if there is a questions.

SEQ. #: 6810

Long Name: Post-Op-Neuro-Stroke Perm

Short Name: CNStrokP

Definition: Indicate whether the patient has a postoperative stroke and the type of stroke (i.e., any confirmed neurological deficit of abrupt onset caused by a disturbance in blood supply to the brain) that did not resolve within 24 hours.

Intent/Clarification: Stroke occurs when the blood supply to part of the brain is suddenly interrupted or when a blood vessel in the brain bursts, spilling blood into the spaces surrounding brain cells. Brain cells die when they no longer receive oxygen and nutrients from the blood or there is sudden bleeding into or around the brain.

The symptoms of a stroke include:

- Sudden numbness or weakness, especially on one side of the body
- Sudden confusion or trouble speaking or understanding speech
- Sudden trouble seeing in one or both eyes
- Sudden trouble with walking, dizziness, or loss of balance or coordination
- Sudden severe headache with no known cause

There are two forms of stroke:

- Ischemic - blockage of a blood vessel supplying the brain
- Hemorrhagic - bleeding into or around the brain

Central events are caused by embolic or hemorrhagic events. Neurological deficits such as confusion, delirium and/or encephalopathic (anoxic or metabolic) events are not to be coded in this field.

Reference: <https://www.ninds.nih.gov/Disorders/All-Disorders/Stroke-Information-Page>

FAQ October 2017: How should embolic stroke be coded?

Answer: Embolic strokes should be coded as ischemic.

SEQ. #: 6815

Long Name: Post-Op-Neuro-Transient Ischemic Attack - TIA

Short Name: CNStrokTTIA

Definition: Indicate whether the patient had a postoperative Transient Ischemic Attack (TIA): Loss of neurological function that was abrupt in onset but with complete return of function within 24 hours.

Intent/Clarification: A transient ischemic attack (TIA) is a transient stroke that lasts up to 24 hours. It occurs when the blood supply to part of the brain is briefly interrupted. TIA symptoms, which usually occur suddenly, are similar to those of stroke but do not last as long. Most symptoms of a TIA disappear within an hour, although they may persist for up to 24 hours.

Symptoms can include:

- Numbness or weakness in the face, arm, or leg, especially on one side of the body
- Confusion or difficulty in talking or understanding speech
- Trouble seeing in one or both eyes; and difficulty with walking, dizziness, or loss of balance and coordination.

Patients who have suffered a TIA have an increased risk of peripheral and coronary artery atherosclerosis, and an increased risk of subsequent heart attack and stroke.

Reference: <https://www.ninds.nih.gov/Disorders/All-Disorders/Stroke-Information-Page>

SEQ. #: 6821

Long Name: Post-Op-Neuro-Encephalopathy

Short Name: CNEnceph

Definition: Indicate the type of postoperative encephalopathy the patient developed, if any.

Intent/Clarification:

Encephalopathy is a term for any diffuse disease of the brain that alters brain function or structure. The hallmark of encephalopathy is an altered mental state. Blood tests, spinal fluid examination, imaging studies, electroencephalograms, and similar diagnostic studies may be used to differentiate the various causes of encephalopathy.

Encephalopathy may be caused by:

- Infectious agent (bacteria, virus, or prion),
- Metabolic or mitochondrial dysfunction,
- Brain tumor or increased pressure in the skull,
- Prolonged exposure to toxic elements (including solvents, drugs, radiation, paints, industrial chemicals, and certain metals),
- Chronic progressive trauma,
- Poor nutrition,
- Lack of oxygen or blood flow to the brain

Depending on the type and severity of encephalopathy, common neurological symptoms are:

- progressive loss of memory and cognitive ability
- subtle personality changes
- inability to concentrate
- lethargy
- progressive loss of consciousness.

Other neurological symptoms may include:

- myoclonus (involuntary twitching of a muscle or group of muscles)
- nystagmus (rapid, involuntary eye movement)
- tremor
- muscle atrophy and weakness
- dementia
- seizures
- loss of ability to swallow or speak

If multiple causes, choose first event.

Reference: <https://www.ninds.nih.gov/Disorders/All-Disorders/Stroke-Information-Page>

- None
- Anoxic
- Embolic
- Drug
- Metabolic
- Intracranial Bleeding
- Other
- Unknown

SEQ. #: 6822

Long Name: Post-Op-Neuro-Coma/Unresponsive State

Short Name: CNComa

Definition: Indicate whether the patient developed a postoperative coma or unresponsive state (not stroke).

Intent/Clarification: A coma, sometimes also called persistent vegetative state, is a profound or deep state of unconsciousness. Persistent vegetative state is not brain-death. An individual in a state of coma is alive but unable to move or respond to his or her environment.

SEQ. #: 6825

Long Name: Post-Op-Neuro-Paralysis

Short Name: CNParal

Definition: Indicate whether the patient had a new postoperative paralysis, paraparesis, or paraplegia related to spinal cord ischemia and not related to a stroke.

Intent/Clarification: Paralysis is a loss of purposeful movement as a result of a neurological injury, drugs or toxins. Loss of motor function may be complete (paralysis) or partial (paresis); unilateral (hemiplegic) or bilateral confined to the lower extremities (paraplegic) or present in all four extremities (quadriplegic); and may be accompanied by increased muscular tension and hyperactive reflexes (spastic) or by loss of reflexes (flaccid). Related to spinal cord ischemia, **not related to stroke**.

SEQ. #: 6826

Long Name: Post-Op-Neuro-Paralysis Type

Short Name: CNParalTy

Definition: Indicate whether the new postoperative paralysis, paraparesis, or paraplegia was transient or permanent.

Intent/Clarification: Related to spinal cord ischemia, **not related to stroke**.

- **Transient** - is non-lasting and of short (< 24 hours) duration.
 - **Permanent** - is enduring, lasting, or without change for more than 24 hours.
-
-

SEQ. #: 6829

Long Name: Post-Op-Neuro-Paresis

Short Name: CNParesis

Definition: Indicate whether postoperative paresis was present

Intent/Clarification:

- Yes
 - No
-
-

SEQ. #: 6830

Long Name: Post-Op-Neuro-Paresis Type

Short Name: CNParesisTy

Definition: Indicate the type of post op paresis

Intent/Clarification: Paresis may be transient or permanent and is related to spinal cord ischemia and not to stroke.

SEQ. #: 6832

Long Name: Post-Op-Phrenic Nerve Injury

Short Name: PhrenNrvInj

Definition: Indicate whether patient has symptoms of phrenic nerve injury, (e.g., immobility or elevation of the diaphragm, etc.).

Intent/Clarification: Traumatic or thermal injury to the phrenic nerve can result in paralysis of the hemi diaphragm on the affected side, resulting in respiratory difficulty.

SEQ. #: 6833

Long Name: Post-Op-Recurrent Laryngeal Nerve Injury

Short Name: RecLarynNrvInj

Definition: Indicate whether patient has symptoms of recurrent laryngeal nerve injury, (e.g., hoarseness, difficulty speaking, etc.).

Intent/Clarification: The recurrent laryngeal nerve controls movement of the larynx. The larynx contains the apparatus for voice production: the vocal cords, and the muscles and ligaments that move the vocal cords. It also controls the flow of air into the lungs. When the recurrent laryngeal nerve is damaged, the movements of the larynx are reduced. This causes voice weakness, hoarseness, or sometimes the complete loss of voice. The changes may be temporary or permanent.

SEQ. #: 6835

Long Name: Post-Op-Pulm-Vent Prolonged

Short Name: CPVntLng

Definition: Indicate whether the patient had prolonged post-operative pulmonary ventilation > 24.0 hours.

The hours of postoperative ventilation time include OR exit until extubation, plus any additional hours following reintubation.

Include (but not limited to) causes such as ARDS, pulmonary edema, and/or any patient requiring mechanical ventilation > 24 hours postoperatively.

Intent/Clarification: To calculate total hours, include initial and additional hours of mechanical ventilation. Extended ventilation may include, but is not limited to, the specific definitional reasons. Example: If a major stroke or coma occurred that required ventilation for life support, code as prolonged if greater than 24 hours. Do not include the hours ventilated if a patient returns to the operating room suite and requires re-intubation as part of general anesthesia but does not require ventilation beyond the time in the operating room (i.e. after OR Exit Time).

SEQ. #: 6840

Long Name: Post-Op-Pulm-Pneumonia

Short Name: CPPneum

Definition: Indicate whether the patient had pneumonia according to the CDC definition.

Intent/Clarification: See the CDC definition below:

<https://www.cdc.gov/nhsn/pdfs/pscmanual/6pscvapcurrent.pdf>

Table 1: Specific Site Algorithms for Clinically Defined Pneumonia (PNU1)

Imaging Test Evidence	Signs/Symptoms/Laboratory
<p>Two or more serial chest imaging test results with at least <u>one</u> of the following^{1,2}:</p> <p>New and persistent or Progressive and persistent</p> <ul style="list-style-type: none"> • Infiltrate • Consolidation • Cavitation • Pneumatoceles, in infants ≤ 1 year old <p>Note: In patients <u>without</u> underlying pulmonary or cardiac disease (e.g., respiratory distress syndrome, bronchopulmonary dysplasia, pulmonary edema, or chronic obstructive pulmonary disease), <u>one definitive</u> imaging test result is acceptable.¹</p>	<p>For ANY PATIENT, at least <u>one</u> of the following:</p> <ul style="list-style-type: none"> • Fever ($>38.0^{\circ}\text{C}$ or $>100.4^{\circ}\text{F}$) • Leukopenia (≤ 4000 WBC/mm³) or leukocytosis ($\geq 12,000$ WBC/mm³) • For adults ≥ 70 years old, altered mental status with no other recognized cause <p>And at least <u>two</u> of the following:</p> <ul style="list-style-type: none"> • New onset of purulent sputum³ or change in character of sputum⁴, or increased respiratory secretions, or increased suctioning requirements • New onset or worsening cough, or dyspnea, or tachypnea⁴ • Rales⁴ or bronchial breath sounds • Worsening gas exchange (e.g., O₂ desaturations (e.g., PaO₂/FiO₂ ≤ 240)², increased oxygen requirements, or increased ventilator demand) <p>ALTERNATE CRITERIA, for infants ≤ 1 year old:</p> <p>Worsening gas exchange (e.g., O₂ desaturations [e.g., pulse oximetry $<94\%$], increased oxygen requirements, or increased ventilator demand)</p> <p>And at least <u>three</u> of the following:</p> <ul style="list-style-type: none"> • Temperature instability • Leukopenia (≤ 4000 WBC/mm³) or leukocytosis ($\geq 15,000$ WBC/mm³) and left shift ($\geq 10\%$ band forms) • New onset of purulent sputum³ or change in character of sputum⁴, or increased respiratory secretions or increased suctioning requirements • Apnea, tachypnea⁴, nasal flaring with retraction of chest wall or nasal flaring with grunting • Wheezing, rales⁴, or rhonchi • Cough • Bradycardia (<100 beats/min) or tachycardia (>170 beats/min) <p>ALTERNATE CRITERIA, for child >1 year old or ≤ 12 years old, at least <u>three</u> of the following:</p> <ul style="list-style-type: none"> • Fever ($>38.0^{\circ}\text{C}$ or $>100.4^{\circ}\text{F}$) or hypothermia ($<36.0^{\circ}\text{C}$ or $<96.8^{\circ}\text{F}$) • Leukopenia (≤ 4000 WBC/mm³) or leukocytosis ($\geq 15,000$ WBC/mm³) • New onset of purulent sputum³ or change in character of sputum⁴, or increased respiratory secretions, or increased suctioning requirements • New onset or worsening cough, or dyspnea, apnea, or tachypnea⁴ • Rales⁴ or bronchial breath sounds • Worsening gas exchange (e.g., O₂ desaturations [e.g., pulse oximetry $<94\%$], increased oxygen requirements, or increased ventilator demand)

SEQ. #: 6845

Long Name: Post-Op-Venous Thromboembolism-VTE

Short Name: CVTE

Definition: Indicate whether the patient developed postoperative venous thrombosis or thromboembolic event.

Intent/Clarification: A clot within a blood vessel is called a thrombus and the process by which it forms is known as thrombosis. It can be damaging as it might block the flow of blood. Also, part of the clot could embolize, or break off, and block a blood vessel further along, cutting off the blood supply to important organs.

Post-operative patients are at risk of forming clots in the lower extremities that could lead to pulmonary embolism. Capture upper and lower extremity events.

SEQ. #: 6850

Long Name: Post-Op-Pulmonary Thromboembolism

Short Name: PulmEmb

Definition: Indicate whether the patient had a pulmonary thromboembolism diagnosed by radiologic study such as V/Q scan, angiogram, or spiral CT.

Intent/Clarification: Pulmonary embolism is a life threatening clot formation in one or more pulmonary arteries causing partial or complete obstruct of blood flow to the lung(s). Pulmonary embolisms must be documented through diagnostic testing.

SEQ. #: 6855

Long Name: Post-Op-Deep Venous Thrombosis

Short Name: DVT

Definition: Indicate whether patient had thrombosis (clot formation) in a deep vein.

Intent/Clarification: Deep vein thrombosis (DVT) is the formation of a blood clot in the deep veins within the body, such as in the leg or pelvis. This kind of thrombosis can occur after surgery and may cause redness, pain and swelling.

SEQ. #: 6860

Long Name: Post-Op-Pleural Effusion Requiring Drainage

Short Name: CPIEff

Definition: Indicate whether a post-operative pleural effusion required drainage via thoracentesis or chest tube insertion.

Intent/Clarification: Interventions include chest tube insertion, needle aspiration or other invasive procedure. May include hemothorax.

FAQ September 2017: A patient experienced a chylothorax postop from a chest tube that was in place from the OR. No new drains were placed. He ultimately was transferred to another acute care hospital that had the ability to perform a lymphsynthigraphy and was diagnosed with a thoracic duct injury. Should this get captured as a complication, and if so how?

Answer: Code pleural effusion requiring drainage for this chylothorax.

SEQ. #: 6865

Long Name: Post-Op-Pneumothorax Requiring Intervention

Short Name: PostOpPneumo

Definition: Indicate whether the patient had a post-operative pneumothorax requiring intervention.

Intent/Clarification: Interventions include chest tube insertion, needle aspiration or other invasive procedure. Do not capture a small pneumothorax followed with serial chest X-rays.

SEQ. #: 6870

Long Name: Post-Op-Renal-Renal Failure

Short Name: CRenFail

Definition: Indicate whether the patient had acute renal failure or worsening renal function resulting in ONE OR BOTH of the following:

1. Increase in serum creatinine level 3.0 x greater than baseline, or serum creatinine level ≥ 4 mg/dL. Acute rise must be at least 0.5 mg/dl
2. A new requirement for dialysis postoperatively.

Intent/Clarification:

The Acute Dialysis Quality Initiative, a multidisciplinary collaboration, defined a range of acute renal dysfunction called the RIFLE classification system. It is used to define grades of severity based on objective measurements.

See highlighted Renal Failure criteria below.

Classifications of Loss and End-stage disease are beyond the current scope of follow-up. Code yes if the patient meets the highlighted RIFLE Failure criteria or if dialysis was newly required post op.

Risk (R) - Increase in serum creatinine level X 1.5 or decrease in GFR by 25%, or UO < 0.5 mL/kg/h for 6 hours

Injury (I) - Increase in serum creatinine level X 2.0 or decrease in GFR by 50%, or UO < 0.5 mL/kg/h for 12 hours, or decrease in GFR by 75%; UO < 0.3 mL/kg/h for 24 hours, or anuria for 12 hours

Failure (F) – Increase in serum creatinine level X 3.0, or serum creatinine \geq mg/dL with at least a 0.5 mg/dL rise, or decrease in GFR by 75%; UO < 0.3 mL/kg/h for 24 hours, or anuria for 12 hours.

Loss (L) - Persistent ARF, complete loss of kidney function > 4 weeks

End-stage kidney disease (E) - Loss of kidney function >3 months

CLARIFICATION: If dialysis (seq# 375) is equal to "No" and if postoperative creatinine level (seq# 6555) is greater than or equal to 3X last creatinine level (seq# 585) or postoperative creatinine (seq# 6555) is greater than or equal to 4.0 with a 0.5 mg/dL rise or new postoperative dialysis (seq# 6875) then, renal failure (seq# 6870) is equal to "Yes".

SEQ. #: 6875

Long Name: Post-Op-Renal-Dialysis Req

Short Name: CRenDial

Definition: Indicate whether the patient had a new requirement for dialysis postoperatively, which may include hemodialysis, peritoneal dialysis.

Intent/Clarification: May include either hemo or peritoneal dialysis. This includes a one-time need for dialysis as well as implementation of longer term therapy.

If the patient was on preoperative peritoneal dialysis and moved to hemodialysis postoperatively, this does not constitute a worsening of the condition and should not be coded as an event. Does not include aquapheresis.

Continuous Veno-Venous Hemofiltration) (CVVH, CVVH-D) and Continuous Renal Replacement Therapy (CRRT) should be coded here as "Yes." (Code Ultra filtration as "No", it is captured in a separate field).

SEQ. #: 6880

Long Name: Post-Op-Dialysis Required After Discharge

Short Name: DialDur

Definition: Indicate whether dialysis was required after hospital discharge.

Intent/Clarification: The intent is to separate patients with possible long term dialysis from those that recovered kidney function prior to discharge.

SEQ. #: 6881

Long Name: Post-Op-Dialysis Duration

Short Name: DialStat

Definition: Indicate the duration of post-discharge dialysis.

Intent/Clarification: This may be temporary or permanent.

FAQ October 2017: Clarify the difference between temporary and permanent dialysis duration.

Answer: Code temporary if the patient no longer requires dialysis at the time of 30 day follow up after discharge.

Code permanent if the patient continues to require dialysis at the time of 30 day follow up after discharge.

SEQ. #: 6885

Long Name: Post-Op-Ultra Filtration

Short Name: CUltraFil

Definition: Indicate whether patient required Ultra filtration.

Intent/Clarification: Ultrafiltration is for fluid overload and is not counted as dialysis. Continuous Veno-Venous Hemofiltration) (CVVH, CVVH-D and Continuous Renal Replacement Therapy (CRRT) should be coded here as “No”, they are considered dialysis and should be captured in Seq# 6875

SEQ. #: 6890

Long Name: Post-Op-Vasc-Iliac/Fem Dissect

Short Name: CVaIFem

Definition: Indicate whether the patient had a dissection occurring in the iliac or femoral arteries.

Intent/Clarification: The origin of the event may have been at the site of cannulation or a preoperative catheterization insertion site, but the dissection occurred post-operatively.

SEQ. #: 6891

Long Name: Post-Op-Vasc-Acute Limb Isch

Short Name: CVaLblsc

Definition: Indicate whether the patient had any complication producing limb ischemia. This may include upper or lower limb ischemia.

Intent/Clarification: Ischemic events are restricted to the arterial system. These do not include venous system events, (i.e. DVT (deep vein thrombosis)). Example: A patient had an IABP removed and experienced an emboli which resulted in a necrotic great toe: Code “Yes” for acute limb ischemia.

SEQ. #: 6892

Long Name: Post-Op-Mechanical Assist Device Related Complication

Short Name: CMAD

Definition: Indicate whether there was a post-operative event related to a mechanical assist device.

Intent/Clarification:

- Yes
- No

SEQ. #: 6893

Long Name: Post-Op-MAD-Cannula / Insertion Site Issue

Short Name: CMADCanIns

Definition: Indicate whether the mechanical assist device related postoperative event included a cannula/insertion site issue.

Intent/Clarification: May include bleeding or infection.

SEQ. #: 6894

Long Name: Post-Op-MAD-Hemorrhagic

Short Name: CMADHem

Definition: Indicate whether there was hemorrhage related to a mechanical assist device

Intent/Clarification: Patients are at increased risk of bleeding due to anticoagulation and anti-platelet therapy, non-pulsatile blood flow leading to blood vessel malformation, and changes in blood-clotting factors.

SEQ. #: 6895

Long Name: Post-Op-MAD-Thrombotic/Embolic

Short Name: CMADThromEm

Definition: Indicate whether there was a thrombotic or embolic event related to a mechanical assist device

Intent/Clarification:

SEQ. #: 6896

Long Name: Post-Op-MAD-Hemolytic

Short Name: CMADHemolytic

Definition: Indicate whether there was a hemolytic event related to a mechanical assist device

Intent/Clarification: Patients may experience clinical signs of hemolysis (anemia, low hematocrit, hyperbilirubinemia) and a plasma free hemoglobin >40 mg/dL within 72 hours of VAD implant. Do not include hemolysis resulting from non-device causes.

SEQ. #: 6897

Long Name: Post-Op-MAD-Infection

Short Name: CMADInf

Definition: Indicate whether there was infection related to a mechanical assist device

Intent/Clarification: May include driveline/cannula infection, pump pocket infection, VAD endocarditis, sternal wound infection, or sepsis.

SEQ. #: 6898

Long Name: Post-Op-MAD-Other

Short Name: CMADOther

Definition: Indicate whether any other mechanical assist device related event occurred

Intent/Clarification: Device malfunctions, VAD thrombus, psychiatric episodes.

SEQ. #: 6900

Long Name: Post-Op-Rhythm Disturbance Requiring Perm Device

Short Name: CRhythmDis

Definition: Indicate whether patient developed a new dysrhythmia requiring insertion of a permanent device. Do not code these device insertions in the reoperation section even if performed in the OR.

Intent/Clarification: Include permanent pacemakers, Implantable cardioverter defibrillators (ICD) and combination devices. Do not code if the patient experiences third degree block and has temporary pacemaker wires inserted, but the block resolves and the patient does not require a permanent pacemaker.

- Pacemaker
 - ICD
 - Pacemaker/ICD
 - Other
 - None
-
-

SEQ. #: 6905

Long Name: Post-Op-Other-Card Arrest

Short Name: COtArrst

Definition: Indicate whether the patient had an acute cardiac arrest documented by one of the following:

- Ventricular fibrillation
- Rapid ventricular tachycardia with hemodynamic instability
- Asystole
- ICD shocks

Intent/Clarification: The cardiac arrest may be precipitated by ventricular fibrillation/tachycardia, asystole, or pulseless electrical activity (PEA). Code yes for sudden events requiring CPR. It is expected that all deaths inevitably have cardiac arrest, but this field is to capture those events that are sudden or acute in occurrence.

SEQ. #: 6906

Long Name: Post-Op-Other-Aortic Endoleak

Short Name: COtAortEndo

Definition: Indicate whether a post-operative endoleak occurred

Intent/Clarification: An **endoleak** is defined as persistent blood flow in the aneurysm sac through and around the endovascular seal and is the most common complication after endovascular aneurysm repair.

SEQ. #: 6907

Long Name: Post-Op-Other-Aortic Endoleak Type

Short Name: COtAortEndoTy

Definition: Indicate the type of endoleak

Intent/Clarification:

A **Type Ia endoleak** is defined as a leak occurring at the proximal seal zone.

A **Type Ib endoleak** is defined as a leak occurring at the distal seal zone.

A **Type II endoleak** is defined as retrograde filling of the aneurysm sac or false lumen in the case of dissection by aortic branch vessels (e.g. left subclavian artery, intercostal arteries, etc.).

A **Type III endoleak** is defined as leakage of blood into the aneurysm sac, or false lumen in the case of dissection, due to either a gap between separate endograft components, or a defect in the fabric of the graft secondary to graft strut fracture or erosion.

A **Type IV endoleak** is defined as the presence of an endoleak secondary to graft porosity. All other types of endoleaks must be definitively ruled out prior to selecting this diagnosis.

A Type **V endoleak**, also known as endotension, is defined as persistent aneurysm expansion in the absence of a confirmed endoleak.

SEQ. #: 6908

Long Name: Post-Op-Other-Aortic Rupture **Short Name:** COtAortRupt

Definition: Indicate whether aortic rupture occurred post op

Intent/Clarification:

SEQ. #: 6909

Long Name: Post-Op-Other-Aortic Dissection

Short Name: CVaAoDis

Definition: Indicate whether the patient had a dissection occurring in any part of the aorta.

Intent/Clarification: This includes ascending, arch, descending, thoracic or abdominal aorta. Aortic dissection is bleeding into or along the wall of the aorta. This does not include an aneurysmal event, unless it goes on to rupture or dissect.

SEQ. #: 6910

Long Name: Post-Op-Other-Aortic DissectionType

Short Name: CVaAoDisTy

Definition: Indicate the type of aortic dissection.

Intent/Clarification: May include antegrade, retrograde or both.

If the dissection extends proximally (i.e. back towards the aortic arch or ascending aorta) beyond the original extent then “**Retrograde**” should be selected.

If the dissection extends distally (i.e. downstream towards the descending or abdominal aorta) beyond the original extent then “**Antegrade**” should be selected.

If the dissection extends both proximally and distally then “**Both**” should be selected.

SEQ. #: 6911

Long Name: Post-Op-Other-Aortic Side Branch Malperfusion

Short Name: COtAortSide

Definition: Indicate whether aortic side branch malperfusion occurred

Intent/Clarification: The intent is to identify if aortic branch vessels have compromised flow in the post-operative period.

SEQ. #: 6912

Long Name: Post-Op-Other-Aortic Stent Graft Induced Entry Tear

Short Name: COtAortTear

Definition: Indicate whether an aortic stent graft induced entry tear occurred

Intent/Clarification:

This is typically due to the septum being fractured by a balloon or endograft, and the result is the creation of a new fenestration/connection between the true and false lumens of the dissection (so-called stent graft induced new entry tear (SINE)). This would typically be determined by the surgeon's assessment of the intraoperative completion angiogram, intravascular ultrasound, and/or transesophageal echocardiography.

SEQ. #: 6914

Long Name: Post-Op-Other-Anticoag Event

Short Name: COtCoag

Definition: Indicate whether the patient had bleeding, hemorrhage, and/or embolic events related to anticoagulant therapy postoperatively.

This may include patients who experience Disseminated Intravascular Coagulopathy (DIC) or Heparin Induced Thrombocytopenia (HIT).

Intent/Clarification: The intent of the field is to capture those patients that bleed, hemorrhage and /or suffer an embolic event related to anticoagulant therapy received post-op.

Abnormal coag lab tests without clinical events are not included.

Patients with DIC or HIT are included.

Patients with bleeding secondarily to surgical suture 'leaking' or general surgical 'oozing' are not to be included.

HIT (Heparin Induced Thrombocytopenia) is diagnosed with Heparin Assay and or D-Dimer laboratory tests only and are more than post-pump excessive bleeding or lower platelet counts. The physiological effects of CPB can reduce post-operative platelet counts as much as 50% within 24 hours.

SEQ. #: 6915

Long Name: Post-Op-Other-Pericardiocentesis

Short Name: COtTamp

Definition: Indicate whether the patient had a pericardiocentesis to remove fluid in the pericardial space compromising cardiac filling.

Intent/Clarification: Tamponade, fluid accumulation between the myocardium and pericardium of the heart, inhibits filling of the heart and results in hemodynamic compromise. Severity of tamponade may dictate the degree of intervention (invasive or non-invasive, surgical or Pericardiocentesis).

THIS FIELD IS FOR THOSE EVENTS THAT DO NOT REQUIRE RETURN TO THE OPERATING ROOM FOR TREATMENT.

SEQ. #: 6920

Long Name: Post-Op-Other-GI Event

Short Name: COtGI

Definition: Indicate whether the patient had a postoperative occurrence of any GI event, including but not limited to:

- GI bleeding requiring transfusion
- Pancreatitis with abnormal amylase/lipase requiring nasogastric (NG) suction therapy
- Cholecystitis requiring cholecystectomy or drainage
- Mesenteric ischemia requiring exploration
- ~~Hepatic failure (remove)~~
- Prolonged ileus
- Clostridium difficile

Intent/Clarification: GI events may require medical management, observational management, or surgical intervention to control.

DO NOT include events such as prolonged nausea and/or vomiting with no other documented physiological cause. Refer to the specific list included within the definition.

FAQ September 2017: Should we code a GI event for an abdominal compartment syndrome leading to a decompressive laparotomy?

Answer: Yes, code GI event for abdominal compartment syndrome in addition to the reoperation other non-cardiac for the decompression laparotomy.

SEQ. #: 6921

Long Name: Post-Op-Other-Liver Dysfunction or Failure

Short Name: COtLiver

Definition: Indicate whether the patient had liver dysfunction or failure.

Intent/Clarification: Refer to physician's documentation for diagnosis.

SEQ. #: 6925

Long Name: Post-Op-Other-Multi Sys Fail

Short Name: COtMSF

Definition: Indicate whether the patient had two or more major organ systems suffer compromised functions.

Intent/Clarification: Major organ systems are neurological, renal, pulmonary, cardiac, vascular or systemic. Multisystem Organ Failure (MSOF) means multiple organ systems have failed and function cannot be recovered by mechanical and/or pharmacological means. End-stage means irreversible organ failure.

SEQ. #: 6930

Long Name: Post-Op-Other-A Fib

Short Name: COtAFib

Definition: Indicate whether the patient experienced atrial fibrillation/flutter (AF) requiring treatment. Exclude patients who were in AFib at the start of surgery.

Intent/Clarification: Include any episode of A-Fib lasting longer than one hour and/or requiring treatment. Capture event(s) in all patients who were not in A-Fib at the start of surgery.

SEQ. #: 6950

Long Name: Post-Op-Other-Other

Short Name: COtOther

Definition: Indicate whether a postoperative event occurred that is not identified in the categories above yet impacts hospital length of stay and/or outcome.

Intent/Clarification: It is advised to restrict the capture of post-operative events to those that create a life threatening event, extended hospitalization, and/or require medical intervention to ward off clinical deterioration.

Discharge / Mortality

SEQ. #: 7000

Long Name: Date of Last Follow-Up

Short Name: LFUDate

Definition: Indicate the date on which the last follow-up was made. If patient dies in the hospital, this value will be the same as the date of death. If no follow-up is made after patient is discharged, this value will be the same as the discharge date.

Intent/Clarification:

This is the date that is last documented in the chart or obtained by contacting the physician's office. Required date format: mm/dd/yyyy

SEQ. #: 7001

Long Name: Mort-30d Status

Short Name: Mt30Stat

Definition: Indicate whether the patient was alive or dead at 30 days post-surgery (whether in hospital or not).

Intent/Clarification:

- Alive
 - Dead
 - Unknown
- -----

SEQ. #: 7002

Long Name: Mort-Op Death-Method Of Verification

Short Name: Mt30StatMeth

Definition: Indicate the primary method used to verify the patient's 30-day mortality status.

Intent/Clarification:

- Phone call to patient or family
 - Letter from medical provider
 - Evidence of life or death in medical record
 - Office visit on or after 30 days after the date of surgery.
 - Social Security Death Master File/NDI
 - Other
- -----

SEQ. #: 7005

Long Name: Discharge / Mortality Status

Short Name: DischMortStat

Definition: Indicate the discharge and current vital status of the patient

Intent/Clarification:

- In hospital, alive
- Died in hospital

- Discharged alive, last known status is alive
- Discharged alive, died after discharge

“In hospital, alive” refers to patient’s that are in the hospital at the 30 day mark that were never discharged. It is provided so sites do not get marked as missing on the required mortality fields for their composite scores/STAR ratings.

SEQ. #: 7008

Long Name: Date of Discharge

Short Name: DischDt

Definition: Indicate the date the patient was discharged from the hospital (acute care) even if the patient is going to a rehab or hospice or similar extended care unit within the same physical facility. If the patient died in the hospital, the discharge date is the date of death.

Intent/Clarification:

Required date format: mm/dd/yyyy

SEQ. #: 7009

Long Name: Discharge Location

Short Name: DisLoctn

Definition: Indicate the location to where the patient was discharged.

Intent/Clarification:

- **Home** (or, temporarily, at the home of a relative)
 - **Extended Care/Transitional Care Unit (TCU)/Rehab** (Code LTAC as Extended Care/Transitional Care Unit/Rehab. Do not count as part of acute care stay.
 - **Other Acute Care Hospital**
 - **Nursing Home**
 - **Hospice**
 - **Left AMA**
 - **Other**
- -----

SEQ. #: 7010

Long Name: Cardiac Rehabilitation Referral

Short Name: CardRef

Definition: Indicate whether advice was given or discussion conducted with the patient (by physician, nurse, or other personnel) regarding the importance of joining a cardiac rehabilitation program, or an appointment made.

Intent/Clarification:

Identify those patients who are referred to post discharge cardiac reconditioning and rehabilitation. Do not count Phase I, in hospital rehab, as “Yes”.

This is a Joint Commission endpoint and is to be documented on every patient. Cardiac rehabilitation programs are many times free standing or external to an acute care setting. Cardiac rehabilitation programs are designed specifically for the patients with cardiac disease who have medical and/or surgical recovery needs.

If the surgery was of Non-Cardiac nature (See Section N), code as “Not Applicable”.

If the patient is discharged with home health but a discussion regarding Phase II Cardiac Rehab occurred, choose “Yes”.

SEQ. #: 7011

Long Name: Smoking Cessation Counseling

Short Name: SmokCoun

Definition: Indicate whether, prior to discharge from the acute care facility, the patient received smoking cessation counseling. Please select "Not Applicable" for those patients with no prior history of smoking or remote (more than 1 year) history.

Intent/Clarification:

This is a Joint Commission endpoint and it must be documented that either literature and/or counseling was offered and provided to the patient.

Counseling should be provided to users of Cigarettes, Pipe, Cigars, Smokeless Cans, Other tobacco products (orbs, strips, sticks, hookah, etc.) It does not include e-cigs.

If the patient was not a smoker on the admission assessment or clinically, mentally or emotionally inappropriate for a referral, select “Not Applicable”.

GENERAL INFORMATION:

FAQ September 2017: When a patient leaves the hospital AMA and there are no prescriptions, how are discharge medications coded?

Answer: Code no.

SEQ. #: 7060

Long Name: Aspirin - Discharge

Short Name: DCASA

Definition: Indicate whether or not the patient was discharged from facility on Aspirin, or if it was contraindicated. The contraindication must be documented in the medical record by a physician, nurse practitioner, pharmacist or physician assistant.

Intent/Clarification:

Includes enteric coated and/or baby aspirin. Aspirin acts to “*decrease*” the blood viscosity and inhibits the clotting of platelets.

- **Yes:** Capture those who receive an order for Aspirin at discharge that contains at least 75mg ASA

- **No:** Patient did not receive an Aspirin order at discharge
- **Contraindicated** - Documented evidence of contraindication. If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record, such as notation of a medication allergy by Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist.

FAQ September 2017: Is the medication bundle excluded for an Isolated CAB due to AMA status?

Answer: No, the medications are not excluded from the composite measures.

SEQ. #: 7070

Long Name: ADP Inhibitors - Discharge

Short Name: DCADP

Definition: Indicate whether or not the patient was discharged from facility on an ADP inhibitor, or if it was contraindicated. The contraindication must be documented in the medical record by a physician, nurse practitioner, pharmacist or physician assistant.

Intent/Clarification:

These medications inhibit adenosine diphosphate (ADP) induced platelet aggregation (clotting) are often used to treat patients with a history of atherosclerotic cardiovascular disease to potentially reduce the incidence of major cardiovascular events (stroke, peripheral arterial disease, etc.).

- **Yes:** Capture those who receive an order for an ADP Inhibitor at discharge
- **No:** Patient did not receive an ADP Inhibitor order at discharge
- **Contraindicated:** Documented evidence of contraindication. If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record, such as notation of a medication allergy by a Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist.

FAQ September 2017: We noticed that there is no variable to capture P2Y12 at discharge in v2.9? How should they be coded.

Answer: Code P2Y12 as an ADP Inhibitor in sequence number 7070.

SEQ. #: 7075

Long Name: Other Antiplatelet - Discharge

Short Name: DCOthAntiplat

Definition: Indicate whether or not the patient was discharged from facility on any other antiplatelet medication, or if it was contraindicated. The contraindication must be

documented in the medical record by a physician, nurse practitioner, pharmacist or physician assistant.

Intent/Clarification:

- **Yes:** Capture those who receive an order for any other antiplatelet medication at discharge.
 - **No:** Patient did not receive any other antiplatelet medication order at discharge
 - **Contraindicated:** Documented evidence of contraindication. If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record (notation of a medication allergy prior to arrival) by a Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist.
-
-

SEQ. #: 7080

Long Name: Direct Thrombin Inhibitors - Discharge

Short Name: DCDirThromIn

Definition: Indicate whether or not the patient was discharged from facility on a direct thrombin inhibitor, or if it was contraindicated. The contraindication must be documented in the medical record by a physician, nurse practitioner, pharmacist or physician assistant.

Intent/Clarification:

Direct thrombin inhibitors (DTIs) are an innovative class of anticoagulants that bind directly to thrombin to inhibit its actions and impede the clotting process.

Bivalent:

- Bivalirudin (transient inhibition - is cleaved by thrombin)
- Lepirudin
- Desirudin

Univalent:

- Argatroban
- Melagatran (and its prodrug ximelagatran)
- Dabigatran

Allosteric Inhibitors:

No allosteric thrombin inhibitor has reached the stage of clinical trials.

- **Yes:** Capture those who receive an order for a thrombin inhibitor at discharge.
- **No:** Patient did not receive a thrombin inhibitor order at discharge.
- **Contraindicated:** Documented evidence of contraindication. If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record, such as notation of a medication allergy by a Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist

SEQ. #: 7085

Long Name: Warfarin (Coumadin) - Discharge

Short Name: DCCoum

Definition: Indicate whether or not the patient was discharged from facility on Warfarin (Coumadin), or if it was contraindicated. The contraindication must be documented in the medical record by a physician, nurse practitioner, pharmacist or physician assistant.

Intent/Clarification: The primary action of Coumadin/Warfarin is to prevent or delay blood coagulation.

- **Yes:** Capture those who receive an order for warfarin at discharge
- **No:** Patient did not receive a warfarin order at discharge
- **Contraindicated:** Documented evidence of contraindication. If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record, such as notation of a medication allergy by a Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist.

SEQ. #: 7090

Long Name: Factor Xa Inhibitors - Discharge

Short Name: DCFactorXa

Definition: Indicate whether or not the patient was discharged from facility on a factor Xa inhibitor, or if it was contraindicated. The contraindication must be documented in the medical record by a physician, nurse practitioner, pharmacist or physician assistant.

Intent/Clarification:

Direct factor Xa inhibitors ('xabans') are a class of anticoagulant drugs which act directly upon Factor X in the coagulation cascade, without using Antithrombin as a mediator.

- Apixaban (Eliquis)
- Betrixaban
- Darexaban
- Edoxaban (Savaysa)
- Otamixaban
- Rivaroxaban (Xarelto)
- Arixtra (Fondaparinux)
(not intended to be inclusive list)
- **Yes:** Capture those who receive an order for a Factor Xa inhibitor medication at discharge.
- **No:** Patient did not receive a Factor Xa inhibitor medication order at discharge
- **Contraindicated** - Documented evidence of contraindication. If a contraindication is documented explicitly as excluded for medical reasons,

or is evidenced clearly within the medical record (notation of a medication allergy prior to arrival) by a Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist.

SEQ. #: 7091

Long Name: Novel Oral Anticoagulant - Discharge

Short Name: DCNovOrAnti

Definition: Indicate whether or not the patient was discharged from facility on a Novel Oral Anticoagulant, or if it was contraindicated. The contraindication must be documented in the medical record by a physician, nurse practitioner, pharmacist or physician assistant.

Intent/Clarification: New agents have been introduced that are collectively referred to as **novel oral anticoagulants (NOACs)** or **directly acting oral anticoagulants (DOACs)**. They have been shown to be as good as or possibly better than the Coumadin with less serious side effects. The newer anticoagulants (NOACs/DOACs), are more expensive than the traditional ones and should be used with care in patients with kidney problems. Additionally, there is no antidote for the factor Xa inhibitors, so it is difficult to stop their effects in the body in cases of emergency (accidents, urgent surgery). Idarucizumab was FDA approved for the reversal of dabigatran in 2015.

- **Yes:** Capture those who receive an order for a Novel Oral Anticoagulant medication at discharge.
- **No:** Patient did not receive a Novel Oral Anticoagulant medication order at discharge
- **Contraindicated** - Documented evidence of contraindication. If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record (notation of a medication allergy prior to arrival) by a Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist.

FAQ September 2017: Clarify which medications should be coded in this category.

Answer: Confusion lies when the medications could be coded in any or all of the three categories, Factor Xa, Thrombin Inhibitors, or NOAC. Initially all three categories were included to differentiate medications that did not have antidotes. With more antidotes available, the three medications seem redundant. Capture the medication according to the manufacturer's category description. The only caveat to remember is that you should only code ONE of the three medication types so as not to have it appear that the patient is receiving more than one. Code consistently at your site.

SEQ. #: 7095

Long Name: Other Anticoagulant - Discharge

Short Name: DCOthAnticoag

Definition: Indicate whether or not the patient was discharged from facility on any other anticoagulant, or if it was contraindicated. The contraindication must be documented in the medical record by a physician, nurse practitioner, pharmacist or physician assistant.

Intent/Clarification: Examples: Heparin (unfractionated), Heparin (Low molecular weight), Enoxaparin/Lovenox, Dalteparin, Tinzaparin

- **Yes:** Capture those who receive an order for any other anticoagulant medication at discharge.
 - **No:** Patient did not receive an order for any other anticoagulant medication at discharge
 - **Contraindicated** - Documented evidence of contraindication. If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record (notation of a medication allergy prior to arrival) by a Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist.
-
-

SEQ. #: 7100

Long Name: ACE or ARB Inhibitors - Discharge

Short Name: DCACE

Definition: Indicate whether or not the patient was discharged from facility on ACE or ARB Inhibitors, or if it was contraindicated or not indicated (no history of CHF or EF>40%). The contraindication must be documented in the medical record by a physician, nurse practitioner, pharmacist or physician assistant.

Intent/Clarification: Primary use is for the treatment of hypertension but is also an essential treatment for congestive heart failure (reduces the workload of the heart). Routine, lifelong use of angiotensin converting enzyme inhibitors (ACEI) or angiotensin receptor blockers (ARB) is recommended for heart failure patients with a lower than usual ejection fraction (40 percent or less). Action is to dilate blood vessels to improve the amount of blood the heart is able to pump and thereby reducing the workload on the heart.

- **Yes** - Capture those who receive an order for an ACE or ARB inhibitor medication at discharge.
 - **No** – Patient did not receive a ACE or ARB inhibitor medication order at discharge
 - **Contraindicated** - Documented evidence of contraindication. If a contraindication is documented explicitly as excluded or medical reasons, or is evidenced clearly within the medical record (notation of a medication allergy prior to arrival) by a Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist.
 - **Not indicated** (no history of CHF or EF > 40%)
-
-

SEQ. #: 7103

Long Name: Amiodarone - Discharge

Short Name: DCAmiodarone

Definition: Indicate whether or not the patient was discharged from facility on Amiodarone, or if it was contraindicated. The contraindication must be documented in the medical record by a physician, nurse practitioner, pharmacist or physician assistant.

Intent/Clarification: Note that this value is specific to Amiodarone, rather than anti-arrhythmic drugs in general. Amiodarone is effective in situations where other anti-arrhythmic may fall short.

- **Yes** - Capture those who receive an order for Amiodarone or Multaq at discharge.
- **No** – Patient did not receive an Amiodarone or Multaq order at discharge.
- **Contraindicated** - Documented evidence of contraindication. If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record (notation of a medication allergy prior to arrival) by Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist.

SEQ. #: 7105

Long Name: Beta Blockers - Discharge

Short Name: DCBeta

Definition: Indicate whether or not the patient was discharged on beta blockers, or if beta blocker was contraindicated. The contraindication must be documented in the medical record by a physician, nurse practitioner, pharmacist or physician assistant.

Intent/Clarification: Beta blockers have been proven to increase survival of cardiac patients following MI and in the perioperative period.

Beta blockers are used for the treatment of high blood pressure, treating chest pain or angina, controlling irregular heart rhythms, slowing ventricular rate response and for the treatment of congestive heart failure.

- **Yes** - Capture those who receive an order for a beta blocker medication at discharge.
- **No** – Patient did not receive a beta blocker medication order at discharge.
- **Contraindicated** - Documented evidence of contraindication. If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record (notation of a medication allergy prior to arrival) by a Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist.

SEQ. #: 7115

Long Name: Lipid Lowering Statin - Discharge

Short Name: DCLipLowStat

Definition: Indicate whether or not the patient was discharged from facility on a Statin, or if it was contraindicated. The contraindication must be documented in the medical record by a physician, nurse practitioner, pharmacist or physician assistant.

Intent/Clarification: Lipid lowering medications block the production of cholesterol and fat. Depending upon the specific medication, each may target unique levels such as HDL (good cholesterol), LDL (bad cholesterol) and triglycerides or polipoprotein B (protein needed to produce cholesterol). They may also reduce the absorption of dietary cholesterol by combining with the cholesterol to remove it from the bloodstream.

Statin medications typically have a generic name ending in the suffix 'statin'. However, some combination statin/non- statin drugs have other generic names. Do not capture non-statins here unless combined with a statin.

- **Yes** - Capture those who receive an order for a statin at discharge.
- **No** – Patient did not receive a statin order at discharge
- **Contraindicated** - Documented evidence of contraindication. If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record (notation of a medication allergy prior to arrival) by a Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist.

SEQ. #: 7120

Long Name: Lipid Lowering - Other - Discharge

Short Name: DCLipLowNonStat

Definition: Indicate whether or not the patient was discharged from facility on a lipid-lowering medication other than a statin, or if it was contraindicated. The contraindication must be documented in the medical record by a physician, nurse practitioner, pharmacist or physician assistant.

Intent/Clarification: Non-statin and other medications prescribed at discharge do not meet the measures according to new Heart Association guidelines.

Lipid lowering medications block the production of cholesterol and fat. Depending upon the specific medication, each may target unique levels such as HDL (good cholesterol), LDL (bad cholesterol) and triglycerides or polipoprotein B (protein needed to produce cholesterol). They may also reduce the absorption of dietary cholesterol by combining with the cholesterol to remove it from the bloodstream. New AHA guidelines favor Statin use and question efficacy of non-statins.

Examples: Fish oils, Niacor, Niaspan, Zetia, Fenofibrate, Tricor, Triglide, Lopid, Colestid, Prevalite, Questran, Welchol

- **Yes** - Capture those who receive an order for a non-statin medication at discharge.
- **No** – Patient did not receive a non-statin medication order at discharge
- **Contraindicated** - Documented evidence of contraindication. If a contraindication is documented explicitly as excluded for medical reasons, or is evidenced clearly within the medical record (notation of a medication allergy prior

to arrival) by a Physician, Nurse Practitioner, Anesthesia, Physician Assistant, or Pharmacist.

SEQ. #: 7121

Long Name: Mort-Date

Short Name: MtDate

Definition: Indicate the date the patient was declared dead.

Intent/Clarification:

Provide the date the patient died in hospital or was discharged alive, died after discharge within 30 days.

Date in the format mm/dd/yyyy.

SEQ. #: 7122

Long Name: Mort-Prim Cause

Short Name: MtCause

Definition: Indicate the PRIMARY cause of death, i.e., the first significant abnormal event which ultimately led to death.

Intent/Clarification: If the patient died due to multiple organ system failure, select the system that either was the initiator of the Multisystem Organ Failure (MSOF) or the primary cause of the patient's demise.

- Cardiac
 - Neurologic
 - Renal
 - Vascular
 - Infection
 - Pulmonary
 - Unknown
 - Other
-
-

SEQ. #: 7123

Long Name: In-Hospital Death location

Short Name: InHospDthLoc

Definition: Indicate the location within the hospital where the patient died.

Intent/Clarification:

- OR During Initial Surgery
- OR During Reoperation
- In-hospital (Other than OR)

SEQ. #: 7124

Long Name: Mort-Op Death

Short Name: MtOpD

Definition: Operative Mortality includes: (1) all deaths, regardless of cause, occurring during the hospitalization in which the operation was performed, even if after 30 days (including patients transferred to other acute care facilities); and (2) all deaths, regardless of cause, occurring after discharge from the hospital, but before the end of the thirtieth postoperative day.

Intent/Clarification: Includes all causes of death including deaths occurring in hospitals when the patient is transferred to a higher level of care.

SEQ. #: 7125

Long Name: Post Discharge Death Location

Short Name: PostDisDthLoc

Definition: Indicate the location where the patient died after being discharged from the original hospitalization.

Intent/Clarification:

- **Home** (or, temporarily, at the home of a relative)
 - **Extended Care Facility/Transitional Care Unit (TCU)** (Code LTAC as Extended Care/Transitional Care Unit/Rehab. Do not count as part of acute care stay.)
 - **Hospice**
 - **Acute Rehabilitation** (Ultimate plan for patient to return home after a short-stay)
 - **Hospital, During Readmission**
 - **Other**
 - **Unknown**
- -----

Readmission

SEQ. #: 7140

Long Name: Readmission

Short Name: Readmit

Definition: Indicate whether the patient was readmitted to the hospital within 30 days of discharge from hospitalization for this surgery. Code yes for inpatient admission to an acute care facility. Do not capture ED or outpatient visits or admission to a skilled facility or nursing home.

Intent/Clarification:

This is not part of the composite score.

The intent is to capture inpatient readmissions to acute care and primary care facilities only where the patient **status is listed as “In-Patient”**.

- Obtain information as close to 30 days from date of discharge as possible.
- It is understood that some readmissions are planned; these are still counted as readmissions.
- To code “Yes”, readmissions do not need to be at same institution where the initial surgical procedure was done.
- Discharge and readmission to a psychiatric care facilities, where the patient is considered an in-patient are to be considered as readmissions.
- Do not include Emergency Department visits or observation status visits unless the ED visits leads to status of in-patient.
- If a patient is readmitted to an in-patient rehabilitation hospital, code “No”.
- If a patient is readmitted to an LTAC, code “No”.
- Do not code transfers to higher level of care, this is considered an extension of the same acute care admission. If the patient was discharged to the “Acute Rehab” floor of the same hospital and then readmitted back as an in-patient back into a nursing floor, code “Yes” to admission as an inpatient is considered “Yes.”
- To align with CMS, 30 day readmission should not be coded for patients who remain in observation units, no matter the duration.

On occasion a patient is readmitted twice within the 30 day time frame from the date of the procedure. This is a Yes/No question, and does not ask how many times readmitted. Any time the patient is readmitted to a hospital \leq 30 days from the date of discharge regardless if the readmission was planned or unplanned, related or unrelated. You code the first readmission only.

SEQ. #: 7145

Long Name: Date of Readmission

Short Name: ReadmitDt

Definition: Indicate the date the patient was readmitted.

Intent/Clarification:

Indicate the date the patient was readmitted with a status of In-patient. If the patient was admitted with the status of “Observation” but later changed to “In-patient” code the date the patient was changed to in-patient status.

Required date format: mm/dd/yyyy

SEQ. #: 7160

Long Name: Readmit Reason

Short Name: ReadmRsn

Definition: Indicate the primary reason that the patient was readmitted as an in-patient.

Intent/Clarification:

The intent is to identify readmissions where conditions have a physiologic relationship to cardiothoracic surgery.

If the patient was readmitted multiple times, use the first readmission to code this section. Example: If the patient was readmitted twice, the first time with pneumonia and the second time with angina, code pneumonia.

Readmit Reason must be completed if known.

- **Angina:** chest pain or discomfort often spreading to the shoulders, arms, and/or neck, caused by inadequate blood supply to the heart; stable or unstable.
- **Anticoagulant Complication - Pharmacological:** relates to a bleeding complication secondary to the administration of an anticoagulant, IIb/IIIa inhibitor or other platelet inhibitor, for example Plavix, Coumadin, ReoPro etc. This is often diagnosed as Sub-therapeutic or Supra-therapeutic INR.
- **Anticoagulant Complications- Valvular:** relates to thrombus forming in, on and around the prosthetic valve.
- **Aortic Complication:** may relate to issues in the native aorta or be secondary to aortic procedures.
- **Arrhythmia / Heart Block:** Patient admitted due to rhythm irregularities that may have required pharmacological, non- invasive, or invasive treatment.
- **Blood Pressure:** (hyper or hypotension)
- **Chest Pain, non-cardiac**
- **Congestive Heart Failure:** May be manifested as pulmonary edema or only identified as “heart failure”. Must have a diagnosis of Congestive Heart Failure.
- **Coronary Artery / Graft Dysfunction:** This may include native vessels and/or conduit restenosis, spasm or dissection.
- **Depression/psychiatric issue**
- **DVT (Deep Venous Thrombosis):** the formation of a blood clot in the deep veins within the body, such as in the leg or pelvis diagnosed by ultrasound.
- **Electrolyte imbalance**
- **Endocarditis:** Confirmed diagnosis of endocarditis by blood culture and/or vegetation on or around a heart valve. This may include native tissue, ring or prosthetic valve involvement.
- **Failure to thrive:** weight loss of more than 5%, decreased appetite, poor nutrition, and physical inactivity, often associated with dehydration, depression, immune dysfunction.
- **GI issue:** may require medical management, observational management or surgical intervention to control.
- **Infection- Conduit Harvest Site:** Use CDC definitions.
- **Infection- Deep Sternum / Mediastinitis:** Use CDC definitions. May or may not require surgical intervention.
- **Mental status changes:** Any other mental status change not diagnosed as Stroke or Transient Ischemic Attack
- **Myocardial Infarction:** MI diagnosis and/or angina diagnosed by the criteria listed in the definition. Prior to coding as MI, verify with discharge diagnosis to assure that the MI was ‘ruled in’ or that the patient’s reported angina was not secondary to chest wall pain, as diagnosed with echocardiography, chest x- ray or other methods.

- **PE (Pulmonary Embolism):** Pulmonary embolisms must be documented through diagnostic testing such as VQ scan, angiogram, or CT. Do not confuse Pulmonary Embolism with Pulmonary Edema, captured under 'Respiratory Complication, Other'.
- **Pericardial Effusion and/or Tamponade:** May or may not require invasive intervention on readmission i.e. re-exploration or pericardial tap.
- **Pericarditis/Post Cardiotomy Syndrome:** Inflammatory reaction involving the pericardium that may include fever, effusion, pain.
- **Pleural Effusion Requiring Intervention:** A pleural effusion is a buildup of fluid between the layers of tissue that line the lungs and chest cavity. Diagnosis is often made through imaging studies. Intervention may consist of Thoracentesis (often by Interventional Radiology), Chest tube, Pleural drain (including pleural catheter or pigtail catheter), or Pleural decortication. Intervention does not necessarily entail an OR visit. Many procedures are done at ICU bedside.
- **Pneumonia:** Pneumonia is an inflammation of the lungs, typically diagnosed by microbiology of sputum cultures. It can be detected by imaging studies but should have confirming evidence. Include aspiration pneumonia. Look for documentation in medical record notes.
- **Renal Failure:** Use "Failure" criteria highlighted in RIFLE criteria.
- **Renal Insufficiency:** dysfunction of the kidneys with accumulation of waste products in the blood.
- **Respiratory Complication, Other:** Include acute respiratory failure (often requiring emergent intubation or ECMO cannulation), hypoxemia, pulmonary edema, respiratory acidosis. Pneumonia is separately captured.
- **Sepsis:** See definition of sepsis in the post-operative events section.
- **Stroke:** Confirmed neurological deficit of abrupt onset caused by a disturbance in blood flow to the brain that did not resolve within 24 hours.
- **TIA (Transient Ischemic Attack):** Neurological dysfunction that lasts less than 24 hours and is completely resolved.
- **Transfusion:**
- **Transplant Rejection:** There are two forms of acute rejection: cellular and vascular. The chances of acute cellular rejection are greatest during the first six months after transplant. Acute vascular rejection is a type of acute rejection that occurs early after transplant (within the first four months) in a small number of patients.
- **VAD Complication:** Any device failure or malfunction of a VAD. Some physiologic complications, such as hemorrhagic stroke, hemolysis, or GI bleeds and be related to VAD complications.
- **Valve Dysfunction:** Can be either structural (i.e. leaflet fracture, impaired leaflet function, calcification) or non- structural (perivalvular leak, hemolytic anemia, pannus obstruction) dysfunction. Is applicable to either a mechanical or tissue valve. Dysfunction related to Endocarditis is captured separately.
- **Vascular Complication, Acute:** Any major arterial or venous circulatory compromise that requires pharmacological, non- invasive or invasive treatment to resolve; i.e. peripheral delivery of TPA, peripheral angioplasty. Include acute limb ischemia that may require fasciotomy or amputation for treatment. DVT (Deep Vein Thrombosis) is captured separately.
- **Wound:** Other (drainage, cellulitis)
- **Other – Related Readmission:** Those conditions that may have a correlation to cardiothoracic surgery.
- **Other – Nonrelated Readmission:** All other reasons for admission, i.e., trauma, cancer, that are not related to the initial cardiac surgery or its complications.

- **Other – Planned Readmission:** Readmission for a procedure that was conditional upon surgical remediation of a cardiac condition. Example: A patient is re-admitted to the hospital after CABG for reasons that were planned prior to cardiac surgery (e.g., colon resection or kidney transplant).
 - **Unknown:** Use this field selection only if there is no information available as to the reason why the patient returned. All effort should be made to identify the reason.
-
-

SEQ. #: 7165

Long Name: Readmit Reason - Primary Procedure

Short Name: ReadmPro

Definition: Indicate the primary procedure that the patient received after being readmitted as an in-patient.

Intent/Clarification:

If the patient was readmitted multiple times, use the first readmission to code this section. Example: If the patient was readmitted twice and did not have any procedures on the first visit, but had a Cath-lab intervention on the second visit, code No Procedure Performed.

- **No Procedure Performed:** There was no invasive or a non-invasive procedure performed. Patient may have been managed by medical observation, pharmacological or other medical therapies. Blood transfusions, ECGs, ordinary x-ray imaging, and IV infusions are not considered 'procedures'.
- **Cath Lab for Valve Intervention:** Valvuloplasty, TAVR, mitral clip and related procedures.
- **Cath Lab for Coronary Intervention (PCI):** Percutaneous coronary intervention, angioplasty, STENT or other coronary occlusive therapies in the Cath Lab.
- **Dialysis:** The patient required new hemo or peritoneal dialysis. May include CRRT.
- **OR for Bleeding:** Bleeding due to pericardial tamponade or related to a prior cardiac surgery. Includes repair of ventricular lacerations. (Note that OR visit is not an absolute requirement. Procedures done at ICU bedside to control mediastinal bleeding are included, as defined under Postoperative Event.)
- **OR for Coronary Artery Intervention:** Any surgical intervention on any of the coronary arteries due to progressive native coronary disease, conduit spasm, occlusion or dissection.
- **OR for Sternal Debridement / Muscle Flap:** Any surgical intervention necessary to debride (clean or remove marginal tissue or muscle) or Plastic Surgeon involvement to perform muscle flap reconstruction for deep sternal wound infection.
- **OR for Valve Intervention:** Any surgical procedure performed (repair and/or replacement) on any heart valve; native, prosthetic or ring/band device.
- **OR for Vascular Procedure:** Any (arterial) vascular surgical procedure required. Examples would include but are not limited to: (femoral hematoma evacuation, PTA, AAA, Carotid Endarterectomy, Fem-Pop bypass etc.)
- **OR for Aorta Intervention:**
- **Pacemaker Insertion / ICD:** Permanent Pacemaker or Implantable Cardioverter Defibrillator for arrhythmia or heart block.

- **Pericardiotomy / Pericardiocentesis:** Pericardiotomy is removal of all or part of the pericardium. Pericardiocentesis is drainage of accumulated fluid from or around the heart that creates hemodynamic compromise for the patient. Pericardiocentesis is typically performed as a non- surgical intervention, but a more invasive approach can be achieved through the surgical procedure of pericardial window.
- **Planned non-cardiac procedure:** Example: Planned colon resection.
- **Thoracentesis / Chest Tube Insertion:** Thoracentesis is a procedure to remove fluid from the space between the lungs and the chest wall called the pleural space. It is done with a needle. For persistent fluid accumulation, a chest tube can be inserted for more long-term drainage.
- **Wound Vac:** Wound Vac therapy promotes surgical wound healing through Negative Pressure Wound Therapy (NPWT). By delivering negative pressure (a vacuum) at the wound site, this helps draw wound edges together, remove infectious materials and actively promote granulation.
- **Other Procedure:** Some type of invasive or non-invasive procedure was performed that is not included in the above referenced list.
- **Unknown:** Use this field selection only if there is no information available as to the treatment/intervention prescribed. All effort should be made to identify the treatment used.

SEQ. #: 7166

Long Name: Readmit Reason - Primary Procedure - Aorta Intervention Type

Short Name: ReadmAortIntTy

Definition: Indicate the type of aortic intervention required during readmission.

Intent/Clarification: Indicate if the patient requires an open or endovascular aorta procedure.

SEQ. #: 7167

Long Name: Readmit Reason - Primary Procedure - Aorta Intervention Indication

Short Name: ReadmAortIntInd

Definition: Select the indication for aortic reintervention

Intent/Clarification: Indications for an aorta reintervention procedure include:

- Rupture
- Endoleak
- Infection
- Dissection
- Expansion
- Loss of side branch patency
- Other

Risk Scores

SEQ. #: 7170

Long Name: Predicted Risk of Mortality

Short Name: PredMort

Definition: Indicate the Predicted Risk of Mortality.

Intent/Clarification: The prediction of risk is calculated within the software according to a DCRI model specified to the vendor. **No user data input is accepted.**

SEQ. #: 7175

Long Name: Predicted Deep Sternal Wound Infx

Short Name: PredDeep

Definition: Indicate the Predicted Risk of Deep Sternal Wound Infection.

Intent/Clarification: The prediction of risk is calculated within the software according to a DCRI model specified to the vendor. **No user data input is accepted.**

SEQ. #: 7180

Long Name: Predicted Reoperation

Short Name: PredReop

Definition: Indicate the Predicted Risk of Reoperation.

Intent/Clarification: The prediction of risk is calculated within the software according to a DCRI model specified to the vendor. **No user data input is accepted.**

SEQ. #: 7185

Long Name: Predicted Permanent Stroke

Short Name: PredStro

Definition: Indicate the Predicted Risk of Permanent Stroke.

Intent/Clarification: The prediction of risk is calculated within the software according to a DCRI model specified to the vendor. **No user data input is accepted.**

SEQ. #: 7190

Long Name: Predicted Prolonged Ventilation

Short Name: PredVent

Definition: Indicate the Predicted Risk of Prolonged Ventilation.

Intent/Clarification: The prediction of risk is calculated within the software according to a DCRI model specified to the vendor. **No user data input is accepted.**

SEQ. #: 7195

Long Name: Predicted Renal Failure

Short Name: PredRenF

Definition: Indicate the Predicted Risk of Renal Failure.

Intent/Clarification: The prediction of risk is calculated within the software according to a DCRI model specified to the vendor. **No user data input is accepted.**

SEQ. #: 7200

Long Name: Predicted Morbidity or Mortality

Short Name: PredMM

Definition: Indicate the Predicted Risk of Morbidity or Mortality.

Intent/Clarification: The prediction of risk is calculated within the software according to a DCRI model specified to the vendor. **No user data input is accepted.**

SEQ. #: 7205

Long Name: Predicted Short Length of Stay

Short Name: Pred6D

Definition: Indicate the Predicted Risk of Short Length of Stay.

Intent/Clarification: The prediction of risk is calculated within the software according to a DCRI model specified to the vendor. **No user data input is accepted.**

SEQ. #: 7210

Long Name: Predicted Long Length of Stay

Short Name: Pred14D

Definition: Indicate the Predicted Risk of Long Length of Stay.

Intent/Clarification: The prediction of risk is calculated within the software according to a DCRI model specified to the vendor. **No user data input is accepted.**

STS Temporary Fields

STS Temporary fields should only be used at the direction of STS. Do not use for local data collection and clear any data that may have been entered prior to submission. See field 7230 below for instructions.

SEQ. #: 7215

Long Name: Temporary Yes/No Field #1

Short Name: TempYN1

Definition: This is a temporary field that should not be used for data collection until expressly instructed to by the STS.

Intent/Clarification: Use only as directed by STS, do not add custom field here.

SEQ. #: 7220

Long Name: Temporary Yes/No Field #2

Short Name: TempYN2

Definition: This is a temporary field that should not be used for data collection until expressly instructed to by the STS.

Intent/Clarification: Use only as directed by STS, do not add custom field here.

SEQ. #: 7225

Long Name: Temporary Date Field

Short Name: TempDt

Definition: This is a temporary field that should not be used for data collection until expressly instructed to by the STS.

Intent/Clarification: Use only as directed by STS, do not add custom field here.

SEQ. #: 7230

Long Name: Temporary Coded Field

Short Name: TempCode

Definition: This is a temporary field that should not be used for data collection until expressly instructed to by the STS.

Intent/Clarification: Use only as directed by STS, do not add custom field here.

SEQ. #: 7235

Long Name: Temporary Text Field

Short Name: TempText

Definition: This is a temporary field that should not be used for data collection until expressly instructed to by the STS.

Intent/Clarification: Use only as directed by STS, do not add custom field here.

Adult Cardiac Anesthesiology

FAQ 2017: If the anesthesiologist is not a participant in the STS Anesthesia module, DCRI will not include the fields in the anesthesia module in the DQR.

SEQ. #: 7310

Long Name: Primary Anesthesiologist Name

Short Name: PrimAnesName

Definition: Indicate the full name of the primary anesthesiologist for the procedure.

Intent/Clarification:

Field must be populated. Missing data or information for an anesthesiologist not on your current contract with the STS will cause your data file submission not to process.

SEQ. #: 7315

Long Name: Primary Anesthesiologist National Provider Identifier

Short Name: PrimAnesNPI

Definition: Indicate the individual-level National Provider Identifier (NPI) of the primary anesthesiologist for the procedure.

Intent/Clarification:

Field must be populated. Missing or inaccurate data will cause your data file submission not to process. It is crucial to enter the correct anesthesiologist identifier.

SEQ. #: 7320

Long Name: Care Team Model

Short Name: AnesCareTeamMod

Definition: Indicate the anesthesia care team assigned for the predominant portion of the procedure.

Intent/Clarification:

Determine the care model primarily responsible for providing anesthesia to the patient intraoperatively. This information can be found on the anesthesia record. Check with your anesthesia team or leave blank if the data is not available.

SEQ. #: 7325


Long Name: Pain Score Baseline

Short Name: PainScorePre

Definition: Indicate the highest baseline (preoperative) pain score on the 0-10 integer scale, or indicate that the score was not recorded.

Intent/Clarification:

Pain score, which is a quality metric, is routinely assessed as part of preoperative holding area check in list. This information should be obtainable from a progress note or similar documentation completed by preoperative nurse closest to the OR Entry time.



SEQ. #: 7330

Long Name: Transfusion Algorithm to Guide Transfusion

Short Name: TransfAlg

Definition: Indicate whether a transfusion algorithm or guideline was used to guide transfusion in the patient.

Intent/Clarification:

A transfusion algorithm or guideline is a predetermined set of treatment plans specific to various patient specific criteria to aid in transfusing the patient. Check with your anesthesia team or leave blank if the data is not available.

SEQ. #: 7335

Long Name: Cell saver volume

Short Name: CellSavVol

Definition: Indicate the volume of cell-saver blood that was transfused intraoperatively. Include any volume started in the OR, even if the infusion completed postoperatively.

Do not include autologous, allogeneic, pump-residual, or chest-tube recirculated blood. Value should be recorded in milliliters.

Intent/Clarification:

Cell-saver blood is blood that the patient loses during surgery which is transfused back to the patient. Time frame includes any cell-saver infusions started intraoperatively regardless if they completion time is after OR Exit date/time. This type of data could be obtained from the Perfusionist record who was assigned to that specific case.

Some hospitals will bag the residual pump blood and the anesthesiologist hangs it and gives some extra protamine. This is not the same as Cell Saver blood and should not be included here.

SEQ. #: 7340

Long Name: Heparin Total Dose

Short Name: TotHep

Definition: Indicate the total dose of heparin that was administered intraoperatively prior to the initiation of first cardiopulmonary bypass.

Include all doses of heparin given prior to the first cardiopulmonary bypass. Value should be recorded in units.

Intent/Clarification:

Heparin administered after OR Entry time and prior to the initiation of cardiopulmonary bypass. Measurement should be recorded in units.

SEQ. #: 7345

Long Name: Heparin Management

Short Name: HepMgmt

Definition: Indicate the method of heparin management used intraoperatively.

Different approaches are utilized to measure the adequacy of heparinization for anticoagulation.

Intent/Clarification:

The adequacy of heparinization determines the coaguability of the patient's blood. Heparin titration based on activated clotting time (ACT) measures how quickly the blood will clot. The larger the number the longer it will take for the blood to clot. Heparin titration based on heparin concentration (Hepcon System) measures the concentration of heparin in the blood. If either of these two measurement are not used to determine the level of heparinization then "other" should be chosen.

SEQ. #: 7350

Long Name: Protamine total dose

Short Name: TotProt

Definition: Indicate the total dose of protamine given intraoperatively to reverse heparinization after first cardiopulmonary bypass.

Value should be recorded in milligrams. Do not include doses given in the ICU.

Intent/Clarification:

Protamine is a medication given used to reverse the effects of heparin within the operating room. Time frame should be after the initiation of cardiopulmonary bypass and prior to ICU admission.

SEQ. #: 7351

Long Name: Antithrombin III Total Dose

Short Name: AntithromDose

Definition: indicate the total dose of antithrombin III

Intent/Clarification:

Antithrombin III is a medication given to enhance the heparin effect to achieve adequate anticoagulation. Time frame should be any amount given within the intraoperative phase after OR Entry time.

SEQ. #: 7360

Long Name: Viscoelastic Testing Used During Operation

Short Name: IntraViscoTest

Definition: Indicate whether viscoelastic testing was used intraoperatively (example: TEG, TEG-FF, or ROTEM).

Viscoelastic testing is a method of measuring coagulation in the blood.

Intent/Clarification:

Viscoelastic testing is used to determine which coagulation products to administer when the patient has an anticipated coagulopathy or non-surgical cause of bleeding. CT anesthesia team or patient's lab record may be useful to see whether any of the above indicated viscoelastic tests has been performed.

SEQ. #: 7365

Long Name: Volatile Agent Used

Short Name: VolAgentUsed

Definition: Indicate whether a volatile agent was used.

Intent/Clarification:

A volatile anesthetic is an inhaled anesthetic administered via an anesthetic gas machine or via the cardiopulmonary bypass machine.

SEQ. #: 7366

Long Name: Volatile Agent - Isoflurane

Short Name: VolAgentIso

Definition: Indicate whether the volatile agent used was Isoflurane

Intent/Clarification:

Indicate if isoflurane was the volatile anesthetic used to provide anesthesia.

SEQ. #: 7367

Long Name: Volatile Agent - Sevoflurane

Short Name: VolAgentSevo

Definition: Indicate whether the volatile agent used was Sevoflurane

Intent/Clarification:

Indicate if sevoflurane the volatile anesthetic used to provide anesthesia.

SEQ. #: 7368

Long Name: Volatile Agent - Desflurane

Short Name: VolAgentDes

Definition: Indicate whether the volatile agent used was Desflurane

Intent/Clarification:

Indicate if desflurane was the volatile anesthetic used to provide anesthesia.

SEQ. #: 7369

Long Name: Volatile Agent - Other

Short Name: VolAgentOth

Definition: Indicate whether any other volatile agent was used

Intent/Clarification:

Although highly unlikely, indicate if any other volatile agents were used to provide anesthesia. Information may be obtained from anesthesia record or perfusion record.

SEQ. #: 7370

Long Name: Volatile Agent Timing - Pre-CPB

Short Name: VolAgentTimPre

Definition: Indicate whether the volatile agent was used prior to the patient being on CPB.

Intent/Clarification:

Time frame of administering a volatile agent is after OR entry and prior to CPB initiation.

SEQ. #: 7375

Long Name: Volatile Agent Timing - During CPB

Short Name: VolAgentTimDur

Definition: Indicate whether the volatile agent was used during the period when patient was on CPB.

Intent/Clarification:

A volatile agent was administered during the use of cardiopulmonary bypass. This information will either come from intraoperative anesthesia chart or perfusion chart. Leave blank if the information is unavailable.

SEQ. #: 7380

Long Name: Volatile Agent Timing - Post CPB

Short Name: VolAgentTimPost

Definition: Indicate whether the volatile agent was used after the patient was taken off CPB.

Intent/Clarification:

Indicate if a volatile agent was administered after the discontinuation of cardiopulmonary bypass and prior to admission to the ICU.

SEQ. #: 7385

Long Name: Volatile Agent Timing - Maintenance (no CPB)

Short Name: VolAgentTimMaint

Definition: Indicate whether a volatile agent was used for maintenance in a non-pump case (no CPB).

Intent/Clarification:

A volatile agent was administered after entry into the OR and prior to ICU admission in off-pump cases.

SEQ. #: 7390

Long Name: Intraop Infusion: Dexmedetomidine

Short Name: DexIntra

Definition: Indicate the use of dexmedetomidine infusion during surgery.

Any use of dexmedetomidine infusion during the intraoperative period, usually but not always, in the post-bypass period.

Intent/Clarification:

Indicate if dexmedetomidine was administered after OR Entry time and prior to OR Exit time

SEQ. #: 7395

Long Name: Intraop Infusion: Propofol

Short Name: PropIntra

Definition: Indicate the use of propofol infusion during surgery.

Any use of a propofol infusion during the intraoperative period, usually but not always, in the post-bypass period.

Intent/Clarification:

Indicate if Propofol was administered by infusion after OR Entry time and prior to OR Exit time.

Specific attention should be paid to exclude bolus Propofol administration during any time of the intraoperative phase.

SEQ. #: 7400

Long Name: Intraop Mgs of Midazolam

Short Name: MidazIntra

Definition: Indicate the intraoperative does of midazolam in milligrams. Enter zero if no midazolam used.

Intent/Clarification:

Record in milligrams the amount of midazolam administered after OR Entry and prior to OR Exit. Record "0mg" if no midazolam was administered intraoperatively.

SEQ. #: 7405

Long Name: Intraop Insulin Total Dose (max units)

Short Name: TotInsulIntra

Definition: Indicate the total units (bolus and infusion) of insulin administered intraoperatively. Enter zero if no insulin was given.

Intent/Clarification:

Record, in units, the amount of insulin administered after OR Entry and prior to OR Exit. This includes bolus and infusion doses. Record "0 units" if no insulin was administered intraoperatively.

SEQ. #: 7410

Long Name: Blood Pressure Baseline (Pre-Anesthetic Induction) - Systolic

Short Name: PreAnesthBPSys

Definition: Indicate the most representative preoperative blood pressure upon arrival in the operating room.

The most representative initial blood pressure (systolic) should be recorded. This number may be an initial single recording or the average or median of a series of BP determinations. In all cases, the values should be recorded in the operating room prior to the induction of anesthesia.

Intent/Clarification:

Record the systolic blood pressure closest to, but prior to induction of anesthesia that is most representative of the patient's preoperative status. If the blood pressure closes to induction is debatably abnormal for the patient (erroneously high or low), then a median of the first five (5) blood pressures measurements by the automated record keeping system obtained after OR Entry may be used.

If this information is not available, leave blank.

SEQ. #: 7415

Long Name: Blood Pressure Baseline (Pre-Anesthetic Induction) - Diastolic

Short Name: PreAnesthBPDia

Definition: Indicate the most representative preoperative blood pressure upon arrival in the operating room.

The most representative initial blood pressure (diastolic) should be recorded. This number may be an initial single recording or the average or median of a series of BP determinations. In all cases, the values should be recorded in the operating room prior to the induction of anesthesia.

Intent/Clarification:

Record the diastolic blood pressure closest to, but prior to induction of anesthesia that is most representative of the patient's preoperative status.

If the blood pressure closes to induction is debatably abnormal for the patient, then a median of blood pressures obtained after OR Entry may be used. If the blood pressure closes to induction is debatably abnormal for the patient (erroneously high or low), then a median of the first five (5) blood pressures measurements by the automated record keeping system obtained after OR Entry may be used.

If this information is not available, leave blank.

SEQ. #: 7420

Long Name: Blood Pressure Baseline (Pre-Anesthetic Induction) - Mean

Short Name: PreAnesthBPMean

Definition: Indicate the most representative preoperative blood pressure upon arrival in the operating room.

The most representative initial blood pressure (mean) should be recorded. This number may be an initial single recording or the average or median of a series of BP determinations. In all cases, the values should be recorded in the operating room prior to the induction of anesthesia.

Intent/Clarification: Record the mean arterial pressure obtained from the arterial line closest to the induction of anesthesia. If the mean arterial pressure closes to induction is debatably abnormal for the patient, then a median of arterial pressures obtained after OR Entry may be used. If the arterial pressure closes to induction is debatably abnormal for the patient (erroneously high or low), then a median of the first five (5) arterial pressures measurements by the automated record keeping system obtained after OR Entry may be used.

If no mean arterial pressure is available, leave blank.

Do not capture mean cuff pressure (NBP).

SEQ. #: 7425

Long Name: Heart Rate Baseline (Pre-Anesthetic Induction)

Short Name: PreAnesthHR

Definition: Indicate the most representative preoperative heart rate upon arrival in the operating room.

The most representative initial heart rate should be recorded. This number may be an initial single recording or the average or median of a series of heart rate determinations. In all cases, the values should be recorded in the operating room prior to the induction of anesthesia. The source of heart rate should derive from the ECG monitor, since pulse rates derived from pulse oximetry/plethysmography or arterial tracings may underestimate the heart rate in tachyarrhythmias and other circumstances.

Intent/Clarification:

Record the heart rate closes to, but prior to induction of anesthesia, that is most representative of the patient's preoperative status. If the heart rate closes to induction is debatably abnormal for the patient, then a median of five (5) heart rates obtained after OR Entry may be used.

If no heart rate is available, leave blank.

SEQ. #: 7430

Long Name: Pulmonary Artery Catheter Used

Short Name: PACIntra

Definition: Indicate the preoperative or intraoperative placement of a pulmonary artery catheter (Swan-Ganz type-catheter).

Placement of a pulmonary artery catheter (PAC) in the preoperative or intraoperative period and use of this catheter during the intraoperative period.

Intent/Clarification:

Identify if a pulmonary artery catheter was placed pre or intra-operatively and used during the intraoperative period.

SEQ. #: 7435

Long Name: Core Temperature Source

Short Name: CoreTempSrc

Definition: Indicate the source of core temperature data used to guide cooling and/or rewarming during cardiac surgery.

Cardiac centers utilize various sites for measuring core temperature during cardiac procedures. These may include the esophageal, bladder, nasopharyngeal, pulmonary artery catheter thermistor, tympanic, or rectal sources. If more than one temperature is being recorded, the value selected as the core should be noted.

Intent/Clarification: Identify what source was used for determining the core temperature. This should coincide with data reported in the Operative section of the adult cardiac surgery data collection form.

SEQ. #: 7440

Long Name: Core Temperature Maximum

Short Name: CoreTempMax

Definition: Indicate the patient's highest core temperature during the procedure in degrees centigrade.

Intent/Clarification:

Indicate the patient's highest core temperature after the induction of anesthesia, prior to OR Exit.

SEQ. #: 7445

Long Name: Nitric Oxide Therapy Begun Intraoperatively

Short Name: NitricOxIntraop

Definition: Indicate the usage of inhaled nitric oxide.

Inhaled nitric oxide is used in the treatment of pulmonary hypertension and right ventricular failure. The intent is to capture the usage of inhaled nitric oxide during the cardiac surgical procedure. Do not record the usage of inhaled vasodilating substances other than nitric oxide in this data field.

Intent/Clarification:

Indicated if nitric oxide was used intraoperatively; after OR Entry but prior to OR Exit.

The Nitric Oxide (NO) machine is kept separate from the anesthesia machine and is often recoded by the Respiratory Therapist. The information is most likely found on the Respiratory Therapist record if not on the anesthesia record.

SEQ. #: 7450

Long Name: Total Crystalloid Administered by Anesthesia Care Team

Short Name: TotCrystAnesth

Definition: Indicate the total volume of intravenous crystalloid administered by the anesthesia care team. The data should be recorded in milliliters. Enter zero if no crystalloid used.

There is continuing controversy as to the risks and benefits of liberal or restrictive intravenous fluid regimens. Record the total volume of all crystalloid intravenous fluids administered by the anesthesia care team. Do not record any blood products in this data field.

Intent/Clarification:

Indicate if crystalloid fluids were administered in the OR by the anesthesia care team.

This does not include fluid administered by perfusion. Record in milliliters. Enter "0" if no crystalloid fluids were administered by anesthesia.

Common crystalloid fluids include 0.9% NaCl, Lactated Ringers, Plasmalyte and D5 ½ 0.9%Saline.

SEQ. #: 7455

Long Name: Total Synthetic Colloid Administered by Anesthesia Care Team

Short Name: TotColloidAnesth

Definition: Indicate the total volume of intravenous synthetic colloid fluid administered by the anesthesia care team. The data should be recorded in milliliters. Enter zero if no synthetic colloid used.

There is continuing controversy as to the risks and benefits of liberal or restrictive intravenous fluid regimens. Record the total volume of all synthetic colloid intravenous fluids administered by the anesthesia care team. Do not record any blood products in this data field.

Intent/Clarification:

Indicate if colloid fluids were administered in the OR by the anesthesia care team. **This does not include fluid administered by perfusion.** Record in milliliters. Enter "0" if no colloid fluids were administered by anesthesia.

Common colloid fluids are Hespan and Voluven.

SEQ. #: 7460

Long Name: Total Albumin Administered by Anesthesia Care Team

Short Name: TotAlbumAnesth

Definition: Indicate the total volume of intravenous human serum albumin fluid administered by the anesthesia care team. The data should be record in milliliters. Enter zero if no albumin used.

There is continuing controversy as to the risks and benefits of liberal or restrictive intravenous fluid regimens. Record the total volume of all human serum albumin fluid administered by the anesthesia care team. Do not record any blood products in this data field.

Intent/Clarification:

Indicate if Albumin was administered in the OR by the anesthesia care team. **This does not include administration by perfusion.** Record in milliliters. Enter "0" if no albumin was administered by anesthesia.

Albumin solutions include: Albumin 5%, Albumin 20% and Plasmanate 5%.

SEQ. #: 7470

Long Name: Intraoperative Glucose Trough Value

Short Name: GlucTroughIntraop

Definition: Indicate the trough value of intraoperative glucose in mg/dl.

Intraoperative glucose values vary widely in cardiac surgery. Administration of glucose containing fluids, stress, insulin, and glucorticoids may all affect intraoperative glycemic levels.

Intent/Clarification:

Indicate the patient's lowest intraoperative glucose level in mg/dL. Time frame is after induction of anesthesia and prior to OR Exit time.

SEQ. #: 7475

Long Name: Vasodilators used

Short Name: VasodillIntraop

Definition: Indicate the usage of intravenous vasodilating drugs administered by continuous infusion during the intraoperative phase of cardiac surgery.

Vasodilators are used commonly in cardiac surgical patients for the control of intraoperative hypertension and for afterload reduction to improve ventricular function. For the purposes of this data field, infusions of milrinone and pure vasodilating drugs, such as nitroglycerin, nitroprusside, and nicardipine, should be recorded.

Intent/Clarification:

Indicate if the patient received continuous infusion of vasodilating drugs intraoperatively. Do not include one-time dose.

Could include but not limited to: Apresoline/hydralazine, nitroglycerin, nitroprusside, nicardipine, Esmolol, and milrinone.

SEQ. #: 7476

Long Name: Intraoperative Processed EEG (BIS)

Short Name: IntraProcEEG

Definition: Indicate whether an intraoperative processed EEG (BIS) was monitored

Intent/Clarification:

Indicate if a processed EEG was utilized intraoperatively regardless if it was a BIS or other similar device.

SEQ. #: 7480

Long Name: Intraoperative Pre-procedure TEE Performed

Short Name: IntraOpPreTEE

Definition: Indicate whether intraoperative TEE was performed pre-procedure.

Intent/Clarification:

Indicate if an intraoperative TEE was performed after OR Entry time after induction, but prior to Incision time.

SEQ. #: 7485

Long Name: Pre-Procedure Left Ventricular Ejection Fraction Measured

Short Name: PreLVEFMeas

Definition: Indicate whether left ventricular ejection fraction was measured

Intent/Clarification:

Indicate if an ejection fraction was measured during the intraoperative TEE after OR Entry time after induction, but prior to Incision time.

This field is a child to Seq # 7480.

SEQ. #: 7490

Long Name: Left Ventricular Ejection Fraction Estimate

Short Name: PreLVEF

Definition: Indicate the estimate of Left Ventricular ejection fraction determined by intraoperative transesophageal echocardiography.

Enter a range of 1-99. If a percentage range is reported, report a whole number using the “mean” (i.e., 50-55% is reported as 53%). If a qualitative description is reported, code the mean value for that range; i.e., normal (50-70%) is coded as 60%.

- Hyperdynamic: >70%
- Normal: 50%–70% (midpoint 60%)
- Mild dysfunction: 40%–49% (midpoint 45%)
- Moderate dysfunction: 30%–39% (midpoint 35%)
- Severe dysfunction: <30%

Note: If no diagnostic report is in the medical record, a value documented in the medical record is acceptable.

Use the defining terms/percentages listed above to remain consistent with Adult Cardiac Database reporting.

SEQ. #: 7495

Long Name: Pre-Procedure Right Ventricular Function

Short Name: PreRVF_x

Definition: Indicate the estimate of RV function determined by intraoperative transesophageal echocardiography.

Intent/Clarification:

Use the Right Ventricular function obtained between OR Entry and incision time, but after induction.

SEQ. #: 7500

Long Name: Mitral Regurgitation

Short Name: PreMR

Definition: Indicate the degree of mitral valve regurgitation from intraoperative transesophageal echocardiography.

Enter the highest level recorded in the chart, i.e., worst performance level. “Moderately severe” should be coded as “severe”.

Intent/Clarification:

Use the degree of mitral valve regurgitation obtained between OR Entry and incision time, but after induction. Enter the highest level of regurgitation in the chart.

SEQ. #: 7505

Long Name: Mitral Stenosis

Short Name: PreMS

Definition: Indicate the degree of mitral valve stenosis from intraoperative transesophageal echocardiography.

Enter the highest level recorded in the chart, i.e., worst performance level. "Moderately severe" should be coded as "severe".

Intent/Clarification:

Use the degree of mitral valve regurgitation obtained between OR Entry and incision time, but after induction. Enter the highest level of mitral valve stenosis in the chart.

SEQ. #: 7510

Long Name: Aortic Regurgitation

Short Name: PreAR

Definition: Indicate the degree of aortic valve regurgitation from intraoperative transesophageal echocardiography.

Enter the highest level recorded in the chart, i.e., worst performance level. "Moderately severe" should be coded as "severe".

Intent/Clarification:

Use the degree of aortic valve regurgitation obtained between OR Entry and incision time, but after induction. Enter the highest level of aortic valve regurgitation in the chart.

SEQ. #: 7515

Long Name: Aortic Stenosis

Short Name: PreAS

Definition: Indicate the degree of aortic valve stenosis from intraoperative transesophageal echocardiography.

Enter the highest level recorded in the chart, i.e., worst performance level. "Moderately severe" should be coded as "severe".

Intent/Clarification:

Use the degree of aortic valve stenosis obtained between OR Entry and incision time, but after induction. Enter the highest level of aortic valve stenosis in the chart.

SEQ. #: 7520

Long Name: Aortic Valve Area Assessed

Short Name: PreAVAAssessed

Definition: Indicate whether the aortic valve areas was assessed from intraoperative transesophageal echocardiography.

Intent/Clarification:

Time frame is after OR Entry time.

SEQ. #: 7525

Long Name: Aortic Valve Area

Short Name: PreAVA

Definition: Indicate the aortic valve area from intraoperative transesophageal echocardiography.

Enter numeric value in square centimeters for aortic valve.

Intent/Clarification:

Report the aortic valve area obtained between OR Entry and incision time, but after induction. Answer in cm².

SEQ. #: 7530

Long Name: Tricuspid Regurgitation

Short Name: PreTR

Definition: Indicate the degree of tricuspid valve regurgitation from intraoperative transesophageal echocardiography.

Enter the highest level recorded in the chart, i.e., worst performance level. "Moderately severe" should be coded as "severe".

Intent/Clarification:

Enter the highest level of tricuspid valve regurgitation obtained between OR Entry and incision time, but after induction.

SEQ. #: 7535

Long Name: Patent Foramen Ovale

Short Name: PrePFO

Definition: Indicate the presence of patent foramen ovale diagnosed by intraoperative transesophageal echocardiography.

Intent/Clarification:

Indicated if a patent foramen ovale was identified on the intraoperative TEE. Time frame is between OR Entry and incision time, but after induction.

SEQ. #: 7540

Long Name: Ascending Aorta Assessed

Short Name: AscAoAssessed

Definition: Indicate whether the ascending aorta was assessed using TEE.

Intent/Clarification:

The ascending aorta includes the area from the aortic root to proximal of the innominate artery. Indicate if a TEE was performed intraoperatively to assess the ascending aorta. Time frame is after OR Entry time prior to incision, but after induction.

SEQ. #: 7545

Long Name: Maximal Ascending Aortic Diameter

Short Name: MxAscAo

Definition: Indicate the maximal diameter of ascending aorta as determined by intraoperative transesophageal echocardiography.

Indicate maximal diameter of ascending aorta in centimeters as determined by intraoperative transesophageal echocardiography.

Intent/Clarification:

Record the maximal diameter of the ascending aorta in centimeters using data obtained from an intraoperative TEE. Time frame is after OR Entry time prior to incision, but after induction.

SEQ. #: 7550

Long Name: Maximal Ascending Aortic Atheroma Thickness

Short Name: MxAscAoThick

Definition: Indicate the maximal ascending aortic atherosclerotic thickness as measured by intraoperative transesophageal echocardiography.

Indicate maximal thickness of ascending aorta plaque in millimeters as determined by intraoperative transesophageal echocardiography. If only intimal thickening and no plaque put numeric value of zero.

Intent/Clarification:

Time frame is after OR Entry time prior to incision, but after induction.

SEQ. #: 7555

Long Name: Ascending Aortic Atheroma Mobility

Short Name: AsAthMo

Definition: Indicate the ascending aortic atheroma mobility as measured by intraoperative transesophageal echocardiography.

Intent/Clarification:

Indicate if there was atheroma mobility within the ascending aorta. Time frame is after OR Entry time prior to incision, but after induction.

EQ. #: 7560

Long Name: Aortic Arch Visualized

Short Name: AoArcVis

Definition: Indicate whether the aortic arch was visualized.

Intent/Clarification:

Indicate if an intraoperative TEE was performed that assessed the aortic arch. The aortic arch is normally located between the innominate artery and left subclavian artery. Time frame is after OR Entry time prior to incision, but after induction.

SEQ. #: 7565

Long Name: Maximal Aortic Arch Atheroma Thickness

Short Name: MxArcAth

Definition: Indicate the maximal aortic arch atherosclerotic thickness as measured by intraoperative transesophageal echocardiography.

Indicate maximal thickness of aortic arch plaque in millimeters as determined by intraoperative transesophageal echocardiography. If only intimal thickening and no plaque put numeric value of zero.

Intent/Clarification:

Time frame is after OR Entry time prior to incision, but after induction.

SEQ. #: 7570

Long Name: Aortic Arch Atheroma Mobility

Short Name: ArcAthMo

Definition: Indicate the aortic arch atheroma mobility as measured by pre-CPB intraoperative transesophageal echocardiography.

Intent/Clarification:

Indicate if aortic arch atheroma mobility was noted on the intraoperative TEE. Time frame is after OR Entry time prior to incision, but after induction.

SEQ. #: 7575

Long Name: Cardiopulmonary Bypass Used

Short Name: CPBUsed

Definition: Indicate whether cardiopulmonary bypass was used.

Intent/Clarification:

Indicate if the patient was placed on cardiopulmonary bypass for any portion of the procedure.

SEQ. #: 7580

Long Name: Retrograde Autologous Priming of CPB Circuit

Short Name: RetrAutolPrim

Definition: Indicate whether retrograde autologous priming was used by the cardiopulmonary perfusion team prior to the onset of cardiopulmonary bypass.

Retrograde autologous priming is technique used by cardiopulmonary perfusionists to minimize hemodilution and hypotension during onset of cardiopulmonary bypass.

Intent/Clarification:

Indicate if retrograde autologous priming was used by Perfusion. This information can usually be obtained in the perfusion record.

SEQ. #: 7585

Long Name: Total Fluids Crystalloid Administered by Perfusion Team

Short Name: TotCrystPerf

Definition: Indicate the total volume of intravenous crystalloid fluids administered by cardiopulmonary perfusion team. The data should be record in milliliters. Enter zero if fluid crystalloid not used by perfusion team.

There is continuing controversy as to the risks and benefits of liberal or restrictive intravenous fluid regimens. Record the total of all crystalloid intravenous fluids given by the cardiopulmonary perfusion team. Do not record any blood products in this data field.

Intent/Clarification:

Record the entire amount of crystalloid fluids administered intravenously by the perfusion team as recorded on the perfusion record. Do not include amount given by anesthesia, this is captured in SEQ. #7450. If input and output amounts are listed, record the input amount.

Common crystalloid fluids include 0.9% NaCl, Lactated Ringers, and D5 ½-0.9% NaCl.

SEQ. #: 7590

Long Name: Total Synthetic Colloid Administered by Perfusion Team

Short Name: TotColloidPerf

Definition: Indicate the total volume of intravenous synthetic colloid fluids (of any concentration) administered by the cardiopulmonary perfusion team. The data should be recorded in milliliters. Enter zero if synthetic colloid not administered by perfusion team.

There is continuing controversy as to the risks and benefits of liberal or restrictive intravenous fluid regimens. Record the total of all synthetic colloid intravenous fluids given by the cardiopulmonary perfusion team. Synthetic colloids of all concentrations and substitution ratios should be included, Do not record any blood products in this data field.

Intent/Clarification:

Record the entire amount of colloid fluids administered intravenously by the perfusion team as recorded on the perfusion record. Do not include amount given by anesthesia, this is captured in SEQ. #7455. If input and output amounts are listed, record the input amount.

Common colloid fluids used in the OR are Hespan and Voluven.

SEQ. #: 7595

Long Name: Total Albumin Administered by Perfusion Team

Short Name: TotAlbumPerf

Definition: Indicate the total volume of intravenous human serum albumin fluids (of any concentration) administered by the cardiopulmonary perfusion team. The data should be recorded in milliliters. Enter zero if albumin not administered by perfusion team.

There is continuing controversy as to the risks and benefits of liberal or restrictive intravenous fluid regimens. Record the total of all human serum albumin intravenous fluids given by the cardiopulmonary perfusion team. Albumin-containing fluids of all concentrations should be included. Do not record any blood products in this data field.

Intent/Clarification:

Record the entire amount of albumin administered intravenously by the perfusion team. Do not include amount given by anesthesia, this is captured in SEQ. #7460.

SEQ. #: 7600

Long Name: Hemofiltration Volume Removed by Perfusion Team **Short Name:** HemofilPerf

Definition: Indicate the total volume of ultrafiltrate removed by the cardiopulmonary perfusion team during cardiopulmonary bypass and during modified ultra-hemofiltration post-CPB. Record the data in milliliters.

Hemofiltration is used to concentrate the red blood cells and plasma proteins in the circulation during and immediately following CPB.

Intent/Clarification:

Indicate the total volume of fluid removed by hemofiltration intraoperatively after the initiation of the initial cardiopulmonary bypass as record on the perfusion record. Record amount in millimeters. Time frame is at the start of the initial cardiopulmonary bypass to admission to the ICU.

SEQ. #: 7605

Long Name: Inotropes used to wean from CPB

Short Name: InotropWeanCPB

Definition: Indicate the usage of inotropic drug infusions to facilitate weaning from cardiopulmonary bypass. For this data field, any drug infusion with inotropic properties, including catecholamines, phosphodiesterase inhibitors, and calcium sensitizers, should be recorded.

Inotropic drugs infusions are used routinely or as required in many cardiac surgical patients during the process of weaning from CPB. Record all usage of drugs with positive inotropic effect, including epinephrine, norepinephrine, dopamine, dobutamine, levosimendan, and milrinone.

Intent/Clarification:

Indicate if inotropes were used to facilitate the weaning process from cardiopulmonary bypass. Select "Yes" if any drug with inotropic property was administered during the weaning process.

Inotropic drugs increase the pumping effect of the heart muscle, making the heart pump stronger. Common inotropic drugs include epinephrine, norepinephrine, dopamine, dobutamine, levosimendan, and milrinone. This also includes drugs with inotropic properties such as catecholamines, phosphodiesterase inhibitors, and calcium sensitizers.

If timing is unclear, obtain clarification regarding timing of the weaning process from the Cardiothoracic Anesthesiology team at your facility.

SEQ. #: 7610

Long Name: Vasopressors used to wean from CPB

Short Name: VasopWeanCPB

Definition: Indicate the usage of vasoconstrictive drugs to facilitate weaning from cardiopulmonary bypass. For this data field, any drug infusion at a dosage range with clinically vasoconstrictive properties, including catecholamines and pure vasoconstrictors, should be recorded.

Low systemic vascular resistance (a.k.a. vasoplegia) is a common condition during cardiopulmonary bypass that may be related to preoperative vasodilating drugs or certain antiarrhythmic drugs. Include purely vasoconstrictive drugs. Also record usage of drugs with inotropic effects that have vasoconstrictive properties in higher doses, such as dopamine and epinephrine.

Intent/Clarification:

Indicate if vasopressors were used to facilitate the weaning process from cardiopulmonary bypass. Select "Yes" for any drug with vasoconstrictive property that was administered, this includes inotropic drugs (such as epinephrine and dopamine) that can be dosed at vasoconstrictive levels or pure vasoconstrictors such as

vasopressin or phenylephrine. Vasoconstrictive drugs constrict the blood vessels raising blood pressure.

Common vasoconstrictor drugs include dopamine, epinephrine, neosynephrine/phenylephrine, norepinephrine (Levophed) and vasopressin.

If timing is unclear, obtain clarification regarding timing of the weaning process from the Cardiothoracic Anesthesiology team at your facility.

SEQ. #: 7615

Long Name: Intraoperative Post-procedure TEE Performed

Short Name: IntraOpPostTEE

Definition: Indicate whether intraoperative TEE was performed post-procedure.

Intent/Clarification:

Indicate if a transesophageal echocardiogram was performed post-procedure intraoperatively. Time frame is after weaning from cardiopulmonary bypass to OR Exit time

SEQ. #: 7620

Long Name: Systolic Anterior Motion of Mitral Valve

Short Name: PostSAM

Definition: Indicate the presence of systolic anterior motion (SAM) of the mitral valve as determined by intraoperative transesophageal echocardiography prior to chest closure.

Choose Yes for any SAM between weaning from CPB and chest closure.

Intent/Clarification:

If a post-procedure TEE was performed, indicate if systolic anterior motion of the mitral valve was noted. Choose "Not assessed" if a post-procedure TEE was performed but systolic anterior motion of the mitral valve was not documented. Time frame is after weaning from cardiopulmonary bypass to OR Exit time.

SEQ. #: 7625

Long Name: Return to CPB for Echo-Related Diagnosis

Short Name: RetCPBEch

Definition: Indicate whether surgical revision was performed based on post procedure intraoperative TEE.

Intent/Clarification:

Indicate if the patient had to be placed back on cardiopulmonary bypass for a surgical revisit as a result from findings on the post-procedure TEE prior to OR Exit time.

SEQ. #: 7630

Long Name: Post-Procedure Left Ventricular Ejection Fraction Measured

Short Name: PostLVEFMeas

Definition: Indicate whether left ventricular ejection fraction was measured post-procedure by intraoperative transesophageal echocardiography.

Intent/Clarification:

Time frame for TEE is the closest time before OR Exit time after final discontinuation of cardiopulmonary bypass time.

SEQ. #: 7635

Long Name: Post-Procedure Left Ventricular Ejection Fraction Estimate **Short Name:** PostLVEF

Definition: Indicate the post-procedure estimate of left ventricular ejection fraction determined by intraoperative transesophageal echocardiography.

Enter a range of 1-99. If a percentage range is reported, report a whole number using the "mean" (i.e., 50-55% is reported as 53%). If a qualitative description is reported, code the mean value for that range; i.e., normal (50-70%) is coded as 60%.

- Hyperdynamic: >70%
- Normal: 50%–70% (midpoint 60%)
- Mild dysfunction: 40%–49% (midpoint 45%)
- Moderate dysfunction: 30%–39% (midpoint 35%)
- Severe dysfunction: <30%

Note: If no diagnostic report is in the medical record, a value documented in the medical record is acceptable.

Use the defining terms/percentages listed above to remain consistent with Adult Cardiac Database reporting.

SEQ. #: 7640

Long Name: Post-Procedure Right Ventricular Function

Short Name: PostRVFx

Definition: Indicate the post-procedure estimate of RV function determined by intraoperative transesophageal echocardiography.

Intent/Clarification:

Choices are normal, mild dysfunction, moderate dysfunction, severe dysfunction, and not assessed. If a range is reported (i.e. mild-moderate) choose the highest range reported. Choose "unknown" if a post-procedure TEE is performed, but right ventricular dysfunction is not documented. Time frame for TEE is the closest time before OR Exit time after final discontinuation of bypass time.

SEQ. #: 7641

Long Name: Intraoperative Cardiac Arrest Related To Anesthesia Care

Short Name: IntraCardArr

Definition: Indicate whether there was a cardiac arrest related to anesthesia care

Intent/Clarification:

Indicate if the patient's heart arrested post-procedure, intraoperatively. Time frame is from induction to ICU arrival time.

SEQ. #: 7645

Long Name: Patient Died Within The OR

Short Name: ORDeath

Definition: Indicate whether the patient died within the OR.

Intent/Clarification:

Time frame is from OR Entry to OR Exit time.

SEQ. #: 7650

Long Name: Core Temperature Upon Entry To ICU/PACU Measured

Short Name: PostTempMeas

Definition: Indicate whether the core temperature was measured upon initial arrival in the ICU/PACU following cardiac surgery.

Intent/Clarification:

Indicate if the core temperature was measured upon arrival to ICU/PACU immediately following cardiac surgery. Core temperature locations include: bladder, rectum, pulmonary artery, esophageal, nasopharyngeal, and tympanic.

SEQ. #: 7655

Long Name: Core Temperature Upon Entry To ICU/PACU

Short Name: PostCoreTemp

Definition: Indicate the core temperature in degrees Centigrade upon initial arrival in the ICU/PACU following cardiac surgery.

The intent is to capture the initial documented core temperature in the intensive care unit, as per the normal routine for core temperature monitoring in the ICU/PACU.

Intent/Clarification:

Document the initial CORE temperature in degrees Celsius upon arrival to the ICU/PACU following cardiac surgery.

SEQ. #: 7660

Long Name: Postoperative INR Measured

Short Name: PostINRMeas

Definition: Indicate whether the International normalized ratio (INR) was measured upon initial arrival in the ICU/PACU following cardiac surgery.

Intent/Clarification:

Document if an initial International Normalized Ratio (INR) was measured upon arrival to the ICU/PACU following cardiac surgery. This lab is usually part of the Prothrombin test (PT/INR).

SEQ. #: 7665

Long Name: First Postoperative INR

Short Name: PostINR

Definition: Indicate the first international normalized ratio (INR) value upon initial arrival in the ICU/PACU following cardiac surgery.

INR is the standard unit used to report the result of a prothrombin (PT) test. The hospital laboratory report should be accessed first when coding this variable. If this is unavailable, then additional source documents may be referenced for lab results.

Intent/Clarification:

Record the first INR value upon arrival to ICU/PACU following cardiac surgery.

FAQ September 2017: What is the timeframe for this component? I have lab results that are 1 hour and 45 min from surgery. Is this too long of a timeframe?

Answer: Laboratory values should be drawn within one hour of arrival in the ICU/PACU.

SEQ. #: 7670

Long Name: WBC Upon Entry To ICU/PACU Measured

Short Name: PostWBCMeas

Definition: Indicate whether the white blood cell count was measured upon initial arrival in the ICU/PACU following cardiac surgery.

Intent/Clarification:

Document if an initial white blood cell count (WBC) was measure upon arrival to the ICU/PACU. This is usually part of the complete blood count (CBC) test.

FAQ September 2017: What is the timeframe for this component? I have lab results that are 1 hour and 45 min from surgery. Is this too long of a timeframe?

Answer: Laboratory values should be drawn within one hour of arrival in the ICU/PACU.

SEQ. #: 7675

Long Name: WBC Upon Entry To ICU/PACU

Short Name: PostWBC

Definition: Indicate the first white blood cell count upon initial arrival in the ICU/PACU following cardiac surgery.

White Blood Cells (leukocytes) are part of the body's immune defense and are often elevated in the presence of infection. The hospital laboratory report should be accessed first when coding this variable. If this is unavailable, then additional source documents may be referenced for lab results.

Intent/Clarification:

Record the first WBC value upon admission to ICU/PACU following cardiac surgery.

FAQ September 2017: What is the timeframe for this component? I have lab results that are 1 hour and 45 min from surgery. Is this too long of a timeframe?

Answer: Laboratory values should be drawn within one hour of arrival in the ICU/PACU.

SEQ. #: 7680

Long Name: Platelets Upon Entry To ICU/PACU Measured

Short Name: PostPltMeas

Definition: Indicate whether the platelet count was measured upon initial arrival in the ICU/PACU following cardiac surgery.

Intent/Clarification:

Document if an initial platelet count (PLT) was measure upon arrival to the ICU/PACU. This is usually part of the complete blood count (CBC) test.

FAQ September 2017: What is the timeframe for this component? I have lab results that are 1 hour and 45 min from surgery. Is this too long of a timeframe?

Answer: Laboratory values should be drawn within one hour of arrival in the ICU/PACU.

SEQ. #: 7685

Long Name: Platelets Upon Entry To ICU/PACU

Short Name: PostPlt

Definition: Indicate the first platelet count upon initial arrival in the ICU/PACU following cardiac surgery.

Platelets are a blood component instrumental in clot formation. The hospital laboratory report should be accessed first when coding this variable. If this is unavailable, then additional source documents may be referenced for lab results.

Intent/Clarification:

Record the first platelet count upon admission to ICU/PACU following cardiac surgery.

FAQ September 2017: What is the timeframe for this component? I have lab results that are 1 hour and 45 min from surgery. Is this too long of a timeframe?

Answer: Laboratory values should be drawn within one hour of arrival in the ICU/PACU.

SEQ. #: 7690

Long Name: Hematocrit Upon Entry To ICU/PACU Measured

Short Name: PostHCTMeas

Definition: Indicate whether the hematocrit value was measured upon initial arrival in the ICU/PACU following cardiac surgery.

Intent/Clarification:

Document if an initial hematocrit level (HCT) was measure upon arrival to the ICU/PACU. This is usually part of the complete blood count (CBC) test or hemoglobin/hematocrit (H/H) test.

SEQ. #: 7695

Long Name: Hematocrit Upon Entry To ICU/PACU

Short Name: PostHCT

Definition: Indicate the first hematocrit value upon initial arrival in the ICU/PACU following cardiac surgery.

Hct, Hematocrit, is the proportion of red cells in the blood. The hospital laboratory report should be accessed first when coding this variable. If this is unavailable, then additional source documents may be referenced for lab results.

Intent/Clarification:

Record the first hematocrit (Hct) level upon admission to ICU/PACU following cardiac surgery.

FAQ September 2017: Is there a timeframe for the results for this component? I have lab results that are 1 hour and 45 min from surgery. Is this too long of a timeframe?

Answer: Laboratory values should be drawn within one hour of arrival in the ICU/PACU.

SEQ. #: 7696

Long Name: Fibrinogen Upon Entry To ICU/PACU Measured

Short Name: PostFibrinMeas

Definition: Indicate whether fibrinogen was measured upon entry to ICU/PACU

Intent/Clarification:

Document if an initial fibrinogen level was measured upon arrival to the ICU/PACU.

SEQ. #: 7697

Long Name: Fibrinogen Upon Entry To ICU/PACU

Short Name: PostFibrin

Definition: Indicate the fibrinogen level upon entry to ICU/PACU

Intent/Clarification:

Record the first fibrinogen level upon admission to the ICU/PACU following cardiac surgery.

FAQ September 2017: Is there a timeframe for the results for this component? I have lab results that are 1 hour and 45 min from surgery. Is this too long of a timeframe?

Answer: Laboratory values should be drawn within one hour of arrival in the ICU/PACU.

SEQ. #: 7700

Long Name: Lactate Upon Entry To ICU/PACU Measured

Short Name: PostLactMeas

Definition: Indicate whether the lactate value was measured upon initial arrival in the ICU/PACU following cardiac surgery.

Intent/Clarification:

Document if an initial Lactate level (Lactic Acid) was measure upon arrival to the ICU/PACU.

SEQ. #: 7705

Long Name: Lactate Upon Entry To ICU/PACU

Short Name: PostLact

Definition: Indicate the value of lactate in **mg/dl mmol/L** upon initial arrival in the ICU/PACU following cardiac surgery. Do not record missing data as a zero value.

Serum lactate is a marker for the duration and severity of malperfusion during critical states. The magnitude of serum lactate has been associated with mortality and adverse outcomes.

Intent/Clarification:

Record the first lactate (lactic acid) level upon admission to the ICU/PACU following cardiac surgery.

FAQ September 2017: Is there a timeframe for the results for this component? I have lab results that are 1 hour and 45 min from surgery. Is this too long of a timeframe?

Answer: Laboratory values should be drawn within one hour of arrival in the ICU/PACU.

SEQ. #: 7710

Long Name: Postop Infusion: Dexmedetomidine

Short Name: DexPost

Definition: Indicate the use of dexmedetomidine infusion after surgery.

Any use of dexmedetomidine infusion during the postoperative period, after transport to the ICU/PACU.

Intent/Clarification:

Indicate if dexmedetomidine was administered after admission to the ICU/PACU following cardiac surgery. Time frame is from OR Exit to Discharge.

SEQ. #: 7715

Long Name: Postop Infusion: Propofol

Short Name: PropPost

Definition: Indicate the use of propofol infusion after surgery.

Any use of a Propofol infusion during the postoperative period, after transport to the ICU/PACU.

Intent/Clarification:

Indicate if the patient received a Propofol infusion after admission to the ICU/PACU following cardiac surgery. Time frame is from OR Exit to Discharge.

This does not include bolus doses.

SEQ. #: 7720

Long Name: Postoperative Delirium

Short Name: PostopDel

Definition: Indicate whether the patient experienced postoperative delirium.

Postoperative altered mental state such as loss of memory and cognitive ability, personality changes, inability to concentrate, or lethargy, without actual evidence of stroke or coma.

Intent/Clarification:

Indicate if the patient experienced postoperative delirium as evidenced by change in mental status (including memory loss, personality changes, lethargy, and changes in cognitive ability) without evidence of a stroke or coma. Refer to physician documentation for diagnosis. Time frame is from OR Exit to Discharge.

Definition of Post-operative Delirium

Post-operative delirium is a state of global brain dysfunction occurring after a surgical procedure, the diagnosis of which is made by establishing:

- An acute disturbance in level of arousal (may be thargy-stupor or hypervigilance-agitation) and an acute disturbance in cognition.
 - Identifying these disturbances as representing an acute change in the patient's baseline level of arousal and cognition requires the establishment of baseline functioning in these areas from corroborative sources including family, friends, and caregivers. *Note:* Even patients with poor baseline levels of cognitive function (i.e. pre-existing Dementia) can develop superimposed delirium.
 - The hallmark cognitive changes associated with delirium is a disturbance in attention (reduced ability to direct, focus, sustain, or shift attention) and awareness (reduced orientation to environment).
- These changes must develop over a short period of time (usually hours to a few days).
- These changes in cognition and level of arousal must demonstrate a pattern of fluctuation in severity during course of the day (i.e. there can be intervening periods of lucidity).
- Additional cognitive disturbances which may manifest during an episode of Delirium:
 - Memory deficits
 - Disorientation
 - Language
 - Visuospatial ability
- Additional behavioral disturbances which may manifest during an episode of delirium:
 - Changes in sleep-wake cycle
 - Hostility
 - Verbal and physical aggression
 - Unintentional self-harm (i.e. self-extubation, removal of catheters, falling out of bed)
 - Uncooperativeness with care
 - Euphoria
 - Hallucinations (visual or auditory)
 - Delusions (typically paranoid)

- Disorganized thinking

SEQ. #: 7725

Long Name: Heparin-Induced Thrombocytopenia (Postop Dx)

Short Name: PostHITAnti

Definition: Indicate whether Heparin Induced Thrombocytopenia, HIT, is confirmed by antibody testing.

Heparin induced thrombocytopenia (HIT) can be defined as any clinical event best explained by platelet factor 4 (PF4)/heparin-reactive antibodies ('HIT antibodies') in a patient who is receiving, or who has recently received heparin. Thrombocytopenia is the most common 'event' in HIT and occurs in at least 90% of patients, depending upon the definition of thrombocytopenia. A very small proportion of patients with HIT develop thrombosis. Alternative (nonheparin) anticoagulant therapy reduces the risk of subsequent thrombosis.

Intent/Clarification:

Indicate if the patient experienced heparin induced thrombocytopenia (HIT) postoperatively. This is evidenced by the presence of HIT antibodies found via specific laboratory test. Consult with your laboratory to determine the test that your facility uses and the number of positive test used to diagnose a patient. Depending upon the test, some facilities require three positive results before confirming the diagnosis. This is sometimes referred to as being "HITA positive" in documentation. Time frame is from OR Exit to Discharge.

SEQ. #: 7730

Long Name: Pain Score POD #3


Short Name: PainScorePOD3

Definition: Indicate the pain score on postoperative day #3 (Integer Rating Scale).

Highest pain score on postoperative day #3 on the 0-10 integer scale, if recorded, or record score as missing.

Intent/Clarification:

Record the highest pain score using the integer scale from 0-10 on postoperative day 3. With a score of "0" indicating no pain and a score of "10" indicating the worse possible pain ever imagined. If the patient was evaluated on a non-numerical scale use the corresponding answer related to the 1-10 scale.



SEQ. #: 7735

Long Name: Pain Score Hospital Discharge

Short Name: PainScoreDisch

Definition: Indicate the pain score on day of discharge (Integer Rating Scale).

Highest pain score recorded on day of discharge on the 0-10 integer scale, if recorded, or record score as missing.

Intent/Clarification:

Record the highest pain score using the integer scale from 0-10 on the day of discharge from the hospital inpatient stay. With a score of "0" indicating no pain and a score of "10" indicating the worse possible pain ever imagined. If the patient was evaluated on a non-numerical scale use the corresponding answer related to the 1-10 scale.

