

## Table of Contents

Introduction.....	2
1. Administrative .....	2
2. Demographics .....	5
3. Noncardiac Congenital Anatomic Abnormalities .....	21
4. Chromosomal Abnormalities .....	22
5. Syndromes .....	23
6. Hospitalization .....	24
7. Preoperative Factors .....	28
8. Diagnosis .....	30
9. Procedures .....	32
10. Procedure-Specific Factors.....	37
11. Operative.....	58
12. CABG Procedures .....	120
13. Valve Procedures .....	121
14. VAD Procedures.....	133
15. Complications.....	139
16. Discharge/Readmission.....	143
17. Patient Process Measures .....	149
18. Anesthesia Administrative .....	157
19. Anesthesia Preoperative .....	159
20. Anesthesia Monitoring.....	164
21. Anesthesia Anesthetic Technique .....	174
22. Anesthesia Airway.....	183
24. Anesthesia Intraoperative Pharmacology.....	186
25. Anesthesia Pharmacology On Arrival To ICU/PACU.....	187
26. Anesthesia ICU/PACU Care .....	188
27. Anesthesia Adverse Events .....	192
28. STS Temporary Fields .....	193

## Introduction

*This manual is intended to clarify field definition and intent. This document contains the most up to date instructions for v. 3.3 data abstraction. Do not refer to old manuals or other data definitions. Please review this document prior to submitting clinical questions. FAQs will be added to the document in red to provide additional examples and clarification. Please do not print this document since it will change frequently. Using the web version will ensure that you have the most up to date information. Occasionally there may be changes or important information that will be highlighted here and will be also included in STS Database Newsletters.*

*Use the Ctrl + F function to search for a number or term of interest.*

### *General Information:*

*The STS data collection forms should be held for two years. If you only collect data directly to the software you are not required to create data collection forms to save.*

*For all questions where the choices include “no” and “unknown”: When a history and physical or a consultation exists in the medical record and the values are not specifically addressed in the documentation, code no. Unknown should be coded only in the circumstance where no clinical documentation exists and the patient cannot give history and supportive documentation.*

## 1. Administrative

SeqNo: 10

Long Name: Participant ID

Short Name: ParticID

Database Table Name: Operations

Data Source: User or Automatic

Format: Text Data Length: 5

Definition: Participant ID is a unique number assigned to each database participant by the STS. A database participant is defined as one entity that signs a Participation Agreement with the STS, submits one data file to the harvest, and gets back one report on their data. The participant ID must be entered into each record.

Each participant's data, if submitted to harvest, must be in one data file. If one participant keeps their data in more than one file (e.g., at two sites), then the participant must combine them back into one file for harvest submission.

If two or more participants share a single purchased software, and enter cases into one database, then the data must be extracted into two different files, one for each participant ID, with each record having the correct participant ID number.

SeqNo: 20

Long Name: STS Data Version

Short Name: DataVrsn

Database Table Name: Operations

Data Source: Automatic  
Format: Text Data Length: 8  
Definition: Version number of the STS Data Specifications/Dictionary, to which each record conforms. It will identify which fields should have data, and what are the valid data for each field. This must be entered into the record automatically by the software at the time the record is created.

---

SeqNo: 30  
Long Name: On-Demand Files Version Number  
Short Name: OnDemandVrsn  
Database Table Name: Operations  
Data Source: Automatic  
Format: Text  
Definition: The version number of the On-Demand lists in use at the time this data record was created or edited. The value is inserted into the record at the time the record is created or is modified by the user. The version numbers will be specified by the STS.

---

SeqNo: 40  
Long Name: Software Vendor Identifier  
Short Name: VendorID  
Database Table Name: Operations  
Data Source: Automatic  
Format: Text Data Length: 8  
Definition: Identifying code (assigned by STS) given to identify software vendor (up to 8 characters). Vendors should use standard name identification across sites. Changes to Software Vendor Identifier must be reported to the STS.

---

SeqNo: 50  
Long Name: Software Version  
Short Name: SoftVrsn  
Database Table Name: Operations  
Data Source: Automatic  
Format: Text Data Length: 20  
Definition: Vendor's software product name and version number identifying the software which created this record. Vendor controls the value in this field.

---

SeqNo: 60  
Long Name: Operation Table Record Identifier  
Short Name: OperationID  
Database Table Name: Operations

---

Data Source: Automatic  
Format: Text  
Definition: An arbitrary, unique value generated by the software that permanently identifies each operation record in the participant's database. The value of the identifier is a combination of a code assigned to the software developer by the STS, and a value generated by the software to create a unique value. Once assigned to a record, this number can never be changed or reused. The data warehouse will use this value to communicate issues about individual records with the participant. This field is the primary key that links this record with the associated records in the Diagnosis, Risk Factors, Preoperative Factors, Procedures, Complications, Anesthesia Adverse Events, Preoperative Medications, Intraoperative Pharmacology, and ICU Pharmacology tables.

---

SeqNo: 70  
Long Name: Operations Link to Demographics Table  
Short Name: PatID  
Database Table Name: Operations  
Data Source: Automatic  
Format: Text  
Definition: An arbitrary, unique value generated by the software that permanently identifies each patient demographic record in the participant's database. This field is the foreign key that links this record with the associated record in the Demographics table.

---

SeqNo: 81  
Long Name: Patient Participating In STS-Related Clinical Trial  
Short Name: ClinTrial  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate which, if any, STS-related clinical trial in which the patient is participating. The STS will assign a code to each clinical trial as they begin collecting data.

---

SeqNo: 82  
Long Name: Patient Participating In STS-Related Clinical Trial - Patient ID  
Short Name: ClinTrialPatID  
Database Table Name: Operations  
Data Source: User  
Format: Text  
Definition: Indicate the patient identifier used to identify the patient in the clinical trial.

Parent Long Name: Patient Participating In STS-Related Clinical Trial  
Parent Short Name: ClinTrial  
Parent Value(s): Is Not "None" And Is Not Missing

---

## 2. Demographics

SeqNo: 90

Long Name: Demographics Table Patient Identifier

Short Name: PatID

Database Table Name: Demographics

Data Source: Automatic

Format: Text

Definition: An arbitrary value (not a recognizable ID like Social Security Number or Medical Record Number) that uniquely and permanently identifies each patient. The value of the identifier is a combination of a code assigned to the software developer by the STS, and a value generated by the software to create a unique value. Once assigned to a patient, this can never be changed or reused.

This field is the primary key that links this demographics record with the associated records in the Non-Cardiac Abnormalities, Noncardiac Congenital Anatomic Abnormalities, Chromosomal Abnormalities, and Syndromes tables.

---

SeqNo: 100

Long Name: Demographics Table Data Version

Short Name: DemogDataVrsn

Database Table Name: Demographics

Data Source: Automatic

Format: Text

Definition: Version number of the STS Data Specifications/Dictionary, to which this Demographics record conforms as assigned by the software. This value will determine which fields should have data and what are the valid data for each field. This must be entered into the record automatically by the software at the time the record is created. See Software Specifications document for description of how this value can be modified after the record was created.

SeqNo: 110

Long Name: Patient National Identification (Social Security Number)

Short Name: PatNationalID

Database Table Name: Demographics

Data Source: User

Format: Text

Definition: Indicate the patient's Social Security Number (SSN). Although this is the Social Security Number in the USA, other countries may have a different National Patient Identifier Number. For example in Canada, this would be the Social Insurance Number.

---

This field should be collected in compliance with state/local privacy laws.

---

SeqNo: 120

Long Name: Medical Record Number

Short Name: MedRecN

Database Table Name: Demographics

Data Source: User

Format: Text Data Length: 25

Definition: Indicate the patient's medical record number at the hospital where surgery occurred.

This field should be collected in compliance with state/local privacy laws.

---

SeqNo: 140

Long Name: Patient Last Name

Short Name: PatLName

Database Table Name: Demographics

Data Source: User

Format: Text Data Length: 50

Definition: Indicate the patient's last name documented in the medical record.

This field should be collected in compliance with state/local privacy laws.

---

SeqNo: 150

Long Name: Patient First Name

Short Name: PatFName

Database Table Name: Demographics

Data Source: User

Format: Text Data Length: 50

Definition: Indicate the patient's first name documented in the medical record.

This field should be collected in compliance with state/local privacy laws.

---

SeqNo: 170

Long Name: Patient Middle Name

Short Name: PatMName

Database Table Name: Demographics

Data Source: User

Format: Text Data Length: 50

Definition: Indicate the patient's middle name or middle initial as documented in the medical record. Leave "blank" if no middle name.

This field should be collected in compliance with state/local privacy laws.

---

---

SeqNo: 180  
Long Name: Patient's Region  
Short Name: PatRegion  
Database Table Name: Demographics  
Data Source: User  
Format: Text Data Length: 50  
Definition: Indicate the region of the country (i.e., state or province) in which the patient permanently resides at time of admission.

---

SeqNo: 190  
Long Name: Patient's Postal Code  
Short Name: PatPostalCode  
Database Table Name: Demographics  
Data Source: User  
Format: Text Data Length: 20  
Definition: Indicate the ZIP Code of the patient's residence. Outside the USA, this data may be known by other names such as Postal Code.

This field should be collected in compliance with state/local privacy laws.

---

SeqNo: 201  
Long Name: Patient's Country  
Short Name: PatientCountry  
Database Table Name: Demographics  
Data Source: User  
Format: Text  
Definition: Indicate the patient's country of residence at time of admission.

This field should be collected in compliance with state/local privacy laws.

---

SeqNo: 202  
Long Name: Birth Location Is Known  
Short Name: BirthLocKnown  
Database Table Name: Demographics  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the location (city, state, country) of the patient's birth is known.

---

SeqNo: 210

---

Long Name: City of Birth  
Short Name: BirthCit  
Database Table Name: Demographics  
Data Source: User  
Format: Text  
Definition: Indicate the city in which the patient was born.

Parent Long Name: Birth Location Is Known  
Parent Short Name: BirthLocKnown  
Parent Value(s): = "Yes"

---

SeqNo: 220  
Long Name: Birth Region  
Short Name: BirthSta  
Database Table Name: Demographics  
Data Source: User  
Format: Text Data Length: 50  
Definition: Indicate the region of the country (i.e., state or province) in which the patient was born.

Parent Long Name: Birth Location Is Known  
Parent Short Name: BirthLocKnown  
Parent Value(s): = "Yes"

---

SeqNo: 231  
Long Name: Country of Birth  
Short Name: BirthCountry  
Database Table Name: Demographics  
Data Source: User  
Format: Text  
Definition: Indicate the country in which patient was born.

This field should be collected in compliance with state/local privacy laws.

Parent Long Name: Birth Location Is Known  
Parent Short Name: BirthLocKnown  
Parent Value(s): = "Yes"

---

SeqNo: 232  
Long Name: Mode of Delivery Known  
Short Name: DelivModeKnown  
Database Table Name: Demographics  
Data Source: User

---



Format: Text (categorical values specified by STS)  
Definition: Indicate whether the mode of delivery is known.

---

SeqNo: 233  
Long Name: Mode of Delivery  
Short Name: DelivMode  
Database Table Name: Demographics  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the mode of delivery.

Parent Long Name: Mode of Delivery Known  
Parent Short Name: DelivModeKnown  
Parent Value(s): = "Yes"

---

SeqNo: 234  
Long Name: Mother's Gravidity And Parity Known  
Short Name: GravParKnown  
Database Table Name: Demographics  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient's mother's gravidity and parity are known.

---

SeqNo: 235  
Long Name: Mother's Gravidity  
Short Name: Gravidity  
Database Table Name: Demographics  
Data Source: User  
Format: Integer  
Definition: Indicate the number of times the mother of the patient has been pregnant, regardless of whether these pregnancies were carried to term. This includes the current pregnancy.

Low Value: 1 High Value: 30  
Parent Long Name: Mother's Gravidity And Parity Known  
Parent Short Name: GravParKnown  
Parent Value(s): = "Yes"

---

SeqNo: 236  
Long Name: Mother's Parity  
Short Name: Parity  
Database Table Name: Demographics

---

Data Source: User  
Format: Integer  
Definition: Indicate the number of >20-week births the patient's mother has had. Pregnancies with multiple babies (twins, triplets, etc.) count as 1 birth.

Low Value: 0 High Value: 30  
Parent Long Name: Mother's Gravidity And Parity Known  
Parent Short Name: GravParKnown  
Parent Value(s): = "Yes"

---

February 2016: The definition is not very specific - are we documenting the mothers parity including this child with heart disease or prior to this child? As an example, a woman who has been pregnant twice (G2), has a 5 year old child, and now has delivered her second child who now has had heart surgery. When entering that case do we consider her a P1 or a P2? **Same as the Gravity, P2 since the child was delivered.**

---

SeqNo: 237  
Long Name: APGAR Scores Known  
Short Name: ApgarKnown  
Database Table Name: Demographics  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient's APGAR scores are known.

---

SeqNo: 238  
Long Name: APGAR Score At 1 Minute  
Short Name: Apgar1  
Database Table Name: Demographics  
Data Source: User  
Format: Integer  
Definition: Indicate the patient's APGAR score at 1 minute after birth.

Low Value: 0 High Value: 10  
Parent Long Name: APGAR Scores Known  
Parent Short Name: ApgarKnown  
Parent Value(s): = "Yes"

---

SeqNo: 239  
Long Name: APGAR Score At 5 Minutes  
Short Name: Apgar5  
Database Table Name: Demographics  
Data Source: User  
Format: Integer

---

Definition: Indicate the patient's APGAR score at 5 minutes after birth.

Low Value: 0 High Value: 10  
Parent Long Name: APGAR Scores Known  
Parent Short Name: ApgarKnown  
Parent Value(s): = "Yes"

---

SeqNo: 240  
Long Name: Mother's Name Known  
Short Name: MatNameKnown  
Database Table Name: Demographics  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the name of patient's biological mother at time of patient's birth is known. If the patient is adopted and the name of the patient's biological mother is not known, indicate whether the name of the patient's adopted mother is known.

---

SeqNo: 250  
Long Name: Mother's Last Name  
Short Name: MatLName  
Database Table Name: Demographics  
Data Source: User  
Format: Text  
Definition: Indicate the last name of patient's biological mother at time of patient's birth, if it is known.  
  
If the patient is adopted, if the last name of the patient's biological mother is known, please enter the last initial of the patient's biological mother.  
  
If the patient is adopted, if the last name of the patient's biological mother is not known, please enter the last name of the patient's adopted mother.  
  
This field should be collected in compliance with state/local privacy laws.  
  
Parent Long Name: Mother's Name Known  
Parent Short Name: MatNameKnown  
Parent Value(s): = "Yes"

---

SeqNo: 260  
Long Name: Mother's First Name  
Short Name: MatFName  
Database Table Name: Demographics  
Data Source: User

---

Format: Text  
Definition: Indicate the first name of patient's biological mother at time of patient's birth, if it is known.

If the patient is adopted, if the first name of the patient's biological mother is known, please enter the first name of the patient's biological mother.

If the patient is adopted, if the first name of the patient's biological mother is not known, please enter the first name of the patient's adopted mother.

This field should be collected in compliance with state/local privacy laws.

Parent Long Name: Mother's Name Known  
Parent Short Name: MatNameKnown  
Parent Value(s): = "Yes"

---

SeqNo: 280  
Long Name: Mother's Middle Name  
Short Name: MatMName  
Database Table Name: Demographics  
Data Source: User  
Format: Text  
Definition: Indicate the middle name of patient's biological mother at time of patient's birth, if it is known.

If the patient is adopted, if the middle name of the patient's biological mother is known, please enter the middle name of the patient's biological mother.

If the patient is adopted, if the middle name of the patient's biological mother is not known, please enter the middle name of the patient's adopted mother.

This field should be collected in compliance with state/local privacy laws.

Parent Long Name: Mother's Name Known  
Parent Short Name: MatNameKnown  
Parent Value(s): = "Yes"

---

SeqNo: 290  
Long Name: Mother's National Identification (Social Security Number) Known  
Short Name: MatSSNKnown  
Database Table Name: Demographics  
Data Source: User  
Format: Text (categorical values specified by STS)

---

**Definition:** Indicate whether the Social Security Number (SSN) of patient's biological mother at time of patient's birth is known.

If the patient is adopted and the SSN of the patient's biological mother is not known, please indicate whether the SSN of the patient's adopted mother is known.

This field should be collected in compliance with state/local privacy laws.

---

SeqNo: 300

Long Name: Mother's National Identification (Social Security Number)

Short Name: MatSSN

Database Table Name: Demographics

Data Source: User

Format: Text

**Definition:** Indicate the Social Security Number (SSN) of patient's biological mother at time of patient's birth, if it is known. Although this is the SSN in the USA, other countries may have a different National Patient Identifier Number. For example in Canada, this would be the Social Insurance Number. If the patient is adopted, if the SSN of the patient's biological mother is known, please enter the SSN of the patient's biological mother.

If the patient is adopted, if the SSN of the patient's biological mother is not known, please enter the SSN of the patient's adopted mother.

This field should be collected in compliance with state/local privacy laws.

Parent Long Name: Mother's National Identification (Social Security Number) Known

Parent Short Name: MatSSNKnown

Parent Value(s): = "Yes"

---

SeqNo: 310

Long Name: Date of Birth

Short Name: DOB

Database Table Name: Demographics

Data Source: User

Format: Date - mm/dd/yyyy

**Definition:** Indicate the patient's date of birth using 4-digit format for year. This field should be collected in compliance with state/local privacy laws.

---

SeqNo: 320

Long Name: Birth Weight Known

Short Name: BirthWtKnown

Database Table Name: Demographics

Data Source: User

---

Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient's birth weight is known.

---

SeqNo: 330  
Long Name: Birth Weight  
Short Name: BirthWtKg  
Database Table Name: Demographics  
Data Source: User  
Format: Real, at least 3 decimal places  
Definition: Indicate the weight in kilograms of the patient at birth.

Low Value: 0.100 High Value: 10.000

Parent Long Name: Birth Weight Known  
Parent Short Name: BirthWtKnown  
Parent Value(s): = "Yes"

---

SeqNo: 340  
Long Name: Gender  
Short Name: Gender  
Database Table Name: Demographics  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the patient's gender at birth as male, female or ambiguous.

---

SeqNo: 350  
Long Name: Premature Birth  
Short Name: Premature  
Database Table Name: Demographics  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient was born prematurely as defined by a gestational period of less than 37 weeks.

---

SeqNo: 360  
Long Name: Gestational Age At Birth Known  
Short Name: GestAgeKnown  
Database Table Name: Demographics  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient's gestational age at birth is known.

---

---

SeqNo: 370  
Long Name: Gestational Age At Birth In Weeks  
Short Name: GestAgeWeeks  
Database Table Name: Demographics  
Data Source: User  
Format: Integer  
Definition: Indicate the patient's estimated gestational age at birth in weeks. This field is a required field for neonates and infants and is an optional field for children and adults.

Low Value: 16 High Value: 44

Parent Long Name: Gestational Age At Birth Known  
Parent Short Name: GestAgeKnown  
Parent Value(s): = "Yes"

---

SeqNo: 380  
Long Name: Multiple Gestation  
Short Name: MultGest  
Database Table Name: Demographics  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient was part of a multiple gestation, such as twins or triplets.

---

SeqNo: 381  
Long Name: Antenatal Diagnosis of Congenital Heart Disease  
Short Name: AntenatalDiag  
Database Table Name: Demographics  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a cardiac anomaly was diagnosed antenatally (e.g., fetal ultrasound).

---

SeqNo: 382  
Long Name: Fundamental Diagnosis  
Short Name: FundDiagnosis  
Database Table Name: Demographics  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: The fundamental diagnosis is a diagnosis that is carried with a patient throughout life through all operations and hospitalizations. The fundamental diagnosis is the most

complex cardiac anomaly or condition (congenital or acquired) of the patient.

Most frequently, the primary diagnosis will also be the fundamental diagnosis. For some operations, however, the fundamental diagnosis and primary diagnosis will be different.

For example, consider a child who underwent repair of subaortic stenosis, subsequently develops complete atrioventricular (AV) block, and undergoes pacemaker placement within the same hospitalization. The primary diagnosis for the pacemaker surgery is "Arrhythmia, Heart block, Acquired", while the fundamental diagnosis is "Aortic stenosis, Subvalvar".

Similarly, a patient who has a complete AV canal defect and undergoes either palliation or repair of the defect has a primary and fundamental diagnosis of "AVC (AVSD), Complete CAVSD". Subsequently, the child develops mitral insufficiency and is re-hospitalized for mitral valve replacement. The primary diagnosis for the mitral valve replacement operation is "Mitral regurgitation", but the fundamental diagnosis is "AVC (AVSD), Complete CAVSD."

The utilization of the fundamental diagnosis field, it is hoped, will clarify designation of a primary diagnosis, and enable greater specificity in the lesion specific report analyses.

March 2016 – What should I code as the fundamental diagnosis for a patient with familial ateriopathy with severe hypoplasia of pulmonary and aortic vascular tree, Proximal Left main coronary ostial stenosis, hypoplastic ascending, transverse and descending aorta, hypoplastic pulmonary arteries, stenotic pulmonary valve?

**Coronary other, hypoplastic arch.**

---

May 2016: A patient is admitted with a diagnosis of recurrent aortic stenosis, subvalvar. He has previously undergone a repair of subvalvar aortic stenosis along with a repair/resection of coarctation of the aorta. I am identifying aortic stenosis, subvalvar as the primary diagnosis for the current surgery, but what would the best choice be for the child's fundamental diagnosis? Would it be related to the coarctation of the aorta which has been repaired or the recurrence of the subvalvar aortic stenosis? **Either one would be appropriate.**

April 2017: What is the best choice for fundamental diagnosis for single ventricle patients with both heterotaxy and unbalanced AV Canal? Single Ventricle Heterotaxy? or Single Ventricle Unbalanced AV Canal? **Single Ventricle Heterotaxy, as this is the most complex of the patient's diagnoses.**

July 2017: We have a patient with a diagnosis of Partial AV canal, severe, discrete juxtaductal coarctation of the aorta and a PDA. She came to the OR for repair of the CoAo and PDA ligation. I'm thinking the fundamental diagnosis should be Coarctation of the Aorta(CoAo), with complex intracardiac anomaly. Is this correct?

**The fundamental diagnosis is the Partial AV canal (valve disease) as could potentially have more lifelong consequence but in this case, ask your surgeon which he/she thinks may cause more long term disease/sequelae.**

SeqNo: 383

Long Name: Race Documented

Short Name: RaceDocumented

Database Table Name: Demographics

---



Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether race is documented.

SeqNo: 390

Long Name: Race - Caucasian

Short Name: RaceCaucasian

Database Table Name: Demographics

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether the patient's race, as determined by the patient or family, includes Caucasian. This includes a person having origins in any of the original peoples of Europe, the Middle East, or North Africa.

Definition source: Standards for Maintaining, Collecting, and Presenting Federal Data on Race and Ethnicity: The minimum categories for data on race and ethnicity for Federal statistics, program administrative reporting, and civil rights compliance reporting. ([www.whitehouse.gov/omb/fedreg/1997standards.html](http://www.whitehouse.gov/omb/fedreg/1997standards.html))

Parent Long Name: Race Documented

Parent Short Name: RaceDocumented

Parent Value(s): = "Yes"

---

SeqNo: 400

Long Name: Race - Black / African American

Short Name: RaceBlack

Database Table Name: Demographics

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether the patient's race, as determined by the patient or family, includes Black / African American. This includes a person having origins in any of the black racial groups of Africa. Terms such as "Haitian" or "Negro" can be used in addition to "Black or African American."

Definition source: Standards for Maintaining, Collecting, and Presenting Federal Data on Race and Ethnicity: The minimum categories for data on race and ethnicity for Federal statistics, program administrative reporting, and civil rights compliance reporting. ([www.whitehouse.gov/omb/fedreg/1997standards.html](http://www.whitehouse.gov/omb/fedreg/1997standards.html))

Parent Long Name: Race Documented

Parent Short Name: RaceDocumented

Parent Value(s): = "Yes"

---

SeqNo: 410  
Long Name: Race - Asian  
Short Name: RaceAsian  
Database Table Name: Demographics  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient's race, as determined by the patient or family, includes Asian. This includes a person having origins in any of the original peoples of the Far East, Southeast Asia, or the Indian subcontinent including, for example, Cambodia, China, India, Japan, Korea, Malaysia, Pakistan, the Philippine Islands, Thailand, and Vietnam.

Definition source: Standards for Maintaining, Collecting, and Presenting Federal Data on Race and Ethnicity: The minimum categories for data on race and ethnicity for Federal statistics, program administrative reporting, and civil rights compliance reporting. ([www.whitehouse.gov/omb/fedreg/1997standards.html](http://www.whitehouse.gov/omb/fedreg/1997standards.html))

Parent Long Name: Race Documented  
Parent Short Name: RaceDocumented  
Parent Value(s): = "Yes"

---

SeqNo: 420  
Long Name: Race - American Indian / Alaskan Native  
Short Name: RaceNativeAm  
Database Table Name: Demographics  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient's race, as determined by the patient or family, includes American Indian / Alaskan Native. This includes a person having origins in any of the original peoples of North and South America (including Central America), and who maintains tribal affiliation or community attachment.

Definition source: Standards for Maintaining, Collecting, and Presenting Federal Data on Race and Ethnicity : The minimum categories for data on race and ethnicity for Federal statistics, program administrative reporting, and civil rights compliance reporting.

Parent Long Name: Race Documented  
Parent Short Name: RaceDocumented  
Parent Value(s): = "Yes"

---

SeqNo: 430  
Long Name: Race - Native Hawaiian / Pacific Islander  
Short Name: RaceNativePacific

---

Database Table Name:	Demographics
Data Source:	User
Format:	Text (categorical values specified by STS)
Definition:	Indicate whether the patient's race, as determined by the patient or family, includes Native Hawaiian / Pacific Islander. This includes a person having origins in any of the original peoples of Hawaii, Guam, Samoa, or other Pacific Islands.
	Definition source: Standards for Maintaining, Collecting, and Presenting Federal Data on Race and Ethnicity: The minimum categories for data on race and ethnicity for Federal statistics, program administrative reporting, and civil rights compliance reporting. ( <a href="http://www.whitehouse.gov/omb/fedreg/1997standards.html">www.whitehouse.gov/omb/fedreg/1997standards.html</a> )
Parent Long Name:	Race Documented
Parent Short Name:	RaceDocumented
Parent Value(s):	= "Yes"

---

SeqNo:	440
Long Name:	Race - Other
Short Name:	RaceOther
Database Table Name:	Demographics
Data Source:	User
Format:	Text (categorical values specified by STS)
Definition:	Indicate whether the patient's race, as determined by the patient or family, includes any other race.
Parent Long Name:	Race Documented
Parent Short Name:	RaceDocumented
Parent Value(s):	= "Yes"

---

SeqNo:	450
Long Name:	Hispanic Or Latino Ethnicity
Short Name:	Ethnicity
Database Table Name:	Demographics
Data Source:	User
Format:	Text (categorical values specified by STS)
Definition:	Indicate if the patient is of Hispanic or Latino ethnicity as determined by the patient / family. Hispanic or Latino ethnicity includes patient report of Cuban, Mexican, Puerto Rican, South or Central American, or other Spanish culture or origin, regardless of race.
SeqNo:	460
Long Name:	Date of Last Follow-Up
Short Name:	LFUDate

---

Database Table Name: Demographics  
 Data Source: User  
 Format: Date - mm/dd/yyyy  
 Definition: Indicate the date on which the last follow-up was made. If patient dies in the hospital, this value will be the same as the date of death. If no follow-up is made after patient is discharged, this value will be the same as the discharge date.

---

SeqNo: 470  
 Long Name: Last Follow-Up New York Heart Association Classification  
 Short Name: LFUNYHA  
 Database Table Name: Demographics  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate the patient's New York Heart Association (NYHA) classification at the time of the last follow-up. If no follow-up is made after patient is discharged, this value will be the same as the classification at the time of their last discharge.

---

July 2017: When filling out this question, what time reference are we using? Is this part of the 30 day assessment? Or is this information pertaining to the admission or surgical event date? In other words-is the last follow up the previous time they had a follow up from the date of admission? Or the follow up that occurred after discharge from this event? **It is the latest time a patient is seen whether from this event or the date of admission.**

---

SeqNo: 480  
 Long Name: Mortality Status At Last Follow-Up  
 Short Name: LFUMortStat  
 Database Table Name: Demographics  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate the mortality status of the patient at the time of the last follow-up. If no follow-up is made after patient is discharged, this value will be the same as the Mortality Status at Hospital Discharge.

---

SeqNo: 490  
 Long Name: Mortality Date  
 Short Name: MtDate  
 Database Table Name: Demographics  
 Data Source: User  
 Format: Date - mm/dd/yyyy  
 Definition: Indicate the patient's date of death.

Parent Long Name: Mortality Status At Last Follow-Up  
 Parent Short Name: LFUMortStat  
 Parent Value(s): = "Dead"

---

### 3. Noncardiac Congenital Anatomic Abnormalities

SeqNo: 510  
 Long Name: Noncardiac Congenital Anatomic Abnormalities Table Unique Record Identifier  
 Short Name: NCAAUniqueID  
 Database Table Name: NCAA  
 Data Source: Automatic  
 Format: Text  
 Definition: Unique identifier for the record in the Noncardiac Congenital Anatomic Abnormalities table.

SeqNo: 520  
 Long Name: Noncardiac Congenital Anatomic Abnormalities Link to Demographics Table  
 Short Name: PatID  
 Database Table Name: NCAA  
 Data Source: Automatic  
 Format: Text  
 Definition: An arbitrary, unique value generated by the software that permanently identifies each patient demographic record in the participant's database. This field is the foreign key that links this record with the associated record in the Demographics table.

SeqNo: 530  
 Long Name: Major Noncardiac Abnormality  
 Short Name: NCAA  
 Database Table Name: NCAA  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate all of the major noncardiac abnormalities the patient has or select None.

April 2016 – If a patient has a condition that was surgically corrected, for example, “imperforate anus”, should this still be recorded as an NCAA? **Yes**

July 2016: I have a patient who was adopted at age of 2 yrs old, and the above information is UNK. How should I code it? Can we add the option of UNK to the next version? **Chromosomal abnormalities can be identified at any time during an individual’s life – not everyone has genetic testing at birth. Same is true with respect to clinical diagnoses related to the other two categories. If not known to be present, then nothing gets coded. There is no indication for an “Unknown” option.**

SeqNo: 540  
 Long Name: Major Noncardiac Abnormality - Other - Specify

Short Name: NCAAOthSp  
Database Table Name: NCAA  
Data Source: User  
Format: Text Data Length: 100  
Definition: Indicate the other major noncardiac abnormality.

Parent Long Name: Major Noncardiac Abnormality  
Parent Short Name: NCAA  
Parent Value(s): = "Other"

---

#### 4. Chromosomal Abnormalities

SeqNo: 550  
Long Name: Chromosomal Abnormalities Table Unique Record Identifier  
Short Name: ChromAbUniqueID  
Database Table Name: ChromAbnormalities  
Data Source: Automatic  
Format: Text  
Definition: Unique identifier for the record in the Chromosomal Abnormalities table.

---

SeqNo: 560  
Long Name: Chromosomal Abnormalities Link to Demographics Table  
Short Name: PatID  
Database Table Name: ChromAbnormalities  
Data Source: Automatic  
Format: Text  
Definition: An arbitrary, unique value generated by the software that permanently identifies each patient demographic record in the participant's database. This field is the foreign key that links this record with the associated record in the Demographics table.

---

SeqNo: 570  
Long Name: Chromosomal Abnormality  
Short Name: ChromAb  
Database Table Name: ChromAbnormalities  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient has one of the following chromosomal abnormalities.

---

SeqNo: 580  
Long Name: Chromosomal Abnormality - Other - Specify

---

Short Name: ChromAbOthSp  
Database Table Name: ChromAbnormalities  
Data Source: User  
Format: Text Data Length: 100  
Definition: Indicate the other chromosomal abnormalities.

Parent Long Name: Chromosomal Abnormality  
Parent Short Name: ChromAb  
Parent Value(s): = "Other chromosomal abnormality"

---

## 5. Syndromes

SeqNo: 590  
Long Name: Syndromes Table Unique Record Identifier  
Short Name: SynUniqueID  
Database Table Name: Syndromes  
Data Source: Automatic  
Format: Text  
Definition: Unique identifier for the record in the Syndromes table.

---

SeqNo: 600  
Long Name: Syndromes Link to Demographics Table  
Short Name: PatID  
Database Table Name: Syndromes  
Data Source: Automatic  
Format: Text  
Definition: An arbitrary, unique value generated by the software that permanently identifies each patient demographic record in the participant's database. This field is the foreign key that links this record with the associated record in the Demographics table.

---

SeqNo: 610  
Long Name: Syndrome  
Short Name: Syndrome  
Database Table Name: Syndromes  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient has a "Syndrome" or "Syndromic abnormality". A "syndrome" is defined as a group of signs and symptoms that occur together, and characterize a particular abnormality [1].

[1]. Tchervenkov CI, Jacobs JP, Weinberg PM, Aiello VD, Beland MJ, Colan SD, Elliott

MJ, Franklin RC, Gaynor JW, Krogmann ON, Kurosawa H, Maruszewski B, Stellin G. The nomenclature, definition and classification of hypoplastic left heart syndrome.

Cardiology in the Young, 2006; 16(4): 339–368, August 2006.

---

June 2016: Does a syndrome like Tourette's belong in the syndrome field, the preoperative factor field (as a neurological deficit), or in both? **Tourette's would be in syndrome, other. It will not impact the patient's recovery.**

January 2017: Is it correct to interpret #160-Fetal Drug Exposure as anything taken that would not normally be recommended or prescribed for use during pregnancy? Or is this strictly illicit drugs? What is the purpose of this field? **Anything taken that would not normally be recommended or prescribed for use during pregnancy should be captured. The field pertains to drugs etc., which may not be illicit but some may be contraindicated during pregnancy.**

SeqNo: 620

Long Name: Syndrome - Other - Specify

Short Name: SyndromeOthSp

Database Table Name: Syndromes

Data Source: User

Format: Text Data Length: 100

Definition: Indicate the other "Syndrome" or "Syndromic abnormality".

Parent Long Name: Syndrome

Parent Short Name: Syndrome

Parent Value(s): = "Other syndromic abnormality"

---

## 6. Hospitalization

SeqNo: 630

Long Name: Hospital Name

Short Name: HospName

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by user) Data Length: 100

Definition: Indicate the full name of the facility where the procedure was performed. Values should be full, official hospital names with no abbreviations or variations in spelling for a single hospital. Values should also be in mixed-case.

---

SeqNo: 640

Long Name: Hospital Zip Code

Short Name: HospZIP

Database Table Name: Operations

---



Data Source: Lookup  
Format: Text Data Length: 20  
Definition: Indicate the ZIP Code of the hospital. Outside the USA, these data may be known by other names such as Postal Code. This field should be collected in compliance with state/local privacy laws.

---

SeqNo: 650  
Long Name: Hospital State  
Short Name: HospStat  
Database Table Name: Operations  
Data Source: Lookup  
Format: Text Data Length: 50  
Definition: Indicate the region of the country (i.e., state or province) in which the hospital is located.

---

SeqNo: 660  
Long Name: Hospital National Provider Identifier  
Short Name: HospNPI  
Database Table Name: Operations  
Data Source: Lookup  
Format: Text  
Definition: Indicate the hospital's National Provider Identifier (NPI). This number, assigned by the Center for Medicare and Medicaid Services (CMS), is used to uniquely identify facilities for Medicare billing purposes.

---

SeqNo: 771  
Long Name: Primary Payor  
Short Name: PayorPrim  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the primary insurance payor for this admission.

---

SeqNo: 772  
Long Name: Primary Payor Medicare Fee For Service  
Short Name: PrimMCCareFFS  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient is covered by Medicare Fee For Service (Part B).

---

Parent Long Name: Primary Payor  
Parent Short Name: PayorPrim  
Parent Value(s): = "Medicare"

---

SeqNo: 773  
Long Name: Secondary (Supplemental) Payor  
Short Name: PayorSecond  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate which if any secondary insurance payor was used for this admission.

Parent Long Name: Primary Payor  
Parent Short Name: PayorPrim  
Parent Value(s): Is Not "None / self" And Is Not Missing

---

SeqNo: 774  
Long Name: Secondary Payor Medicare Fee For Service  
Short Name: SecondMCareFFS  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient is covered by Medicare Fee For Service (Part B).

Parent Long Name: Secondary (Supplemental) Payor  
Parent Short Name: PayorSecond  
Parent Value(s): = "Medicare"

---

SeqNo: 780  
Long Name: Date of Admission  
Short Name: AdmitDt  
Database Table Name: Operations  
Data Source: User  
Format: Date - mm/dd/yyyy  
Definition: Indicate the date the patient was admitted to the hospital. For those patients who originally enter the hospital in an out-patient capacity (i.e., catheterization), but then are not discharged, the admit date is the date of the patients entry into the hospital.

---

May 2016: What is meant by 'hospitalization'? Does this include periods of time when a patient was cared for at an OSH prior to the admission to our facility for surgery? **It is the date of admission to your facility.**

---

SeqNo: 781  
 Long Name: Location From Which Patient Admitted  
 Short Name: AdmitFromLoc  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate the location from which the patient was admitted.

---

SeqNo: 790  
 Long Name: Date of Surgery  
 Short Name: SurgDt  
 Database Table Name: Operations  
 Data Source: User  
 Format: Date - mm/dd/yyyy  
 Definition: Indicate the date of surgery which equals the date the patient enters the OR or equivalent.

---

August 2016: We had a patient who needed an angiogram during his operation. We don't have a hybrid suite so he left the OR on ECMO to go to the cath lab, then came back to the OR and placed back on bypass to finish his operation. Should I code this as one surgery or two? **This would be considered one surgery. The reason is if your facility did have a hybrid suite this situation would be considered one procedure.**

SeqNo: 800  
 Long Name: Height in Centimeters  
 Short Name: HeightCm  
 Database Table Name: Operations  
 Data Source: User  
 Format: Real  
 Definition: Indicate the height of the patient in centimeters at the time of surgery.

Low Value: 15.0    High Value: 250.0

---

SeqNo: 810  
 Long Name: Weight in Kilograms  
 Short Name: WeightKg  
 Database Table Name: Operations  
 Data Source: User  
 Format: Real, at least 3 decimal places  
 Definition: Indicate the weight of the patient in kilograms at the time of surgery.

Low Value: 0.001    High Value: 200.000

---

April 2017: What is the purpose of this field? Is it most accurate to capture the patient's dry (dosing) weight or the patient's actual (daily) weight? **Interpret the specs literally. Use the weight that had been most recently recorded at the time of, or just preceding surgery. This number would typically be used by anesthesiologist to calculate drug dosages, and by the Perfusionist to determine pump flow rates, etc. The weight among infants and neonates is a covariate in the Mortality Risk Model that is used to calculate O:E ratios and report risk-adjusted mortality rates (AMR).**

SeqNo: 820

Long Name: Patient Age In Days

Short Name: AgeDays

Database Table Name: Operations

Data Source: User or Automatic

Format: Integer

Definition: Calculate the patient's age in days at the time of the surgery procedure. The patient's age will be calculated by the software from the date of birth and the date of surgery.

Low Value: 0      High Value: 40150

## 7. Preoperative Factors

SeqNo: 830

Long Name: Preoperative Factor Table Unique Record Identifier

Short Name: PoFUniqueID

Database Table Name: PreopFactors

Data Source: Automatic

Format: Text

Definition: Unique identifier for the record in the Preoperative Factors table.

SeqNo: 840

Long Name: Preoperative Factor Link to Operations Table

Short Name: OperationID

Database Table Name: PreopFactors

Data Source: Automatic

Format: Text

Definition: An arbitrary, unique value generated by the software that permanently identifies each operation record in the participant's database. This field is the foreign key that links the Preoperative Factor record with the associated record in the Operations table.

SeqNo: 850

Long Name:	Preoperative Factor
Short Name:	PreopFactor
Database Table Name:	PreopFactors
Data Source:	User
Format:	Text (categorical values specified by STS)
Definition:	Indicate the factors that are present preoperatively that may impact the patient's outcome.

May 2016: Specifically in regards to PreopFactor 470 - should we code 'mechanical ventilation to treat cardiorespiratory failure' if a patient is intubated for this reason at an OSH prior to coming to our hospital for surgery? **If the patient was intubated to treat respiratory failure then you should indicate 'yes' even if the patient was intubated at an OSH.**

June 2016: I am not always sure what to include in each of the categories: For example: Induced exercise asthma should be included in asthma? **Yes, if they are receiving treatment.** ADD/ADHD should be included in neurological deficit? **No** For renal dysfunction, do you check the lab results or if it is stated in the H&P or in another note that the patient has renal dysfunction it is enough? **There should be documentation supporting renal dysfunction in the medical record.** A 9 yo with trisomy 18 who is non-verbal, non ambulatory and has cognitive function of 3 months old, would you code him as neuro deficit or is it under trisomy 18 in chromosomal abnormalities? **This is not a neuro deficit.** Is renal dysfunction for life or in the last year? **It is not for life or a specific time period. Look at how the patient is prior to surgery.**

July 2016: If a patient has a tracheostomy and is vented at time of OR entry, do you code both "Tracheostomy present" and "Mechanical ventilation to treat cardiorespiratory failure" as preop factors? **Yes**

August 2016: patient had a gastrostomy tube which was removed previously, however, patient has a fistula at the Gtube site. Would I count Gastrostomy present? The fistula is constantly draining onto the abdomen. **A fistula from the stomach to the outside is a "gastrostomy" so it would be appropriate to code "gastrostomy present."**

October 2016: Would 'failure to thrive' be documented under 'other preoperative factors'? And would if it is documented, is it documented even if the reason the patient is failure to thrive because the heart isn't allowing them to grow? **Failure to thrive is not captured in the CHSD; it is your choice whether or not to capture it.**

January 2017: Should inhaled steroids be included in the "currently taking steroids" fields for preoperative factors? **Inhaled steroids should not be included as they are generally taken for reactive airway disease and are not the equivalent of systemic steroid ingestion. The intent of the field was probably related to factors such as: 1) potential increased infection risk, 2) potential impact on healing, and 3) potential for adrenal suppression (and need for "stress" steroid coverage).**

April 2017: For mechanical vent in the pre-operative period, does guidance remain to capture this variable when the patient has experienced mechanical ventilation at any time in the admission, prior to procedure? For example, a patient undergoes CPB in index procedure and is extubated post-op. He then returns to OR for a minor procedure, where he is re-intubated for surgery. Since he was not intubated at the time of the second OR Entry Date/Time, do we still capture Mechanical Vent in Pre-op risk? **No, only include this risk factor if the patient had previously required mechanical ventilation for respiratory failure.**

April 2017: Pt was admitted in Jan 2017 with a possible mass on mitral valve. R/o endocarditis. No fever, neg cultures not on long term antibiotics discharged on Coumadin for thrombus. Pt returned on 3/23 with TIA symptoms and TEE showed a thrombus or vegetation on mitral valve. Again no fever no elevated labs. To OR to remove mitral ring and found in OR slime like material around mitral ring and this time positive for lactobacillus. So here is my question. On risk factors do i mark Endocarditis yes, active and culture other since it was found

in this admission or does it get marked no since it was not known prior to this admission? **Yes, code as endocarditis as the patient was diagnosed in the OR and had histologic evidence of endocarditis.**

May 2017: If a mother uses tobacco products during pregnancy would this count as a preop factor of '570 - Tobacco use'? Would it count as '166 - Fetal drug exposure' under syndromes? **No, maternal smoking is currently not captured currently in the database.**

July 2017: If a patient was treated for NEC at another institution and then transferred to ours for transplant evaluation and later transplant, would NEC be coded as a preoperative factor? **Technically, no, do not include as a preoperative factor.**

August 2017: Should Autism be included as a preoperative factor for neurological deficit? **No, based on historic answers of what constitutes a neurologic deficit (cognitive vs. motor).**

September 2017: An ENT patient sustains an intraoperative aortic injury during a jejunal interposition procedure, cardiac surgery is called in for repair. Should the preoperative factors be coded at the time this case "turns" cardiac, or from when the patient enters the OR for the ENT procedure? **Code the risk factors from when the case turns cardiac (cardiac surgeon enters the OR)**

## 8. Diagnosis

SeqNo: 870

Long Name:	Diagnosis Table Unique Record Identifier
Short Name:	DiagUniqueID
Database Table Name:	Diagnosis
Data Source:	Automatic
Format:	Text
Definition:	Unique identifier for the record in the Diagnosis table.

SeqNo: 880

Long Name:	Diagnosis Link to Operations Table
Short Name:	OperationID
Database Table Name:	Diagnosis
Data Source:	Automatic
Format:	Text
Definition:	An arbitrary, unique value generated by the software that permanently identifies each operation record in the participant's database. This field is the foreign key that links the Diagnosis record with the associated record in the Operations table.

SeqNo: 890

Long Name:	Diagnoses
Short Name:	Diagnosis
Database Table Name:	Diagnosis
Data Source:	User
Format:	Text (categorical values specified by STS)
Definition:	Indicate all diagnoses noted at the time of the surgical procedure or documented by

preoperative studies. This entry may duplicate the Fundamental Diagnosis.

February 2016: which diagnosis would most accurately reflect a diagnosis of inducible ventricular tachycardia found during an EP study (this would not be the patient's primary diagnosis) **"Arrhythmia"**

June 2016: An 8.2kg 7 month old patient underwent LVAD implantation at our institution. She subsequently recovered and presented on 5/23/16 for LVAD explantation. It was discovered during this procedure that there was a diaphragmatic hernia secondary to dense adhesions of the VAD apical cannula to the surface of the diaphragm. What would be the best way to capture this hernia in the diagnosis section? **Complications of VAD implantation; Diaphragmatic, other.**

August 2016: status post other procedures: are status post procedures just for cardiac? Patient is status post a Roux en Y jejunostomy. Would I code that? **It is not necessary to code for that type of gastrointestinal procedure, the s/p codes are mainly for cardiac procedures. Any of the "S/P" codes in the Data Collection form may be used, so you could conceivably code this as 6040=Status post - Miscellaneous procedure, Other or as 11777=Status post - Other procedure.**

October 2016: I am always puzzled how to code the "Open sternum with open (or closed) skin" diagnosis. When it is a past occurrence does it get carried over to the diagnosis list for a new visit? Or does it get dropped and only the SP Delayed sternal closure is coded? **Once the sternum is closed it is no longer a viable diagnosis. Becomes S/P Delayed Sternal Closure.**

November 2016: A patient with history of previous TOF repair was admitted to our facility for pulmonary valve replacement. He was also found to have a "floating"/aberrant left subclavian artery that was causing ischemic pain to his left arm. How would the diagnosis of aberrant Left SC artery be captured? **This cannot be captured on the short list. You can use "cardiac, other"**

December 2016: There is a procedure for valvuloplasty or valve replacement of a common AV Valve, but I can't find a diagnosis for Common AV valve insufficiency. Is there some other diagnosis I should be using? **Is this Left, Right, or Common? Use whichever side is worse as the diagnosis.**

August 2017: A patient with a history of PV stenosis (420) underwent a balloon valvuloplasty for the stenosis. When identifying diagnoses for this patient going forward, would the patient still carry the diagnosis of PV stenosis in addition to the diagnosis of s/p cardiac cath procedure, therapeutic, balloon dilation (6570) or would I only identify the patient as s/p card cath procedure, therapeutic, balloon dilation? **In this example, the fundamental diagnosis would be PV stenosis and would be carried as a definition always. Include both diagnoses.**

September 2017: Should I code an RVOT pseudoaneurysm as "aneurysm, other" or "aneurysm, ventricular, right (including pseudoaneurysm)? Would the related procedure then be "RVOT procedure" or "aneurysm, ventricular, right, repair"? **Code aneurysm, ventricular, right (including pseudoaneurysm) with aneurysm, ventricular, right, repair**

October 2017: I am looking for the best diagnosis that captures this anomaly: DILV, DORV(S,L,L) Upstairs/Downstairs heart with dominant inferior LV and hypoplastic superior RV Criss cross AV valves with straddling TV. **Likely Single ventricle, DILV for primary diagnosis. Refer to surgeons.**

SeqNo: 900

Long Name: Primary Diagnosis Indicator

Short Name: PrimDiag

Database Table Name: Diagnosis

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate the diagnosis of primary importance at the time of this surgical procedure.

Example: fundamental diagnosis of Tetralogy of Fallot. The current Diagnoses are both pulmonary insufficiency and residual ventricular septal defect. In this case, pulmonary insufficiency will be flagged as the primary diagnosis.

---

February 2016 – Where would I capture inducible ventricular tachycardia as a pre-operative diagnosis?

**Capture as pre-operative arrhythmia, ventricular; code as secondary diagnosis, arrhythmia.**

August 2017: I am having trouble with selecting the correct primary diagnosis for a patient who is being cannulated for ECMO. The patient has a fundamental diagnosis of AVC (transitional), but has not undergone surgery for it. She is being cannulated for severe resp failure/hypoxemia and cardiogenic shock. She has right heart failure mostly due to pulmonary hypertensive crises. Her pulmonary hypertension is not primary pulmonary hypertension. What would I select as a primary diagnosis for the ECMO cannulation? Also for this same patient who then recovers sufficiently to be de-cannulated, what would I select as a primary diagnosis for the de-cannulation? **Select the rationale for why the patient is requiring ECMO (shock, cardiac failure – they short list to cardiac, other).**

---

## 9. Procedures

SeqNo: 910

Long Name: Procedures Table Unique Record Identifier

Short Name: ProcUniqueID

Database Table Name: Procedures

Data Source: Automatic

Format: Text

Definition: Unique identifier for the record in the Procedures table.

---

SeqNo: 920

Long Name: Procedures Link to Operations Table

Short Name: OperationID

Database Table Name: Procedures

Data Source: Automatic

Format: Text

Definition: An arbitrary, unique value generated by the software that permanently identifies each operation record in the participant's database. This field is the foreign key that links the Procedure record with the associated record in the Operations table.

---

May 2016: There are currently six short-list options for Coarctation Repair in the STS Congenital Database.

They are "End to End", "End to Extended End", "Subclavian flap", "Patch Aortoplasty", "Interposition graft", and "Other". My surgeon dictates and describes his coarctation repairs as "Coarctation Repair, End to Side". How would you code this using current Coarctation Repair options. Our surgeon can provide more detail of the operation if needed. **Until the codes are updated to include this, use 'end to end – extended'**

---

SeqNo: 930

---



Long Name:	Procedures
Short Name:	Procedure
Database Table Name:	Procedures
Data Source:	User
Format:	Text (categorical values specified by STS)
Definition:	Indicate ALL procedures that were performed during this surgical procedure.

---

12/2015: What surgical procedure do I choose for re-implantation of anomalous right subclavian into right carotid? **Other cardiac unless in the setting of vascular ring repair. Usually an add on procedure. Not analyzed.**

February 2016 : What procedure would I select for a patient who has a reduction of the right atrium / removal RAA done? **Cardiac Procedure, other.**

February 2016: A patient with a previous TOF repair was admitted for a pulmonic valve replacement. A bioprosthetic pulmonary valve was placed and then the RVOT was reconstructed with a gore-tex patch for the roof. A cryoablation was performed along the anterior portion of the VSD patch as inducible VT had been mapped to this area. How would I capture this procedure, particularly the re-construction of the RVOT and the cryoablation? **Code separate RV procedure.**

February 2016: Really more of a clarification - we use the IPCCC Long list, for the procedure, PA Reconstruction (plasty), Reduction pulmonary artery arterioplasty - it maps to the STS term of Cardiac procedure, other. Is this working as designed? It seems as all other PA reconstruction procedures are mapping to the STS Term of PA, reconstruction (plasty). And, cardiac procedure, other does not have a STAT score. **Put the PA Plasty in the first place to get the STAT score.**

May 2016: A patient was admitted with a diagnosis of aortic stenosis, subvalvar. As part of the repair, the patient has a myectomy performed but the patient does not carry a diagnosis of IHSS. The only option for identifying a myectomy done as part of a subvalvar repair for aortic stenosis is if the myectomy was done for IHSS. How would I best capture this patient's procedure given he does not have IHSS? **Use 2100; it does not have to be for IHSS only.**

June 2016: Patient goes to OR for temporary pacemaker done by the surgeons then has a Hybrid later in the same admission. Temporary pacemaker is NO CPB Cardiovascular with a STAT score. Am I correct that the pacemaker will be the index operation? Is there anything that could be done differently so that the Hybrid is the index? **Yes, the pacemaker is the index procedure.**

August 2016: How would I best capture a Bentall procedure with re-implantation of coronary arteries? This was a re-do procedure. The child had previously had an aortic root replacement approx 13 years ago. **Coronary re-implantation is an element of a Bentall Procedure (so an additional code for this is not required). If it is truly a Bentall (which includes composite aortic root and aortic valve replacement), would code as "720= Aortic root replacement, Mechanical" or "715= Aortic root replacement, Bioprosthetic" depending on the type of prosthesis.**

August 2016: How would I capture a procedure where only a biopsy of mediastinal lymph nodes and a mediastinal mass was done? The surgeon documented "Chamberlain procedure for biopsy of mediastinal mass." **Use 2020=Thoracic and/or mediastinal procedure, other. You could also use 1860= Mediastinal procedure, or 1970= Mediastinal exploration. A Chamberlain Procedure is a "mediastinotomy" that is generally done through a small transverse incision next to the upper sternum, with or without resection of a piece of costal cartilage. Nowadays, a fiber-optic scope may be inserted, though traditionally the mediastinum was explored with surgical instruments, digitally, and with a rigid metal scope designed for this purpose. In most instances, some lymph nodes are sampled.**

August 2016: A patient underwent replacement of his Aortic valve with a bioprosthesis. He also required patch enlargement of his aortic root in order to accommodate an appropriate sized valve. How would this procedure best be captured on the DCF? **It depends on the type of annular enlargement: If not done as a Konno procedure, code: 770= Other annular enlargement procedure. This would encompass Nick's, Manougian, and other types of annular enlargement.**

October 2016: A few cases where the Glenn takes over.

I have 3 Ebstein's patients who had Ebstein's repair (stat 1.6) and Glenn (stat 0.5) **The Glenn takes over because of the procedure specific factors.**

Another case of congenitally corrected TGA repair, arterial switch and ASO (double switch) (stat 3.4) with Glenn. **This is correct.**

MAPCA unifocalization (stat 1.6) Rastelli (stat 0.9) and AVC (stat 0.8). **The AVC takes over because of the PSF. MVR (stat) and Glenn (0.4) The Glenn is primary**

A TOF patient, S/P TOF repair current had a tricuspid valvuloplasty, aortic valvuloplasty and VSD. The surgeon wants the tricuspid valvuloplasty to be the primary procedure. **The aortic valvuloplasty is the primary procedure.**

Ebstein's repair patient who had a tricuspid valvuloplasty (stat 0.7) and Glenn (0.5). We would like the tricuspid valvuloplasty to be the primary procedure. **The Ebstein's repair is the primary procedure.**

November 2016: How would I capture a bypass done from the left carotid artery to the left subclavian artery? This was a secondary procedure done during a concomitant pulmonary valve replacement. **You can use 2020 or 2040**

January 2017: I need a procedural code for a right jugular vein to right carotid artery ringed Gore-tex graft shunt that my surgeon performed on a failed fontan with a BDG with ongoing cyanosis. **Use Shunt to Other.**

February 2017: patient had PDA device inserted. The next day it was discovered that the device had embolized into right pulmonary artery. Patient then underwent foreign body retrieval. A week later the patient returned to hospital for surgical closure of PDA. How do I document the status post procedures? Are they 6610 device implantation and 7120 for foreign body removal, or instead of 6610, should it be 7110 for device implantation attempted. **You should document 6610 device implantation and 7120 for foreign body removal.**

February 2017: When a Ross procedure (Stat category 2) is an RV-PA conduit (Stat category 3) assumed as part of that or should it be coded separately? **It is part of the Ross. The Ross is the Primary Procedure. You can include the RV-PA in the valve section.**

April 2017: **PA Banding vs. Hybrid bilateral bands-** If a patient goes to the OR and undergoes a PA banding, regardless of the patient's diagnosis, I select procedure 1640. However, if the surgical plan includes a staged surgical or combination approach during the same hospitalization, I select 2160. For example, if the notes read the patient is to first undergo PA banding (bilateral banding) and then in the next 24-48 hours will return to the hybrid suite for ductal stenting, I select the hybrid procedure.

**Definitions from Specs: 1640 PA banding (PAB)** Placement of a pulmonary artery band, any type. **2160 Hybrid Approach "Stage 1", Application of RPA & LPA bands** A "Hybrid Procedure" is defined as a procedure that combines surgical and transcatheter interventional approaches. The term "Hybrid approach" is used somewhat differently than the term "Hybrid Procedure". A "Hybrid approach" is defined as any of a group of procedures that fit into the general silo of procedures developed from the combined use of surgical and transcatheter interventional techniques. Therefore, not all procedures classified as "Hybrid approach" are truly "Hybrid Procedures".

April 2017: How do you code a "Sano" shunt when it is the primary procedure with no other associated procedures (ie. a TOF with pulmonary atresia that receives a systemic to pulmonary shunt in the form of a "Sano")? If I code the systemic to pulmonary shunt, then it will not be included in my analysis because there is not an associated STAT score. If I choose to code it as an RV to PA conduit, it is a lower STAT score then it

should be and it does not represent the intent of the procedure. **Code as RV to PA conduit for now and will need to be addressed with the upgrade.**

June 2017: When we do an arch reconstruction with a BDG we get to list the arch as the primary procedure. When we do an arch reconstruction with a Fontan, we have to list Fontan as primary as it is not listed in the exceptions list. Can we get Fontan added to the exceptions list? **Per the current primary procedure rules, the fontan would be the primary procedure. It will be suggested to be changed in the next upgrade.**

August 2017: What is the difference between a Kawashima Procedure and a Bilateral Bidirectional Glenn? My surgeon wrote both throughout the chart. The diagnosis listed is complex congenital heart disease consisting of the following: heterotaxy syndrome with mirror image dextrocardia, unbalanced right ventricle dominant AV canal, double outlet right ventricle, malposed great vessels, mild pulmonary stenosis, right aortic arch, interrupted inferior vena cava with azygous continuation, bilateral superior vena cavae, and unobstructed pulmonary venous return to a confluence behind the right sided atrium. On his op note, the procedures listed are bilateral bidirectional Glenn and subtotal PA banding. What would be my primary diagnosis and primary procedure? **Kawashima is a type of cavopulmonary anastomosis used when a patient has an interrupted IVC. If a Kawashima was done, enter that procedure. The primary diagnosis is Single ventricle, Heterotaxy.**

September 2017: Surgeon is requesting suggestion for coding "RPA translocation (12 mm Gore-Tex graft)". Background information: Patient has Truncus-IAA, repaired in 2010, had previous stenting of RPA 2011, 2015, 2017. May 2017 - Had RPA arterioplasty using 8 mm split ringed Gore-Tex graft. Had persistent left lung collapse, therefore had ascending aorta repair May 2017. End of May still had problem with ventilating the left lung, therefore performed Translocation RPA anterior to the ascending aorta to give the bronchus more space between the ascending and descending aorta posteriorly. **There is no current way to code this procedure as a stand-alone procedure. Code as Cardiac, Other.**

September 2017: For the following list of operation, which procedure codes should be used: PREOPERATIVE DIAGNOSES: 1. Distal ascending and proximal arch aortic aneurysm. 2. Marfan syndrome. 3. Status post Bentall type procedure with 27 millimeter Carbomedics mechanical valve composite graft. 4. Status post aortic valve-sparing root replacement with 18 millimeter Hemashield graft, mitral valve repair with 25 millimeter Duran annuloplasty ring, and tricuspid valve repair. POSTOPERATIVE DIAGNOSES: Same. OPERATIONS PERFORMED: 1. Redo sternotomy. 2. Total arch replacement with insertion of 28 millimeter multi-branched Vascutek graft under hypothermic circulatory arrest. 3. Aorta to innominate artery bypass with 8 millimeter Vascutek graft. 4. Aorta to left common carotid artery bypass with 8 millimeter Vascutek graft. **Primary procedure is the aortic arch repair.**

October 2017: When we have coded our Norwood patients, we have always coded the manner of arch reconstruction first and source of pulmonary blood flow second. We have used the Norwood modifier for source of pulmonary blood flow as opposed to palliation, shunt....etc... We are now being told that we cannot use the modifier in the second field as source of pulmonary blood flow and to go back years and correct the data. Is this true? If so, why is the modifier field in place for the Norwood patients? Finally, when one queries the database for number of shunts we get both shunts and index operations and shunts as parts of larger operations, which are complete entities. **This was a change multiple harvests ago (related to mapping of the procedures within the database) asking centers to use only the accepted sources of pulmonary blood flow. Refer to the Report Overview (pg. 19) of the most recent data harvest to see those accepted sources.**

October 2017: We have a very sick child whom has undergone multiple sternal washouts and the surgeon is the one performing all of the wound vac management. The surgeon is performing washouts and wound vac management every three days. Does each washout/ wound vac change get counted as a new case?

**Capture all cases the surgeons perform.**

November 2017: Pt born with multiple VSDs and an ASD. S/P PA Banding, Ao Arch Repair and PDA Closure, surgical as newborn. Our surgeon has coded the primary dx for this event ASD, Secundum and the primary

procedure as PA Band adjustment and also lists ASD Repair, primary closure. Do we need to change the primary procedure to ASD Repair? **No, primary procedure is PA band adjustment based on the stat scores of the procedures.**

November 2017: Dx: Marfan syndrome and enlarged aortic root/thoracic aortic aneurysm and 2+ aortic valve regurgitation. Procedures include: The valve was then re-implanted within the tube graft and the coronary artery buttons attached to the side with Teflon doughnuts. Would you code 1380-Ao Aneurysm Repair or 735-Ao root replacement, valve sparing, or both? **Code both, with the primary the 735, Aortic root replacement.**

November 2017: How is a revised shunt coded? **Code the same as the initial procedure.**

SeqNo: 940

Long Name: Primary Procedure Indicator

Short Name: PrimProc

Database Table Name: Procedures

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether this procedure is considered the PRIMARY Procedure performed during this operation.

Note that the primary procedure is determined at the data warehouse using the methodology published in the Journal of Thoracic and Cardiovascular Surgery ("An empirically based tool for analyzing mortality associated with congenital heart surgery" Sean M. O'Brien, David R. Clarke, Jeffrey P. Jacobs, Marshall L. Jacobs, Francois G. Lacour-Gayet, Christian Pizarro, Karl F. Welke, Bohdan Maruszewski, Zdzislaw Tobota, Weldon J. Miller, Leslie Hamilton, Eric D. Peterson, Constantine Mavroudis and Fred H. Edwards J Thorac Cardiovasc Surg 2009;138:1139-1153 DOI:10.1016/j.jtcvs.2009.03.071). If the above methodology does not return a primary procedure, this field will be used to designate primary procedure.

12/2015: What is the primary procedure when a patient undergoes the following repairs during the same operative setting: tricuspid valvuloplasty, pulmonary valvuloplasty, patch VSD repair, primary closure ASD PFO? The tricuspid valve repair has the highest STAT score but when completed in the setting of the VSD repair, the VSD would be designated as primary. How is this rule impacted with the pulmonary valve repair? **Code as VSD**

February 2016: I don't see the Kawashima operation on our PSF list but it is technically superior cavo with unilateral bidirectional... shouldn't it have one? No. **Patients should not be coded as Glens**

April 2016: What is the primary procedure in the following scenario: The surgeon initially places a central shunt and separates from bypass. Subsequently, the patient declines and is replaced on bypass, the central shunt is taken down and a modified BT shunt is instead placed. I want to be accurate in my coding and list all procedures that were completed, however the central shunt would be considered the primary procedure and the patient never left the operating room with one. **Code BT Shunt as primary**

May 2016: Operative diagnosis: Sinus venosus ASD, PAPVR, secundum ASD. Procedure: Patch closure of secundum ASD, atrial septectomy, repair of PAPVR using atrial baffle. Sinus venosus ASD left alone due to small size. How should this operation be coded? **Use 260; PAPVC repair; only code the PAPVR. Do not code the ASD patch closure.**

December 2016: I have two patients that underwent a bilateral bidirectional Glenn procedure AND a TAPVR repair within the same operation. What is the appropriate way to code this? We do not want to lose out on a STAT 4 category procedure, but my understanding is that due to procedure specific factors, the Glenn will be the primary procedure however the TAPVR is a higher STAT score. This is similar to when a DKS is performed at the same time as a Glenn and it was recently discussed in the AQO meeting that an exception will be made for this combination when analyzed by DCRI. Is there any chance that a TAPVR repair can be made as an exception as well? **The Glenn is the Primary Procedure.**

---

## 10. Procedure-Specific Factors

SeqNo: 949

Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure

Short Name: PSFPrimProc

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate which, if any, of the following "benchmark operations" was the primary procedure for this operation.

---

SeqNo: 950

Long Name: Procedure-Specific Factors - Apical VSD

Short Name: PSFApicalVSD

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether Apical VSD was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure

Parent Short Name: PSFPrimProc

Parent Value(s): = "VSD repair, Primary closure", "VSD repair, Patch", "VSD repair, Device", "Arterial switch operation (ASO) and VSD repair" or "Arterial switch procedure and VSD repair + Aortic arch repair"

---

October 2016: Can you please tell me the difference between Apical VSD and Straddling AV valve. And, do we need to document on both Sequence #950 and #951 whether the VSD repair is primary closure, patch or device? **The Cardiologist or ECHO report should document whether it is an Apical VSD or a Straddling AV valve. Yes, both seq. 950 and 951 should be documented 'Yes' or 'No'.**

---

SeqNo: 951

Long Name: Procedure-Specific Factors - Straddling AV valve

Short Name: PSFStradAVVal

Database Table Name: Operations

---

Data Source:	User
Format:	Text (categorical values specified by STS)
Definition:	Indicate whether Straddling AV valve was a factor.
Parent Long Name:	Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure
Parent Short Name:	PSFPrimProc
Parent Value(s):	= "VSD repair, Primary closure", "VSD repair, Patch", "VSD repair, Device", "Arterial switch operation (ASO) and VSD repair" or "Arterial switch procedure and VSD repair + Aortic arch repair"
<hr/>	
SeqNo: 952	
Long Name:	Procedure-Specific Factors - Major coronary crossing RVOT - Coronary anomaly restricting RVOT enlargement, (LAD from RCA etc.)
Short Name:	PSFMajCorRVOT
Database Table Name:	Operations
Data Source:	User
Format:	Text (categorical values specified by STS)
Definition:	Indicate whether Major coronary crossing RVOT - Coronary anomaly restricting RVOT enlargement, (LAD from RCA etc.) was a factor.
Parent Long Name:	Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure
Parent Short Name:	PSFPrimProc
Parent Value(s):	= "TOF - AVC (AVSD) repair", "TOF repair, No ventriculotomy", "TOF repair, Ventriculotomy, Nontransanular patch", "TOF repair, Ventriculotomy, Transanular Patch", "TOF repair, RV-PA conduit", "TOF - Absent pulmonary valve repair", "Pulmonary atresia
<hr/>	
SeqNo: 953	
Long Name:	Procedure-Specific Factors - VSD, Multiple, Repair
Short Name:	PSFVSDMultRep
Database Table Name:	Operations
Data Source:	User
Format:	Text (categorical values specified by STS)
Definition:	Indicate whether VSD, Multiple, Repair was a factor.
Parent Long Name:	Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure
Parent Short Name:	PSFPrimProc
Parent Value(s):	= "TOF - AVC (AVSD) repair", "TOF repair, No ventriculotomy", "TOF repair, entriculotomy, Nontransanular patch", "TOF repair, Ventriculotomy, Transanular patch", "TOF repair, RV-PA conduit", "TOF - Absent pulmonary valve repair", "Pulmonary atresia
<hr/>	
SeqNo: 954	
Long Name:	Procedure-Specific Factors - Restrictive VSD
Short Name:	PSFRestrictVSD
Database Table Name:	Operations
Data Source:	User



Format: Text (categorical values specified by STS)  
 Definition: Indicate whether Restrictive VSD was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors – Primary Procedure  
 Parent Short Name: PSFPrimProc  
 Parent Value(s): = "TOF - AVC (AVSD) repair", "TOF repair, No ventriculotomy", "TOF repair, Ventriculotomy, Nontransanular patch", "TOF repair, Ventriculotomy, Transanular patch", "TOF repair, RV-PA conduit", "TOF - Absent pulmonary valve repair", "Pulmonary atresia"

---

SeqNo: 955  
 Long Name: Procedure-Specific Factors - Hypoplastic branch pulmonary arteries (diminished pulmonary vascular bed)  
 Short Name: PSFHypoBrPulmArt  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether Hypoplastic branch pulmonary arteries (diminished pulmonary vascular bed) was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
 Parent Short Name: PSFPrimProc  
 Parent Value(s): = "TOF - AVC (AVSD) repair", "TOF repair, No ventriculotomy", "TOF repair, Ventriculotomy, Nontransanular patch", "TOF repair, Ventriculotomy, Transanular patch", "TOF repair, RV-PA conduit", "TOF - Absent pulmonary valve repair", "Pulmonary atresia"

---

SeqNo: 956  
 Long Name: Procedure-Specific Factors - AV Valve regurgitation grade 3 and 4 (Severe AV Valve regurgitation)  
 Short Name: PSFAVRegurg34  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether AV Valve regurgitation grade 3 and 4 (Severe AV Valve regurgitation) was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors – Primary Procedure  
 Parent Short Name: PSFPrimProc  
 Parent Value(s): = "TOF - AVC (AVSD) repair", "AVC (AVSD) repair, Complete (CAVSD)", "Bidirectional cavopulmonary anastomosis (BD CPA) (bidirectional Glenn)", "Glenn (unidirectional cavopulmonary anastomosis) (unidirectional Glenn)", "Bilateral bidirectional"

---

SeqNo: 957  
 Long Name: Procedure-Specific Factors - Double orifice left atrioventricular valve

---

Short Name: PSFDoubOrif  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether Double orifice left atrioventricular valve was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors – Primary Procedure  
 Parent Short Name: PSFPrimProc  
 Parent Value(s): = "TOF - AVC (AVSD) repair" or "AVC (AVSD) repair, Complete (CAVSD)"

---

SeqNo: 958  
 Long Name: Procedure-Specific Factors - Single papillary muscle in the left ventricle and/or parachute left atrioventricular valve  
 Short Name: PSFSingPap  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether Single papillary muscle in the left ventricle and/or parachute left atrioventricular valve was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors – Primary Procedure  
 Parent Short Name: PSFPrimProc  
 Parent Value(s): = "TOF - AVC (AVSD) repair" or "AVC (AVSD) repair, Complete (CAVSD)"

---

SeqNo: 959  
 Long Name: Procedure-Specific Factors - Hypoplastic posterior mural leaflet  
 Short Name: PSFHypoPostMLeaf  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether Hypoplastic posterior mural leaflet was a factor.  
 Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
 Parent Short Name: PSFPrimProc  
 Parent Value(s): = "TOF - AVC (AVSD) repair" or "AVC (AVSD) repair, Complete (CAVSD)"

---

SeqNo: 960  
 Long Name: Procedure-Specific Factors - Atrioventricular septal defect with ventricular imbalance: dominant left ventricle, hypoplastic right ventricle  
 Short Name: PSFASDDomLeft  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether Atrioventricular septal defect with ventricular imbalance: dominant left ventricle and hypoplastic right ventricle was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure

---



Parent Short Name: PSFPrimProc  
Parent Value(s): = "TOF - AVC (AVSD) repair" or "AVC (AVSD) repair, Complete (CAVSD)"

SeqNo: 961  
Long Name: Procedure-Specific Factors - Atrioventricular septal defect with ventricular imbalance: dominant right ventricle, hypoplastic left ventricle  
Short Name: PSFASDDomRight  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Atrioventricular septal defect with ventricular imbalance: dominant right ventricle and hypoplastic left ventricle was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "TOF - AVC (AVSD) repair" or "AVC (AVSD) repair, Complete (CAVSD)"

---

SeqNo: 962  
Long Name: Procedure-Specific Factors - Common atrioventricular valve with unbalanced commitment of valve to left ventricle  
Short Name: PSFCAVLeft  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Common atrioventricular valve with unbalanced commitment of valve to left ventricle was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "TOF - AVC (AVSD) repair" or "AVC (AVSD) repair, Complete (CAVSD)"

---

SeqNo: 963  
Long Name: Procedure-Specific Factors - Common atrioventricular valve with unbalanced commitment of valve to right ventricle  
Short Name: PSFCAVRight  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Common atrioventricular valve with unbalanced commitment of valve to right ventricle was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors – Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "TOF - AVC (AVSD) repair" or "AVC (AVSD) repair, Complete (CAVSD)"

---

SeqNo: 964  
Long Name: Procedure-Specific Factors - Moderate to severe systemic ventricular dysfunction  
Short Name: PSFModSevSVD  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Moderate to severe systemic ventricular dysfunction was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Bidirectional cavopulmonary anastomosis (BDCPA) (bidirectional Glenn)", "Glenn (unidirectional cavopulmonary anastomosis) (unidirectional Glenn)", "Bilateral bidirectional cavopulmonary anastomosis (BBDCPA) (bilateral bidirectional Glenn)", "HemiF"

---

SeqNo: 965  
Long Name: Procedure-Specific Factors - Systemic ventricular outflow tract obstruction (subaortic obstruction)  
Short Name: PSFSysVentObs  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Systemic ventricular outflow tract obstruction (subaortic obstruction) was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Bidirectional cavopulmonary anastomosis (BDCPA) (bidirectional Glenn)", "Glenn (unidirectional cavopulmonary anastomosis) (unidirectional Glenn)", "Bilateral bidirectional cavopulmonary anastomosis (BBDCPA) (bilateral bidirectional Glenn)", "HemiF"

---

SeqNo: 966  
Long Name: Procedure-Specific Factors - Ventricular dominance  
Short Name: PSFVentDom  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate ventricular dominance.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Bidirectional cavopulmonary anastomosis (BDCPA) (bidirectional Glenn)", "Glenn (unidirectional cavopulmonary anastomosis) (unidirectional Glenn)", "Bilateral bidirectional cavopulmonary anastomosis (BBDCPA) (bilateral bidirectional Glenn)",

HemiF

---

SeqNo: 970  
 Long Name: Procedure-Specific Factors - Posterior coronary loop: circumflex coming off the RCA  
 Short Name: PSFPostLoopCirc  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether Posterior coronary loop: circumflex coming off the RCA was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors – Primary Procedure  
 Parent Short Name: PSFPrimProc  
 Parent Value(s): = "Arterial switch operation (ASO)", "Arterial switch procedure + Aortic arch repair",  
 "Arterial switch operation (ASO) and VSD repair" or "Arterial switch procedure and  
 VSD repair + Aortic arch repair"

---

SeqNo: 971  
 Long Name: Procedure-Specific Factors - Posterior Coronary Loop: left trunk coming off the RCA  
 Short Name: PSFPostLoopLeftTrunc  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether Posterior Coronary Loop: left trunk coming off the RCA was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
 Parent Short Name: PSFPrimProc  
 Parent Value(s): = "Arterial switch operation (ASO)", "Arterial switch procedure + Aortic arch repair",  
 "Arterial switch operation (ASO) and VSD repair" or "Arterial switch procedure and  
 VSD repair + Aortic arch repair"

---

SeqNo: 972  
 Long Name: Procedure-Specific Factors - Double Coronary Loops  
 Short Name: PSFDoubleLoops  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether Double Coronary Loops (inverted origin of right and left coronary  
 arteries) was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
 Parent Short Name: PSFPrimProc  
 Parent Value(s): = "Arterial switch operation (ASO)", "Arterial switch procedure + Aortic arch repair",  
 "Arterial switch operation (ASO) and VSD repair" or "Arterial switch procedure and

---

VSD repair + Aortic arch repair"

---

SeqNo: 973

Long Name: Procedure-Specific Factors - Single Coronary Ostium

Short Name: PSFSingOst

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether Single coronary ostium was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors – Primary Procedure

Parent Short Name: PSFPrimProc

Parent Value(s): = "Arterial switch operation (ASO)", "Arterial switch procedure + Aortic arch repair",  
"Arterial switch operation (ASO) and VSD repair" or "Arterial switch procedure and  
VSD repair + Aortic arch repair"

---

SeqNo: 974

Long Name: Procedure-Specific Factors - Intramural coronary

Short Name: PSFIntramuralCor

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether Intramural coronary was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure

Parent Short Name: PSFPrimProc

Parent Value(s): = "Arterial switch operation (ASO)", "Arterial switch procedure + Aortic arch repair",  
"Arterial switch operation (ASO) and VSD repair" or "Arterial switch procedure and  
VSD repair + Aortic arch repair"

---

SeqNo: 975

Long Name: Procedure-Specific Factors - Large infundibular coronary artery from LAD

Short Name: PSFLgInfundArt

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether Large infundibular coronary artery from LAD was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure

Parent Short Name: PSFPrimProc

Parent Value(s): = "Arterial switch operation (ASO)", "Arterial switch procedure + Aortic arch repair",  
"Arterial switch operation (ASO) and VSD repair" or "Arterial switch procedure and  
VSD repair + Aortic arch repair"

---

---

SeqNo: 976  
Long Name: Procedure-Specific Factors - Malaligned commissures  
Short Name: PSFMalComm  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Malaligned commissures was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors – Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Arterial switch operation (ASO)", "Arterial switch procedure + Aortic arch repair",  
"Arterial switch operation (ASO) and VSD repair" or "Arterial switch procedure and  
VSD repair + Aortic arch repair"

---

SeqNo: 977  
Long Name: Procedure-Specific Factors - Take down of a commissure  
Short Name: PSFTakeDownComm  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Take down of a commissure was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Arterial switch operation (ASO)", "Arterial switch procedure + Aortic arch repair",  
"Arterial switch operation (ASO) and VSD repair" or "Arterial switch procedure and  
VSD repair + Aortic arch repair"

---

SeqNo: 978  
Long Name: Procedure-Specific Factors - Aorto-pulmonary diameter mismatch  
Short Name: PSFAortoPulMis  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Aorto-pulmonary diameter mismatch was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Arterial switch operation (ASO)", "Arterial switch procedure + Aortic arch repair",  
"Arterial switch operation (ASO) and VSD repair" or "Arterial switch procedure and  
VSD repair + Aortic arch repair"

---

SeqNo: 979  
Long Name: Procedure-Specific Factors - Side by side vessels  
Short Name: PSFSideBySide  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Side by side vessels was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Arterial switch operation (ASO)", "Arterial switch procedure + Aortic arch repair",  
"Arterial switch operation (ASO) and VSD repair" or "Arterial switch procedure and  
VSD repair + Aortic arch repair"

---

SeqNo: 980  
Long Name: Procedure-Specific Factors - Posterior native aorta  
Short Name: PSFPostNatAorta  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Posterior native aorta was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors – Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Arterial switch operation (ASO)", "Arterial switch procedure + Aortic arch repair",  
"Arterial switch operation (ASO) and VSD repair" or "Arterial switch procedure and  
VSD repair + Aortic arch repair"

---

SeqNo: 981  
Long Name: Procedure-Specific Factors - Subaortic obstruction/ conal septum malalignment  
Short Name: PSFSubAObs  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Subaortic obstruction / conal septum malalignment was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Arterial switch operation (ASO)", "Arterial switch procedure + Aortic arch repair",  
"Arterial switch operation (ASO) and VSD repair" or "Arterial switch procedure and  
VSD repair + Aortic arch repair"

---

SeqNo: 982  
Long Name: Procedure-Specific Factors - Bicuspid native aortic valve (Bicuspid neopulmonary valve)  
Short Name: PSFBicusNatAortic  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Bicuspid native aortic valve (Bicuspid neopulmonary valve) was a factor.  
Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors – Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Arterial switch operation (ASO)", "Arterial switch procedure + Aortic arch repair",  
"Arterial switch operation (ASO) and VSD repair" or "Arterial switch procedure and  
VSD repair + Aortic arch repair"

---

SeqNo: 983  
Long Name: Procedure-Specific Factors - Bicuspid native pulmonary valve (Bicuspid neo-aortic valve)  
Short Name: PSFBicusNatPulm  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Bicuspid native pulmonary valve (Bicuspid neo-aortic valve) was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Arterial switch operation (ASO)", "Arterial switch procedure + Aortic arch repair",  
"Arterial switch operation (ASO) and VSD repair" or "Arterial switch procedure and  
VSD repair + Aortic arch repair"

---

SeqNo: 984  
Long Name: Procedure-Specific Factors - Truncus type 3 (PA Branches from PDA or descending aorta)  
Short Name: PSFTruncType3  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Truncus type 3 (PA Branches from PDA or descending aorta) was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Truncus arteriosus repair" or "Truncus + Interrupted aortic arch repair (IAA)"

---

SeqNo: 985  
Long Name: Procedure-Specific Factors - Abnormal coronary

---

Short Name: PSFAbnormalCor  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Abnormal coronary was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors – Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Truncus arteriosus repair" or "Truncus + Interrupted aortic arch repair (IAA)"

SeqNo: 986  
Long Name: Procedure-Specific Factors - Truncal valve regurgitation (moderate to severe)  
Short Name: PSFTruncValRegurg  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Truncal valve regurgitation (moderate to severe) was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors – Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Truncus arteriosus repair" or "Truncus + Interrupted aortic arch repair (IAA)"

---

SeqNo: 987  
Long Name: Procedure-Specific Factors - Truncal Valve stenosis (moderate to severe)  
Short Name: PSFTruncValSten  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Truncal valve stenosis (moderate to severe) was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors – Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Truncus arteriosus repair" or "Truncus + Interrupted aortic arch repair (IAA)"

---

SeqNo: 988  
Long Name: Procedure-Specific Factors - Source of pulmonary blood flow: Shunt - systemic artery-to-pulmonary artery  
Short Name: PSFSrcPulFloShuntSys  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)

---



**Definition:** Indicate whether Source of pulmonary blood flow: Shunt - systemic artery-to-pulmonary artery was a factor.

**Parent Long Name:** Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure

**Parent Short Name:** PSFPrimProc

**Parent Value(s):** = "Norwood procedure"

---

**SeqNo:** 989

**Long Name:** Procedure-Specific Factors - Source of pulmonary blood flow: Shunt - ventricle-to-pulmonary artery

**Short Name:** PSFSrcPulFloShuntVent

**Database Table Name:** Operations

**Data Source:** User

**Format:** Text (categorical values specified by STS)

**Definition:** Indicate whether Source of pulmonary blood flow: Shunt - ventricle-to-pulmonary artery was a factor.

**Parent Long Name:** Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure

**Parent Short Name:** PSFPrimProc

**Parent Value(s):** = "Norwood procedure"

---

**SeqNo:** 990

**Long Name:** Procedure-Specific Factors - Source of pulmonary blood flow: Superior caval vein-to-pulmonary

**Short Name:** PSFSrcPulFloSuper

**Database Table Name:** Operations

**Data Source:** User

**Format:** Text (categorical values specified by STS)

**Definition:** Indicate whether Source of pulmonary blood flow: Superior caval vein-to-pulmonary artery was a factor.

**Parent Long Name:** Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure

**Parent Short Name:** PSFPrimProc

**Parent Value(s):** = "Norwood procedure"

---

**SeqNo:** 991

**Long Name:** Procedure-Specific Factors - Ascending aorta < 2 mm

**Short Name:** PSFAscAortaLT2

**Database Table Name:** Operations

**Data Source:** User

**Format:** Text (categorical values specified by STS)

**Definition:** Indicate whether Ascending aorta < 2 mm was a factor.

---

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Norwood procedure"

---

SeqNo: 992  
Long Name: Procedure-Specific Factors - Aortic atresia  
Short Name: PSFAortAtresia  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Aortic atresia was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Norwood procedure"

---

SeqNo: 993  
Long Name: Procedure-Specific Factors - Aortic stenosis  
Short Name: PSFAortSten  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Aortic stenosis was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Norwood procedure"

---

SeqNo: 994  
Long Name: Procedure-Specific Factors - Mitral atresia  
Short Name: PSFMitralAtresia  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Mitral atresia was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Norwood procedure"

---

SeqNo: 995

---

Long Name: Procedure-Specific Factors - Mitral stenosis  
Short Name: PSFMitralSten  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Mitral stenosis was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Norwood procedure"

---

SeqNo: 996  
Long Name: Procedure-Specific Factors - Sinusoids  
Short Name: PSFSinusoids  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the presence of sinusoids was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Norwood procedure"

---

SeqNo: 997  
Long Name: Procedure-Specific Factors - Intact atrial septum  
Short Name: PSFIntactAtrSep  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Intact atrial septum was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Norwood procedure"

---

SeqNo: 998  
Long Name: Procedure-Specific Factors - Obstructed pulmonary venous return with severely restrictive ASD  
Short Name: PSFObsPulVenRet  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)

---

**Definition:** Indicate whether Obstructed pulmonary venous return with severely restrictive ASD was a factor.

**Parent Long Name:** Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure

**Parent Short Name:** PSFPrimProc

**Parent Value(s):** = "Norwood procedure"

---

**SeqNo:** 999

**Long Name:** Procedure-Specific Factors - Aberrant right subclavian artery

**Short Name:** PSFAberrantRtSubclav

**Database Table Name:** Operations

**Data Source:** User

**Format:** Text (categorical values specified by STS)

**Definition:** Indicate whether Aberrant right subclavian artery was a factor.

**Parent Long Name:** Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure

**Parent Short Name:** PSFPrimProc

**Parent Value(s):** = "Norwood procedure"

---

**SeqNo:** 1001

**Long Name:** Procedure-Specific Factors - TV Repair

**Short Name:** PSFTVRep

**Database Table Name:** Operations

**Data Source:** User

**Format:** Text (categorical values specified by STS)

**Definition:** Indicate whether TV Repair was a factor.

**Parent Long Name:** Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure

**Parent Short Name:** PSFPrimProc

**Parent Value(s):** = "Ebstein's repair"

---

**SeqNo:** 1002

**Long Name:** Procedure-Specific Factors - TV Repair - Monocusp

**Short Name:** PSFTVRepMono

**Database Table Name:** Operations

**Data Source:** User

**Format:** Text (categorical values specified by STS)

**Definition:** Indicate whether TV Repair - Monocusp was a factor.

**Parent Long Name:** Procedure-Specific Factors - TV Repair

**Parent Short Name:** PSFTVRep

**Parent Value(s):** = "Yes"

---

12/2015: If a patient has a VAD, and then gets a heart transplant and has the VAD removed during the same operation, what should the OpType be coded as? According to the STS definition of VAD with CPB, it says that it includes operations to insert or remove the VAD. So should it be VAD with CPB or should it just be CPB? **It should be CPB. Even if patient comes in on VAD or ECMO it is still CPB.**

February 2016: I am unclear about coding the operation type for a patient whose procedure was aborted after skin incision. The only thing that was performed was a thoracotomy. The patient decompensated after skin incision and nothing further was done. Should this be coded as operation type: Thoracic or as the originally intended procedure of NO CPB Cardiovascular? **Code as Op Type-Other.**

February 2016: Mortality case where 1st procedure, primary procedure was VAD w/CPB (salvage case, as pt was on ECMO at time of OR). Additionally had ASD repair at time of VAD implant(atrial fenestration closure as atrial septectomy performed at bedside prior to VAD not by our surgeons). Op type was VAD w/ CPB. 2nd procedure ECMO decan and delayed sternal closure, which was coded as ECMO procedure. Question is once case goes to DCRI for analysis, will the primary procedure be changed to ASD repair instead of VAD implant which will change the Op type to CPB case and will move this in mortality analysis. **DCRI does not change the OP type. VAD=only cases that go to OR for VAD.**

SeqNo: 1004

Long Name: Procedure-Specific Factors - TV Repair - Bileaflet Repair  
 Short Name: PSFTVRepBileaf  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether TV Repair - Bileaflet Repair was a factor.

Parent Long Name: Procedure-Specific Factors - TV Repair  
 Parent Short Name: PSFTVRep  
 Parent Value(s): = "Yes"

SeqNo: 1006

Long Name: Procedure-Specific Factors - TV Repair - Cone Repair - 360 Degrees Leaflet Approximation  
 Short Name: PSFTVRepCone  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether TV Repair - Cone Repair - 360 Degrees Leaflet Approximation was a factor.

Parent Long Name: Procedure-Specific Factors - TV Repair  
 Parent Short Name: PSFTVRep  
 Parent Value(s): = "Yes"

SeqNo: 1008

Long Name: Procedure-Specific Factors - Sebening Stitch (Anterior RV Papillary Muscle To Ventricular Septum)

Short Name: PSFSebening

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether Sebening Stitch (Anterior RV Papillary Muscle To Ventricular Septum) was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors – Primary Procedure

Parent Short Name: PSFPrimProc

Parent Value(s): = "Ebstein's repair"

---

SeqNo: 1009

Long Name: Procedure-Specific Factors - Annular Reduction

Short Name: PSFAnnRed

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether Annular Reduction was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure

Parent Short Name: PSFPrimProc

Parent Value(s): = "Ebstein's repair"

---

SeqNo: 1010

Long Name: Procedure-Specific Factors - Annular Reduction - Plication

Short Name: PSFAnnRedPlic

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether Annular Reduction - Plication was a factor.

Parent Long Name: Procedure-Specific Factors - Annular Reduction

Parent Short Name: PSFAnnRed

Parent Value(s): = "Yes"

---

SeqNo: 1012

Long Name: Procedure-Specific Factors - Annular Reduction - Partial Ring (C-Shaped Anterior And Inferior

Short Name: PSFAnnRedPartial

Database Table Name: Operations

---

Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Annular Reduction - Partial Ring (C-Shaped Anterior And Inferior Annulus) was a factor.

Parent Long Name: Procedure-Specific Factors - Annular Reduction  
Parent Short Name: PSFAnnRed  
Parent Value(s): = "Yes"

---

SeqNo: 1014  
Long Name: Procedure-Specific Factors - Annular Reduction - Eccentric Ring (Inferior Annulus)  
Short Name: PSFAnnRedEccent  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Annular Reduction - Eccentric Ring (Inferior Annulus) was a factor.

Parent Long Name: Procedure-Specific Factors - Annular Reduction  
Parent Short Name: PSFAnnRed  
Parent Value(s): = "Yes"

---

SeqNo: 1016  
Long Name: Procedure-Specific Factors - Atrialized RV Plication  
Short Name: PSFAtrialRVPlc  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Atrialized RV Plication was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Ebstein's repair"

---

SeqNo: 1018  
Long Name: Procedure-Specific Factors - Atrialized RV Resection  
Short Name: PSFAtrialRVRes  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Atrialized RV Resection was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure  
Parent Short Name: PSFPrimProc

---

Parent Value(s): = "Ebstein's repair"

---

SeqNo: 1020

Long Name: Procedure-Specific Factors - ASD/PFO Closure

Short Name: PSFASDPFO

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether ASD/PFO Closure was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure

Parent Short Name: PSFPrimProc

Parent Value(s): = "Ebstein's repair"

---

SeqNo: 1022

Long Name: Procedure-Specific Factors - Reduction Atrioplasty

Short Name: PSFRedAtrio

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether Reduction Atrioplasty was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure

Parent Short Name: PSFPrimProc

Parent Value(s): = "Ebstein's repair"

---

SeqNo: 1023

Long Name: Procedure-Specific Factors - Arrhythmia Surgery

Short Name: PSFArrSurg

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether Arrhythmia Surgery was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors - Primary Procedure

Parent Short Name: PSFPrimProc

Parent Value(s): = "Ebstein's repair"

---

SeqNo: 1024

Long Name: Procedure-Specific Factors - Arrhythmia Surgery - Cavotricuspid Isthmus Ablation

---



Short Name: PSFArrSurgCavo  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Arrhythmia Surgery - Cavotricuspid Isthmus Ablation was a factor.  
  
Parent Long Name: Procedure-Specific Factors - Arrhythmia Surgery  
Parent Short Name: PSFArrSurg  
Parent Value(s): = "Yes"

---

SeqNo: 1026  
Long Name: Procedure-Specific Factors - Arrhythmia Surgery - Modified Right Atrial Maze  
Short Name: PSFArrSurgModMaze  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Arrhythmia Surgery - Modified Right Atrial Maze was a factor.  
  
Parent Long Name: Procedure-Specific Factors - Arrhythmia Surgery  
Parent Short Name: PSFArrSurg  
Parent Value(s): = "Yes"

---

SeqNo: 1028  
Long Name: Procedure-Specific Factors - Arrhythmia Surgery - Left Atrial Cox Maze  
Short Name: PSFArrSurgCoxMaze  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Arrhythmia Surgery - Left Atrial Cox Maze was a factor.  
  
Parent Long Name: Procedure-Specific Factors - Arrhythmia Surgery  
Parent Short Name: PSFArrSurg  
Parent Value(s): = "Yes"

---

SeqNo: 1030  
Long Name: Procedure-Specific Factors - Arrhythmia Surgery - Pulmonary Vein Isolation  
Short Name: PSFArrSurgPulmIso  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Arrhythmia Surgery - Pulmonary Vein Isolation was a factor.  
  
Parent Long Name: Procedure-Specific Factors - Arrhythmia Surgery

---

Parent Short Name: PSFArrSurg  
Parent Value(s): = "Yes"

---

SeqNo: 1032  
Long Name: Procedure-Specific Factors - Bidirectional Cavopulmonary Anastomosis  
Short Name: PSFBiCavoAnast  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Bidirectional Cavopulmonary Anastomosis was a factor.

Parent Long Name: Procedure-Specific Factors - Procedure-Specific Factors – Primary Procedure  
Parent Short Name: PSFPrimProc  
Parent Value(s): = "Ebstein's repair"

---

## 11. Operative

SeqNo: 1054  
Long Name: Procedure Location  
Short Name: ProcLoc  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the location where the operation/procedure was performed.

---

SeqNo: 1055  
Long Name: Status  
Short Name: Status  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the clinical status of the patient prior to entering the operating room.

---

SeqNo: 1056  
Long Name: Operation Type  
Short Name: OpType  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the type of primary surgical procedure performed.

---

Harvest Codes and Value Definitions:

Code:	Value:	Definition:
1	CPB Cardiovascular	<p>If the procedure is cardiovascular (includes the heart, great vessels, or any branches of the great vessels), and cardiopulmonary bypass is used, this should be chosen as the case category. Do not choose this case category for operations that are not cardiovascular, even if cardiopulmonary bypass is used (<i>see Op Type 9, below</i>)</p> <p>Most lung transplants involve anastomosis to the left atrium (as well as anastomosis to distal main PA or central branch PA). I would consider this a cardiovascular procedure. Transplant, Lung(s) is a STAT Category 3 procedure</p> <p><del>If cardiopulmonary bypass is used, this must be chosen as the case category whether the procedure is thoracic (e.g., tracheal reconstruction) or cardiovascular in nature.</del></p>
2	No CPB Cardiovascular	<p>If the procedure is cardiovascular, but cardiopulmonary bypass is not used, this must be chosen as the case category. This includes any procedure that includes the heart, great vessels, or any of the branches from the great vessels, where CPB is not used. Examples include but are not limited to: coarctation of the aorta repair, creation of a systemic-to-pulmonary artery shunt, patent ductus arteriosus ligation. A delayed sternal closure is included in this category. <del>If a pericardial window done for cancer, it should be classified as a Cardiac Operation (Operation type = No CPB Cardiovascular).</del></p> <ul style="list-style-type: none"> <li><i>pericardial drainage/pericardial window procedure for cancer = Thoracic Procedure</i></li> <li><i>pericardial drainage/pericardial window procedure for cardiac disease = No CPB Cardiovascular</i></li> </ul>
9	CPB Non-Cardiovascular	<p>Procedures that are done with bypass support that do not involve a concomitant cardiovascular procedure. For example, tracheal surgery, neurosurgical procedures, resuscitation and rewarming of drowning victims. These cases are not included in the numerator or denominator of mortality calculations or reports. Tracheal reconstructions done on CPB, without a concomitant cardiovascular procedure are Op Type 9 - CPB Non-cardiovascular. This would pertain, for example to a slide tracheoplasty or tracheal patch-plasty done on CPB. But, if the operation also includes a cardiovascular procedure (as in operation for PA sling with both tracheal repair and</p>

		division/reimplantation of pulmonary artery), then it would be CPB Cardiovascular.
3	ECMO	If ECMO cannulation or decannulation is the primary procedure performed, this category must be chosen. However, if ECMO is initiated for support at the end of another type procedure (i.e., CPB, No CPB Cardiovascular), that procedure takes precedence and the category code would not be ECMO.
4	Thoracic	If a procedure is performed on a structure within the chest cavity but does not involve the cardiac chambers or vessels, it would be a Thoracic category case (for example, lobectomy, pectus excavatum/carinatum repair, anterior spine exposure). There will be thoracic cases that require cardiopulmonary bypass (e.g., some types of tracheal reconstructions). In those cases, the use of cardiopulmonary bypass takes precedence and the case would not be Thoracic, but <del>CPB</del> rather CPB Non-Cardiovascular.
5	Interventional Cardiology	If an interventional device (e.g., occluder, stent) is placed in the operating room as the primary procedure performed, this category must be chosen. However, if in the course of another type procedure (i.e., CPB, No CPB Cardiovascular), an interventional device is placed in addition to the other procedure, the other category takes precedence and the case would not be Interventional Cardiology.
6	VAD Operation Done With CPB	Ventricular Assist Device procedure done with CPB. This includes operations to insert the VAD or to remove the VAD
7	VAD Operation Done Without CPB	Ventricular Assist Device procedure done without CPB. This includes operations to insert the VAD, to remove the VAD, or any procedure performed while on the VAD.
8	Non-cardiac, Non-thoracic procedure on cardiac patient with cardiac anesthesia	Any non-cardiac or non-thoracic procedure such as a general surgical procedure with anesthesia provided by cardiac anesthesiology because of the patient's underlying cardiac physiology.
777	Other	All other procedures that do not fall within the above definitions should be coded as category Other. This would include but not be limited to supportive minor procedures (e.g., line placements)

---

#### ECMO or CPB Examples

Procedures performed while a patient is on ECMO can be coded as Op Type "ECMO" ***if they are done exclusively for the purpose of facilitating ECMO support.***

A patient is admitted to the hospital and requires emergent ECMO cannulation. The next day the patient is taken to the OR for BT Shunt placement done on ECMO. Op Type?

**If the BT Shunt is done on ECMO, this should be coded as Op Type CPB because the ECMO circuit is functioning as a CPB circuit in this situation.**

A patient who winds up on ECMO and has a shunt revision while on ECMO support is a *different type of scenario*, and under those circumstances it is most likely that the procedure “shunt revision” *should be considered to be of Op Type CPB, with the understanding that ECMO circuit is being used to provide CPB support.*

If the patient was transitioned from ECMO to bypass in the OR and then transitioned back to ECMO at the end of the case you would code the **op type as CPB with the pre-op risk factor of ECMO and the post-op complication of ECMO.**

If the patient was de-cannulated from ECMO prior to the placement of the shunt and the shunt was done with no support you would code the Op Type as **No CPB Cardiovascular.**

Patient arrives in CVICU and needs cannulation on ECMO. The consent and discussion with the family is for ECMO. Patient is cannulated for ECMO in the OR and during the procedure it is noted that the patient has excessive pulmonary blood flow and needs a PAB to help control pulmonary blood flow and the patient is better supported on ECMO.

**This situation would be coded as ECMO.**

Patient has bleeding requiring mediastinal exploration while on ECMO.

**This situation would be coded as ECMO.**

Patient returns to the OR for an unbalanced AVC repair while on ECMO. The consent and the operative report note that the case was for the repair. Case completed and the patient returns to the CVICU on ECMO.

**This situation is coded as a CPB Cardiovascular case since the case is a cardiovascular procedure even if the patient returns to the CVICU on ECMO. This would also be the index procedure for the patient.**

Patient arrives in the CVICU. The patient needs a PAB. The consent and the operative report identify that the patient is going to have a PAB. During the procedure the patient is identified to need ECMO as well.

**This is coded as a CV case with or without CPB depending on the operative report, if the PAB was done while on ECMO (cannulation occurred before the PAB) the op type is CPB.**

Patient arrives in CVICU and needs cannulation for ECMO. The consent and discussion with the family is for ECMO. The patient is cannulated for ECMO.

**This is coded as ECMO.**

Patient is decannulated from ECMO. Chest is closed.

**This is coded as ECMO.**

**FAQs:**

August 2016: We have a non-congenital heart kid who was put on ECMO by general surgery. Days later our congenital surgeons re-cannulated him, repaired a nick in his atrium and drained a pericardial effusion, both due to his initial cannulation. Since he has a normal heart and was already on ECMO, would this be considered Op Type ECMO, or CPB? **This is Op Type ECMO.**

---

March 2017: ECMO only was done via a small transverse incision superior and to the right clavicle. When it was removed, the removal was done at bedside. Is this type of ECMO procedure supposed to be captured, since it sounds like it was a percutaneous insertion and not an actual surgical procedure? **If the cardiac surgeons performed the procedure and the patient was put on ECMO, include as an ECMO procedure in the database.**

---

August 2016: Patient is admitted for wound drainage/debridement, sternum is left open planned. Patient then undergoes a 'delayed sternal closure'. What op type would you code for the 'delayed sternal closure'? **This is Op Type Thoracic. In this case the procedure was not related to the heart.**

August 2016: A patient had a PDA closure and a tracheal procedure done. Cardiopulmonary bypass was initiated utilizing an ECMO circuit for the procedure. ECMO was discontinued at the end of the procedure. Should the optype for this procedure be CPB cardio? **If ECMO was initiated exclusively for the procedure, code either Op Type ECMO or Op Type CPB, non-cardiovascular. Only code this as a cardiovascular procedure if the main reason to undertake the procedure was to close the ductus.**

October 2016: A patient is admitted to the hospital and requires emergent ECMO cannulation. The following day the patient is taken to the OR for BT shunt placement done on ECMO.

- 1.) What should the op type be coded as? **If the BT Shunt is done on ECMO, we should code as Op Type = CPB because the ECMO circuit is functioning as a CPB circuit in this setting.**
  - 2.) If the patient was transitioned from ECMO to bypass in the OR and then transitioned back to ECMO at the end of the case would you code op type as CPB with preop risk factor of ECMO and postop complication of ECMO? **Yes. Op Type = CPB, preop risk factor of ECMO and postop complication of ECMO**
  - 3.) If the patient was de-cannulated from ECMO prior to the placement of the shunt and the shunt was done with no support would you code Op Type as No CPB? **Yes. Op Type = No CPB Cardiovascular**
- How should CPB information be entered for cases done on ECMO? **Use the surgery start and end times as minutes on pump.**

March 2017: If a patient is on ECMO and a mediastinal exploration or other mediastinal procedure is performed, does this count as an ECMO procedure or a nonCPB procedure? When this was asked as a question on the conference call, approximately half of the centers were entering these cases as ECMO and the other half as non-CPB. **If major structural repairs were completed while on ECMO, the operation type is CPB. Otherwise, minor procedures done while on ECMO would remain operation type ECMO (e.g.; mediastinal explorations).**

April 2017: I have a question regarding assignment of the Operation Type for a patient. The patient had DDD pacemaker implanted and VAD. You normally do not see these two procedures together, so I am wondering should I go with "no CPB cardiovascular" for the pacemaker or "VAD operation with CPB" or alternate option may also be "CPB", as I wanted to show that the patient was on CPB? **Was CPB used to put the VAD in, then Operation type VAD with CPB; if CPB not used the operation type is VAD without CPB.**

April 2017: A patient was transferred in on ECMO for a VAD implant as a bridge to transplant. The patient had a total of three surgeries with us. First surgery was Berlin Heart implant/ecmo decannulation. Second surgery was delayed sternal closure. Third surgery was mediastinal exploration for bleeding. Several days later, patient suffered a subdural bleed, parents withdrew support and the patient died. What should be the operation types for each of these procedures? Which one is the index? The VAD cases are difficult. What if this patient didn't

expire and had a successful heart transplant as the 4th surgery and was discharged to home. Which operation would be the index procedure?

1. VAD with or without CPB depending on what was done
2. VAD without CPB
3. VAD without CPB

As VAD cases are not included in analysis, there really is not an index operation.

If the patient hadn't expired, the first three surgeries would remain VAD cases. However, the 4<sup>th</sup> surgery for heart transplant would be Op Type: CPB Cardiovascular, and become the patient's index operation.

July 2017: Patient's procedure was resection of left radiocephalic AV fistula and repair of left radial artery with patch. I am not sure what operation type would be best. **Other procedure or Non-cardiac, Non-thoracic procedure on a cardiac patient – whichever fits best.**

September 2017: Patient (CG 3/8) had LVAD implantation, mitral and tricuspid valvuloplasty, and conduit replacement. Is this OpType VAD w/ CPB, or CPB cardiovascular? **CPB Cardiovascular**

October 2017: We have a patient admitted with diagnosis of Ebstein's anomaly in cardiogenic shock and cannulated to ECMO. Subsequently taken to the OR for a Mee shunt and ECMO decannulation. Three weeks later taken to the OR for Cone procedure. What should the operation type be for the Mee shunt? If it is anything but ECMO, the Cone procedure will not be the index case for this admission. If this is acceptable with everyone, then so be it. **It is a CPB case and the Mee Shunt is the index case.**

SeqNo: 1057

Long Name: Near Infrared Spectroscopy (NIRS) Cerebral Metrics Used  
 Short Name: NIRSCerUsed  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether cerebral oximetry was monitored.

---

SeqNo: 1058

Long Name: Near Infrared Spectroscopy (NIRS) Cerebral Metrics Used - Preoperatively  
 Short Name: NIRSCerPre  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether cerebral oximetry was monitored during the preoperative period.

Parent Long Name: Near Infrared Spectroscopy (NIRS) Cerebral Metrics Used  
 Parent Short Name: NIRSCerUsed  
 Parent Value(s): = "Yes"

---

SeqNo: 1059

Long Name: Near Infrared Spectroscopy (NIRS) Cerebral Metrics Used - Intraoperatively

---

Short Name: NIRSCerIntra  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether cerebral oximetry was monitored during the intraoperative period.

Parent Long Name: Near Infrared Spectroscopy (NIRS) Cerebral Metrics Used  
Parent Short Name: NIRSCerUsed  
Parent Value(s): = "Yes"

---

SeqNo: 1060  
Long Name: Near Infrared Spectroscopy (NIRS) Cerebral Metrics Used - Postoperatively  
Short Name: NIRSCerPost  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether cerebral oximetry was monitored during the postoperative period.

Parent Long Name: Near Infrared Spectroscopy (NIRS) Cerebral Metrics Used  
Parent Short Name: NIRSCerUsed  
Parent Value(s): = "Yes"

---

SeqNo: 1061  
Long Name: Near Infrared Spectroscopy (NIRS) Somatic Metrics Used  
Short Name: NIRSSomUsed  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether somatic oximetry was monitored.

---

SeqNo: 1062  
Long Name: Near Infrared Spectroscopy (NIRS) Somatic Metrics Used - Preoperatively  
Short Name: NIRSSomPre  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether somatic oximetry was monitored during the preoperative period.

Parent Long Name: Near Infrared Spectroscopy (NIRS) Somatic Metrics Used  
Parent Short Name: NIRSSomUsed  
Parent Value(s): = "Yes"

---



---

SeqNo: 1063  
Long Name: Near Infrared Spectroscopy (NIRS) Somatic Metrics Used - Intraoperatively  
Short Name: NIRSSomIntra  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether somatic oximetry was monitored during the intraoperative period.

Parent Long Name: Near Infrared Spectroscopy (NIRS) Somatic Metrics Used  
Parent Short Name: NIRSSomUsed  
Parent Value(s): = "Yes"

---

SeqNo: 1064  
Long Name: Near Infrared Spectroscopy (NIRS) Somatic Metrics Used - Postoperatively  
Short Name: NIRSSomPost  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether somatic oximetry was monitored during the postoperative period.

Parent Long Name: Near Infrared Spectroscopy (NIRS) Somatic Metrics Used  
Parent Short Name: NIRSSomUsed  
Parent Value(s): = "Yes"

---

SeqNo: 1065  
Long Name: Time Patient Entered the OR  
Short Name: OREntryT  
Database Table Name: Operations  
Data Source: User  
Format: Time - hh:mm (24-hour clock)  
Definition: Indicate to the nearest minute (using 24-hour clock) the time the patient entered the OR. If the procedure was performed in a location other than the OR, record the time when the sterile field was set up.

---

SeqNo: 1066  
Long Name: Skin Incision Start Time  
Short Name: SStartT  
Database Table Name: Operations  
Data Source: User  
Format: Time - hh:mm (24-hour clock)  
Definition: Indicate to the nearest minute (using 24-hour clock) the time the skin incision was

made.

---

SeqNo: 1067

Long Name: Endotracheal Intubation was Performed

Short Name: Intubate

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether an endotracheal intubation was performed.

---

SeqNo: 1068

Long Name: Intubation Date and Time

Short Name: IntubateDT

Database Table Name: Operations

Data Source: User

Format: Date/Time - mm/dd/yyyy hh:mm

Definition: Indicate the date (mm/dd/yyyy) and time (hh:mm) (24 hour clock) ventilatory support started. Capture the intubation closest to the surgical start time.

If the patient was intubated upon admission and remained intubated until the surgical start time, capture this intubations date and time.

If the patient was admitted intubated (intubated at another institution) and remained continually intubated until the surgical start time, capture the patient's admission date and time.

If the patient was admitted with a tracheostomy in place without ventilatory support, capture the date and time closest to the surgical start time that ventilatory support was initiated.

If the patient was admitted with a tracheostomy in place receiving chronic ventilatory support, capture the admission date and time.

If the intubation date and time is otherwise unknown, enter the date and time the patient entered the operating room.

Do not alter the previously established date and time that ventilatory support was initiated for scenarios including, but not limited to, interruptions in ventilatory support due to accidental extubation /de-cannulation, elective tube change etc.

Parent Long Name: Endotracheal Intubation was Performed

Parent Short Name: Intubate

Parent Value(s): = "Yes"

---

SeqNo: 1069

Long Name: Initial Extubation Date and Time  
Short Name: ExtubateDT  
Database Table Name: Operations  
Data Source: User  
Format: Date/Time - mm/dd/yyyy hh:mm  
Definition: Indicate the date (mm/dd/yyyy) and time (hh:mm) (24 hour clock) ventilatory support initially ceased after surgery. Capture the extubation closest to the surgical stop time.

If the patient has a tracheostomy and is separated from the mechanical ventilator postoperatively within the hospital admission, capture the date and time of separation from the mechanical ventilator closest to the surgical stop time.

If the patient expires while intubated or /cannulated and on the ventilator, capture the date and time of expiration.

If patient discharged on chronic ventilatory support, capture the date and time of discharge.

Parent Long Name: Endotracheal Intubation was Performed  
Parent Short Name: Intubate  
Parent Value(s): = "Yes"

---

SeqNo: 1070

Long Name: Extubated In The Operating Room  
Short Name: ExtubInOR  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the endotracheal tube was removed in the OR or at the end of the procedure if it was performed in another location.

Parent Long Name: Endotracheal Intubation was Performed  
Parent Short Name: Intubate  
Parent Value(s): = "Yes"

---

February 2017: If the patient is extubated in the OR but is reintubated before leaving the OR, how should this question be answered? **'Yes' to #1070, extubated in OR. You would answer 'Yes' to #1017, reintubated after initial post-operative extubation also.**

---

SeqNo: 1071

---

Long Name: Re-Intubated After Initial Postoperative Extubation  
 Short Name: ReIntubate  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether the patient was re-intubated after the initial postoperative extubation.

Parent Long Name: Endotracheal Intubation was Performed  
 Parent Short Name: Intubate  
 Parent Value(s): = "Yes"

August 2016: If a patient is intubated for the congenital cardiac surgery and extubated following the surgery, then intubated again later for another procedure and extubated in the OR following the other procedure, is this counted as a reintubation? **Yes, this is counted as a re-intubation but it is not a complication. The patient was re-intubated for a procedure and not as a result of respiratory failure.**

SeqNo: 1072

Long Name: Final Extubation Date and Time  
 Short Name: FinExtubDT  
 Database Table Name: Operations  
 Data Source: User  
 Format: Date/Time - mm/dd/yyyy hh:mm  
 Definition: Indicate the date (mm/dd/yyyy) and time (hh:mm) (24 hour clock) ventilatory support last ceased prior to discharge after surgery. Capture the extubation time closest to discharge.

If the patient has a tracheostomy and is separated from the mechanical ventilator more than once postoperatively within the hospital admission, capture the date and time of separation from the mechanical ventilator closest to the hospital discharge.

If the patient expires while intubated or cannulated and on the ventilator capture the date and time of expiration.

If the patient was discharged on chronic ventilatory support capture the date and time of discharge.

Parent Long Name: Re-Intubated After Initial Postoperative Extubation  
 Parent Short Name: ReIntubate  
 Parent Value(s): = "Yes"

May 2017: How to document child who is trached and has no final extubation date. **The extubation date will be the date and time the patients discharged from the facility.**

SeqNo: 1073

Long Name: Time of Skin Closure

Short Name: SIStopT  
 Database Table Name: Operations  
 Data Source: User  
 Format: Time - hh:mm (24-hour clock)  
 Definition: Indicate to the nearest minute (using 24-hour clock) the time the skin incision was closed. If patient leaves the operating room with an open incision, collect the time dressings were applied to the incision.

---

SeqNo: 1074  
 Long Name: Time Patient Exited the OR  
 Short Name: ORExitT  
 Database Table Name: Operations  
 Data Source: User  
 Format: Time - hh:mm (24-hour clock)  
 Definition: Indicate to the nearest minute (using 24-hour clock) the time the patient exits the operating room. If the procedure was performed in a location other than the OR, record the time when the sterile field was taken down.

---

SeqNo: 1075  
 Long Name: Procedure Extended Through Midnight  
 Short Name: MultiDay  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether the procedure continued through midnight from one day to the next.

---

March 2017: Is Multiday determined by OR entry time or by skin incision start time? For example, the patient enters the OR at 23:19 on 10/24. Skin incision is at 00:51 on 10/25. SI stop is at 8:06 on 10/25, and OR exit is at 8:17 on 10/25. Is this multiday? **Yes, use OR entry and exit as the parameters for determination of multiday procedures.**

---

SeqNo: 1076  
 Long Name: Surgeon  
 Short Name: Surgeon  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by user)      Data Length: 100  
 Definition: Indicate the name of the primary surgeon performing this surgical procedure. The name, NPI and signature of all surgeons contributing data to the database must be on file with the STS for data files to be accepted.

Parent Long Name: Operation Type  
 Parent Short Name: OpType

---

Parent Value(s): = "CPB Cardiovascular", "No CPB Cardiovascular", "CPB Non-Cardiovascular", "ECMO", "Thoracic", "VAD Operation Done With CPB", "VAD Operation Done Without CPB." or "Other"

---

SeqNo: 1078

Long Name: Surgeon National Provider Identifier

Short Name: SurgNPI

Database Table Name: Operations

Data Source: Lookup

Format: Text

Definition: Indicate the individual-level National Provider Identifier (NPI) of the surgeon performing the procedure.

Parent Long Name: Operation Type

Parent Short Name: OpType

Parent Value(s): = "CPB Cardiovascular", "No CPB Cardiovascular", "CPB Non-Cardiovascular", "ECMO", "Thoracic", "VAD Operation Done With CPB", "VAD Operation Done Without CPB." or "Other"

---

SeqNo: 1079

Long Name: Taxpayer Identification Number

Short Name: TIN

Database Table Name: Operations

Data Source: User

Format: Text

Definition: Indicate the group-level Taxpayer Identification Number for the Taxpayer holder of record for the Surgeon's National Provider Identifier that performed the procedure.

Parent Long Name: Operation Type

Parent Short Name: OpType

Parent Value(s): = "CPB Cardiovascular", "No CPB Cardiovascular", "CPB Non-Cardiovascular", "ECMO", "Thoracic", "VAD Operation Done With CPB", "VAD Operation Done Without CPB." or "Other"

---

SeqNo: 1080

Long Name: Reoperation Within This Admission

Short Name: ReOpInAdm

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether this is a second, or third (or more) operation within the same hospital admission.

---

Parent Long Name: Operation Type  
 Parent Short Name: OpType  
 Parent Value(s): = "CPB Cardiovascular", "No CPB Cardiovascular", "CPB Non-Cardiovascular", "ECMO", "Thoracic", "VAD Operation Done With CPB", "VAD Operation Done Without CPB." or "Other"

August 2017: My surgeon performed a VSD/PFO closure. The patient was moved from the OR table but not out of the operating room when they noticed bleeding, placed him back on the table, opened the sternum and cauterized a bleeder. There is only one Op Note and start and stop times were not noted separately even though the skin was closed and reopened. Is this a reoperation for bleeding or would I list it as mediastinal exploration along with the VSD & PFO procedures? **One operative case, include mediastinal exploration and code the complication bleeding, requiring reoperation.**

SeqNo: 1090

Long Name: Number of Prior Cardiothoracic Operations  
 Short Name: PrvCtOpN  
 Database Table Name: Operations  
 Data Source: User  
 Format: Integer  
 Definition: Indicate, prior to this admission's surgical procedure, how many cardiothoracic (heart or great vessels) surgical procedures were performed with or without cardiopulmonary bypass (CPB). Also include lung procedures utilizing CPB or tracheal procedures utilizing CPB. See Operation Type for further clarification.

Low Value: 0 High Value: 200

Parent Long Name: Operation Type  
 Parent Short Name: OpType  
 Parent Value(s): = "CPB Cardiovascular", "No CPB Cardiovascular", "CPB Non-Cardiovascular", "ECMO", "Thoracic", "VAD Operation Done With CPB", "VAD Operation Done Without CPB." or "Other"

March 2017: Are diagnostic and/ or therapeutic Heart Caths considered prior cardiac cath? Same question for ECMO? **Include only cardiac cases (CPB or No CPB Cardiovascular cases only).**

August 2017: Is an Aortapexy considered a cardiothoracic surgery because it involves one of the great vessels? **Yes.**

SeqNo: 1100

Long Name: Number of Prior CPB Cardiothoracic Operations  
 Short Name: PrvOCtOpN  
 Database Table Name: Operations  
 Data Source: User  
 Format: Integer  
 Definition: Indicate how many cardiothoracic surgical procedures were performed on this patient, prior to this surgical procedure, utilizing CPB (do not include CPB support or ECMO support).

Low Value: 0 High Value: 50

Parent Long Name: Operation Type

Parent Short Name: OpType

Parent Value(s): = "CPB Cardiovascular", "No CPB Cardiovascular", "CPB Non-Cardiovascular", "ECMO", "Thoracic", "VAD Operation Done With CPB", "VAD Operation Done Without CPB" or "Other"

SeqNo: 1130

Long Name: Cross Clamp Time - No CPB

Short Name: XClampTmNC

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the total number of minutes the aorta is completely cross-clamped during this surgical procedure. Enter zero if no cross-clamp was used.

Low Value: 0 High Value: 600

Parent Long Name: Operation Type

Parent Short Name: OpType

Parent Value(s): = "No CPB Cardiovascular"

SeqNo: 1140

Long Name: CPB Blood Prime

Short Name: CPBPrimed

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether the CPB circuit was primed with blood other than the patient's own blood.

Parent Long Name: Operation Type

Parent Short Name: OpType

Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

SeqNo: 1150

Long Name: Cardiopulmonary Bypass Time

Short Name: CPBTm

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the total number of minutes that systemic return is diverted into the cardiopulmonary bypass (CPB) circuit and returned to the systemic system. This time



period (Cardiopulmonary Bypass Time) includes all periods of cerebral perfusion and sucker bypass. This time period (Cardiopulmonary Bypass Time) excludes any circulatory arrest and modified ultrafiltration periods. If more than one period of CPB is required during the surgical procedure, the sum of all the CPB periods will equal the total number of CPB minutes. Enter zero if cardiopulmonary bypass technique was not used.

Low Value: 0      High Value: 999

Parent Long Name: Operation Type

Parent Short Name: OpType

Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

SeqNo: 1160

Long Name: Cross Clamp Time - CPB

Short Name: XClampTm

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the total number of minutes that the coronary circulation is mechanically isolated from systemic circulation, either by an aortic cross clamp or systemic circulatory arrest. This time period (Cross Clamp Time) includes all intervals of intermittent or continuous cardioplegia administration. If more than one cross clamp period is required during this surgical procedure, the sum of the cross clamp periods is equal to the total number of cross clamp minutes. Enter zero if the coronary circulation was never mechanically isolated from systemic circulation, either by an aortic cross clamp or systemic circulatory arrest. For the following two operations: (1) "Transplant, Heart", and (2) "Transplant, Heart and lung", the field "Cross Clamp Time" will be defined as the cross clamp time of the donor heart. Therefore, these two operations represent the only operations where the field "Cross Clamp Time" can be greater than the field "Cardiopulmonary Bypass Time".

Low Value: 0      High Value: 600

Parent Long Name: Operation Type

Parent Short Name: OpType

Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

SeqNo: 1170

Long Name: Circulatory Arrest Time

Short Name: DHCATm

Database Table Name: Operations

Data Source: User

Format: Integer

**Definition:** Indicate the total number of minutes of complete cessation of blood flow to the patient. This time period (Circulatory Arrest Time) excludes any periods of cerebral perfusion. If more than one period of circulatory arrest is required during this surgical procedure, the sum of these periods is equal to the total duration of circulatory arrest. Enter zero if circulatory arrest technique was not used.

**Low Value:** 0      **High Value:** 200

**Parent Long Name:** Operation Type

**Parent Short Name:** OpType

**Parent Value(s):** = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

---

**SeqNo:** 1180

**Long Name:** Patient Temperature Monitoring Site - Bladder

**Short Name:** TempSiteBla

**Database Table Name:** Operations

**Data Source:** User

**Format:** Text (categorical values specified by STS)

**Definition:** Indicate whether the bladder monitoring site was utilized during this procedure to determine lowest and highest patient temperature during cardiopulmonary bypass.

**Parent Long Name:** Operation Type

**Parent Short Name:** OpType

**Parent Value(s):** = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

---

**SeqNo:** 1190

**Long Name:** Lowest Core Temperature - Bladder

**Short Name:** LowCTmpBla

**Database Table Name:** Operations

**Data Source:** User

**Format:** Real

**Definition:** Indicate the lowest temperature (Celsius) achieved during cardiopulmonary bypass as recorded using the bladder monitoring site.

**Low Value:** 1.0      **High Value:** 37.0

**Parent Long Name:** Patient Temperature Monitoring Site - Bladder

**Parent Short Name:** TempSiteBla

**Parent Value(s):** = "Yes"

---

**SeqNo:** 1200

---

Long Name: Patient Temperature Monitoring Site - Esophageal  
Short Name: TempSiteEso  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the esophageal monitoring site was utilized during this procedure to determine lowest and highest patient temperature during cardiopulmonary bypass.

Parent Long Name: Operation Type  
Parent Short Name: OpType  
Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

---

SeqNo: 1210

Long Name: Lowest Core Temperature - Esophageal  
Short Name: LowCTmpEso  
Database Table Name: Operations  
Data Source: User  
Format: Real  
Definition: Indicate the lowest temperature (Celsius) achieved during cardiopulmonary bypass as recorded using the esophageal monitoring site.

Low Value: 1.0      High Value: 37.0

Parent Long Name: Patient Temperature Monitoring Site - Esophageal  
Parent Short Name: TempSiteEso  
Parent Value(s): = "Yes"

---

SeqNo: 1220

Long Name: Patient Temperature Monitoring Site - Nasopharyngeal  
Short Name: TempSiteNas  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the nasopharyngeal monitoring site was utilized during this procedure to determine lowest and highest patient temperature during cardiopulmonary bypass.

Parent Long Name: Operation Type  
Parent Short Name: OpType  
Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

SeqNo: 1230

Long Name: Lowest Core Temperature - Nasopharyngeal  
Short Name: LowCTmpNas  
Database Table Name: Operations

---

Data Source: User  
Format: Real  
Definition: Indicate the lowest temperature (Celsius) achieved during cardiopulmonary bypass as recorded using the nasopharyngeal monitoring site.

Low Value: 1.0      High Value: 37.0

Parent Long Name: Patient Temperature Monitoring Site - Nasopharyngeal  
Parent Short Name: TempSiteNas  
Parent Value(s): = "Yes"

---

SeqNo: 1240

Long Name: Patient Temperature Monitoring Site - Rectal  
Short Name: TempSiteRec  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the rectal monitoring site was utilized during this procedure to determine lowest and highest patient temperature during cardiopulmonary bypass.  
Parent Long Name: Operation Type  
Parent Short Name: OpType  
Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

---

SeqNo: 1250

Long Name: Lowest Core Temperature - Rectal  
Short Name: LowCTmpRec  
Database Table Name: Operations  
Data Source: User  
Format: Real  
Definition: Indicate the lowest temperature (Celsius) achieved during cardiopulmonary bypass as recorded using the rectal monitoring site.

Low Value: 1.0      High Value: 37.0

Parent Long Name: Patient Temperature Monitoring Site - Rectal  
Parent Short Name: TempSiteRec  
Parent Value(s): = "Yes"

---

SeqNo: 1260

Long Name: Patient Temperature Monitoring Site - Tympanic  
Short Name: TempSiteTym  
Database Table Name: Operations  
Data Source: User

---

Format: Text (categorical values specified by STS)  
Definition: Indicate whether the tympanic monitoring site was utilized during this procedure to determine lowest and highest patient temperature during cardiopulmonary bypass.

Parent Long Name: Operation Type  
Parent Short Name: OpType  
Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

---

SeqNo: 1270  
Long Name: Lowest Core Temperature - Tympanic  
Short Name: LowCTmpTym  
Database Table Name: Operations  
Data Source: User  
Format: Real  
Definition: Indicate the lowest temperature (Celsius) achieved during cardiopulmonary bypass as recorded using the tympanic monitoring site.

Low Value: 1.0      High Value: 37.0

Parent Long Name: Patient Temperature Monitoring Site - Tympanic  
Parent Short Name: TempSiteTym  
Parent Value(s): = "Yes"

---

SeqNo: 1280  
Long Name: Patient Temperature Monitoring Site - Other  
Short Name: TempSiteOth  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether any other monitoring site was utilized during this procedure to determine lowest and highest patient temperature during cardiopulmonary bypass.

Parent Long Name: Operation Type  
Parent Short Name: OpType  
Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

---

SeqNo: 1290  
Long Name: Lowest Core Temperature - Other  
Short Name: LowCTmpOth  
Database Table Name: Operations  
Data Source: User  
Format: Real  
Definition: Indicate the lowest temperature (Celsius) achieved during cardiopulmonary bypass as

recorded using the other monitoring site.

Low Value: 1.0 High Value: 37.0

Parent Long Name: Patient Temperature Monitoring Site - Other  
 Parent Short Name: TempSiteOth  
 Parent Value(s): = "Yes"

SeqNo: 1300

Long Name: Cooling Time  
 Short Name: CoolTime  
 Database Table Name: Operations  
 Data Source: User  
 Format: Integer  
 Definition: Indicate the number of minutes of active cooling on cardiopulmonary bypass.

Low Value: 0 High Value: 200

Parent Long Name: Operation Type  
 Parent Short Name: OpType  
 Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

April: Is this question looking for the number of minutes it takes to cool the patient to the temperature, or should it include the minutes that they are held at the lowest temperature? **From the specs, cooling time is the number of minutes of active cooling on cardiopulmonary bypass which would include the time it takes to cool the patient as well as the time the patient is cooled.**

June 2017: After listening to the recording of the April Eddie call, I was confused while listening to the discussion of cooling time and how to determine the correct time of cooling and how "active" cooling time plays a part. From the explanation on the call and the way it was added in the data definitions, the cooling time should include the active cooling time in addition to the time it takes to maintain the desired low temp. This would clearly result in a much higher time than what our facility has been entering. We currently do not include the maintenance of the desired temp, but only the time it took to achieve the temp. If the maintenance time should be included, then the total time for cooling would be the majority of the bypass time. Please clarify the correct way and maybe the intent of this field for a better understanding. **From the specs, and the definition above, cooling time is the number of minutes of active cooling on cardiopulmonary bypass which would include the time it takes to cool the patient as well as the time the patient is cooled.**

SeqNo: 1310

Long Name: Rewarming Time  
 Short Name: RewarmTime  
 Database Table Name: Operations  
 Data Source: User  
 Format: Integer  
 Definition: Indicate the number of minutes from the initiation of rewarming until the target rewarming temperature is achieved.

Low Value: 0 High Value: 500

Parent Long Name: Operation Type

Parent Short Name: OpType

Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

---

July 2017: Question from our perfusionist: Trying to reconcile our CPB cooling and rewarming times. The help file mentions "target" temperature. The question I have is which temperature are we supposed to use for the "target" – Bladder, Naso, Esophageal, Rectal, Venous Blood? **Location of monitoring the temperature does not matter. The target temperature is referring to the temp you are trying to achieve or maintain.**

SeqNo: 1320

Long Name: Cerebral Perfusion Utilized

Short Name: CPerfUtil

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether cerebral perfusion was performed.

Parent Long Name: Operation Type

Parent Short Name: OpType

Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

---

SeqNo: 1330

Long Name: Cerebral Perfusion Time

Short Name: CPerfTime

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the total number of minutes cerebral perfusion was performed. This would include antegrade or retrograde cerebral perfusion strategies.

Low Value: 1 High Value: 999

Parent Long Name: Cerebral Perfusion Utilized

Parent Short Name: CPerfUtil

Parent Value(s): = "Yes"

---

SeqNo: 1340

Long Name: Cerebral Perfusion Cannulation Site - Innominate Artery

Short Name: CPerfCanInn

Database Table Name: Operations

Data Source: User

---

Format: Text (categorical values specified by STS)  
Definition: Indicate whether the innominate artery cannulation site was utilized for cerebral perfusion.

Parent Long Name: Cerebral Perfusion Utilized  
Parent Short Name: CPerfUtil  
Parent Value(s): = "Yes"

---

SeqNo: 1350  
Long Name: Cerebral Perfusion Cannulation Site - Right Subclavian  
Short Name: CPerfCanRSub  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the right subclavian cannulation site was utilized for cerebral perfusion.

Parent Long Name: Cerebral Perfusion Utilized  
Parent Short Name: CPerfUtil  
Parent Value(s): = "Yes"

---

SeqNo: 1360  
Long Name: Cerebral Perfusion Cannulation Site - Right Axillary Artery  
Short Name: CPerfCanRAx  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the right axillary artery cannulation site was utilized for cerebral perfusion.

Parent Long Name: Cerebral Perfusion Utilized  
Parent Short Name: CPerfUtil  
Parent Value(s): = "Yes"

---

SeqNo: 1370  
Long Name: Cerebral Perfusion Cannulation Site - Right Carotid Artery  
Short Name: CPerfCanRCar  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the right carotid artery cannulation site was utilized for cerebral perfusion.

Parent Long Name: Cerebral Perfusion Utilized  
Parent Short Name: CPerfUtil

---



Parent Value(s): = "Yes"

---

SeqNo: 1380

Long Name: Cerebral Perfusion Cannulation Site - Left Carotid Artery

Short Name: CPerfCanLCar

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether the left carotid artery cannulation site was utilized for cerebral perfusion.

Parent Long Name: Cerebral Perfusion Utilized

Parent Short Name: CPerfUtil

Parent Value(s): = "Yes"

---

SeqNo: 1390

Long Name: Cerebral Perfusion Cannulation Site - Superior Vena Cava

Short Name: CPerfCanSVC

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether the superior vena cava cannulation site was utilized for cerebral perfusion.

Parent Long Name: Cerebral Perfusion Utilized

Parent Short Name: CPerfUtil

Parent Value(s): = "Yes"

---

SeqNo: 1400

Long Name: Cerebral Perfusion Periods

Short Name: CPerfPer

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of periods of cerebral perfusion. For example, if the cerebral perfusion time is a total of 20 minutes and the patient received 4 separate 5 minute periods of cerebral perfusion, the cerebral perfusion periods would be 4.

Low Value: 1 High Value: 20

Parent Long Name: Cerebral Perfusion Utilized

Parent Short Name: CPerfUtil

Parent Value(s): = "Yes"

---

---

SeqNo: 1410  
Long Name: Cerebral Perfusion Flow Rate  
Short Name: CPerFlow  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the cerebral perfusion flow rate in milliliters per kilogram (mL/kg) per minute.

Low Value: 1 High Value: 999

Parent Long Name: Cerebral Perfusion Utilized  
Parent Short Name: CPerUtil  
Parent Value(s): = "Yes"

---

SeqNo: 1420  
Long Name: Cerebral Perfusion Temperature  
Short Name: CPerTemp  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the perfusate temperature (Celsius) maintained during cerebral perfusion.

Low Value: 1 High Value: 37

Parent Long Name: Cerebral Perfusion Utilized  
Parent Short Name: CPerUtil  
Parent Value(s): = "Yes"

---

SeqNo: 1430  
Long Name: Arterial Blood Gas Management During Cooling  
Short Name: ABldGasMgt  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the arterial blood gas management strategy utilized during the cooling phase of cardiopulmonary bypass prior to initiation of circulatory arrest or cerebral perfusion.

Parent Long Name: Operation Type  
Parent Short Name: OpType

---

Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

---

SeqNo: 1440

Long Name: Hematocrit Prior to Circulatory Arrest or Cerebral Perfusion

Short Name: HCTPriCircA

Database Table Name: Operations

Data Source: User

Format: Real

Definition: Indicate the last hematocrit value prior to initiation of circulatory arrest or cerebral perfusion.

Low Value: 5.0      High Value: 70.0

Parent Long Name: Operation Type

Parent Short Name: OpType

Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

---

SeqNo: 1450

Long Name: Cardioplegia Delivery

Short Name: CplegiaDeliv

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate the delivery method of cardioplegia if used.

Parent Long Name: Operation Type

Parent Short Name: OpType

Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

---

12/2015: If a patient is given more than one kind of cardioplegia solution (e.g. Buckberg and del Nido) in the same operative event, which do I capture? In one case, one was arresting, the rest were maintenance; but in another, they were both arresting. **Indicate the first one.**

---

SeqNo: 1460

Long Name: Cardioplegia Type

Short Name: CplegiaType

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate the type of cardioplegia used.

Parent Long Name: Cardioplegia Delivery

Parent Short Name: CplegiaDeliv

Parent Value(s): = "Antegrade", "Retrograde" or "Both"

---

SeqNo: 1470

Long Name: Cardioplegia Solution

Short Name: CplegiaSolution

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate the cardioplegia solution used during this procedure.

Parent Long Name: Cardioplegia Delivery

Parent Short Name: CplegiaDeliv

Parent Value(s): = "Antegrade", "Retrograde" or "Both"

SeqNo: 1490

Long Name: Cardioplegia Number Of Doses

Short Name: CplegiaDose

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of doses of cardioplegia administered

Low Value: 1      High Value: 50

Parent Long Name: Cardioplegia Delivery

Parent Short Name: CplegiaDeliv

Parent Value(s): = "Antegrade", "Retrograde" or "Both"

SeqNo: 1640

Long Name: Hematocrit - First after initiating CPB

Short Name: HCTFirst

Database Table Name: Operations

Data Source: User

Format: Real

Definition: Indicate the first hematocrit measured after initiating CPB.

Low Value: 5.0      High Value: 70.0

Parent Long Name: Operation Type

Parent Short Name: OpType

Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

**July 2016: Which HCT values (first, last, and first off) do you record if a patient has multiple bypass runs? If multiple bypass runs occur in the course of a single OR procedure, list the first Hct value during the first**

**bypass run. This may not be the most meaningful value in the world, but coding will be consistent with all other cases.**

January 2017: As a rule, my organization and surgical team do not chart HCT, but rather HGB during their cases. Thus, I have no charted HCT for the preoperative or operative periods. Can I do a calculated HCT, or bypass this entire field. **Leave it blank.** If I bypass the whole series, will our data be rejected? **No**

SeqNo: 1650

Long Name: Hematocrit - Last Measured During CPB  
 Short Name: HCTLast  
 Database Table Name: Operations  
 Data Source: User  
 Format: Real  
 Definition: Indicate the last hematocrit measured during CPB.

Low Value: 5.0 High Value: 70.0

Parent Long Name: Operation Type  
 Parent Short Name: OpType  
 Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

SeqNo: 1660

Long Name: Hematocrit - Post-CPB and Post-Protamine  
 Short Name: HCTPost  
 Database Table Name: Operations  
 Data Source: User  
 Format: Real  
 Definition: Indicate the hematocrit measured post-CPB following protamine administration.

Low Value: 5.0 High Value: 70.0

Parent Long Name: Operation Type  
 Parent Short Name: OpType  
 Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

SeqNo: 1670

Long Name: Ultrafiltration Performed After CPB  
 Short Name: Ultrafiltration  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether ultrafiltration was performed after CPB.

Parent Long Name: Operation Type

Parent Short Name: OpType  
 Parent Value(s): = "CPB Cardiovascular", "CPB Non-Cardiovascular" or "VAD Operation Done With CPB"

**November 2016:** Is this sequence meant to capture only ultrafiltration done after CPB? Our understanding based on information from the perfusionist is that CUF is performed on CPB while MUF is performed post CPB. If this is the case, CUF wouldn't ever be selected, so should it be a choice? Or is the intent to capture all ultrafiltration done during the operation? **It is appropriate to be confused about this field: The wording "Indicate whether ultrafiltration was performed after CPB," is confusing.**

**As a temporary work-around, until such language can be changed in the specifications, I would suggest interpreting this field as pertaining to "ultrafiltration after the initiation of cardiopulmonary bypass,"** From a practical standpoint, this then includes the period of cpb support and the period immediately following cessation of cpb support. Conventional ultrafiltration, if used, is a process that takes place during the period of cpb support. Modified ultrafiltration (MUF), if used, is a process that is carried out after cessation of cardiopulmonary bypass support, before removal of arterial and venous cannulae from the patient. I think an additional question was raised earlier, and in terms of coding this field in the CHSD, processes to promote hemoconcentration other than MUF, carried out after cessation of cardiopulmonary bypass support, should not be coded as either conventional ultrafiltration (CUF) or modified ultrafiltration (MUF). While this may encompass use of a "cell-saver" or other centrifugation device, it may also involve the hemoconcentrating filter that is suitable for CUF or MUF. So, for example, use of the same hemofiltration filter for the purpose of concentrating the volume of blood-containing fluid left in the bypass circuit and the end of the case should not be coded as ultrafiltration, unless it involves removal of fluid directly from, and reinfusion of blood directly to the patient via the cpb circuit and cannulae, as is done in both forms of ultrafiltration.

SeqNo: 1770  
 Long Name: Pulmonary Vascular Resistance Measured  
 Short Name: PVRMeas  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether the Pulmonary Vascular Resistance (PVR) in Woods units was measured by cardiac catheterization prior to this operation.

Parent Long Name: Operation Type  
 Parent Short Name: OpType  
 Parent Value(s): = "CPB Cardiovascular", "No CPB Cardiovascular", "CPB Non-Cardiovascular", "ECMO", "Thoracic", "VAD Operation Done With CPB", "VAD Operation Done Without CPB" or "Other"

**April 2016:** This item says "prior to this operation". Does this mean during this hospitalization? **Yes.**  
 What is the time frame to report a PVR prior to surgery? **6 Months.**

SeqNo: 1780  
 Long Name: Pulmonary Vascular Resistance  
 Short Name: PVR

Database Table Name: Operations  
 Data Source: User  
 Format: Real  
 Definition: If the patient's weight is greater than or equal to 40 kilograms, indicate the pulmonary vascular resistance (in Wood units) as measured by cardiac catheterization.

Low Value: 0.0      High Value: 100.0

Parent Long Name:  
 Parent Short Name: PVRMeas|WeightKg  
 Parent Value(s): = "Yes"|>=40

---

SeqNo: 1790  
 Long Name: Pulmonary Vascular Resistance Index  
 Short Name: PVRI  
 Database Table Name: Operations  
 Data Source: User  
 Format: Real  
 Definition: If the patient's weight is less than 40 kilograms, indicate the Pulmonary Vascular Resistance Index (in Wood units x m2) as measured by cardiac catheterization.

Low Value: 0.0      High Value: 100.0

Parent Long Name:  
 Parent Short Name: PVRMeas|WeightKg  
 Parent Value(s): = "Yes"|<40

---

SeqNo: 2461  
 Long Name: Autologous Transfusion  
 Short Name: AutologousTrans  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether the patient was transfused with any autologous blood products that had been collected prior to surgery (e.g. self-donated).

SeqNo: 2462  
 Long Name: Cell Saver/Salvage Blood Returned to Patient  
 Short Name: CellSavSal  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether cell saver/salvage blood was infused during this procedure.

---

SeqNo: 2743

Long Name: Transfusion of Non-Autologous Blood Products During or After Procedure  
 Short Name: TransfusBldProdAny  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether the patient received non-autologous (self-donated) blood products during or after this procedure.

July 2016: For patients that have 2 major operations during the same admission, on the second operation, are we supposed to answer this question "No" even though product was used for pump prime and intraoperatively, because all products are supposed to be tied to the index operation?

**DCRI has requested that all blood product usage be tied back to the index operation for a given admission (this will be a point of discussion at the upcoming AQO). So, for example, if a patient has Norwood operation and remains in the hospital for several months and then has a Glenn, all the blood products would be listed in the original Norwood as transfusion after 24 hours. I know this doesn't make a lot of clinical sense, which is why we are discussing this (times).and intubation/reintubation.**

December 2016: Where should pooled platelets be entered? **Document it in Seq. 2763.**

SeqNo: 2744

Long Name: Transfusion of Blood Products During Procedure  
 Short Name: Transfusion  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether the patient received a transfusion during this procedure.

Parent Long Name: Transfusion of Non-Autologous Blood Products During or After  
 Parent Short Name: TransfusBldProdAny  
 Parent Value(s): = "Yes"

SeqNo: 2751

Long Name: Blood Products Transfused - Packed Red Blood Cells (PRBC) - Units Transfused During Procedure  
 Short Name: BldProdPRBCDur  
 Database Table Name: Operations  
 Data Source: User  
 Format: Integer  
 Definition: Indicate the number of units of Packed Red Blood Cells (PRBC) the patient received during the procedure (including CPB PRIME). Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.



Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0      High Value: 100

Parent Long Name: Transfusion of Blood Products During Procedure  
 Parent Short Name: Transfusion  
 Parent Value(s): = "Yes"

SeqNo: 2754

Long Name: Blood Products Transfused - Fresh Frozen Plasma (FFP) - Units Transfused During Procedure

Short Name: BldProdFFPDur

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Fresh Frozen Plasma (FFP) the patient received during the procedure (including CPB PRIME). Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0      High Value: 100

Parent Long Name: Transfusion of Blood Products During Procedure  
 Parent Short Name: Transfusion  
 Parent Value(s): = "Yes"

SeqNo: 2757

Long Name: Blood Products Transfused - Fresh Plasma - Units Transfused During Procedure

Short Name: BldProdFreshPDur

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Fresh Plasma (<72 Hours Post-collection, never frozen) the patient received during the procedure (including CPB PRIME). Unit(s) are defined

as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0

High Value: 100

Parent Long Name: Transfusion of Blood Products During Procedure

Parent Short Name: Transfusion

Parent Value(s): = "Yes"

SeqNo: 2760

Long Name: Blood Products Transfused - Single Donor Platelet Pheresis - Units Transfused During Procedure

Short Name: BldProdSnglPlatDur

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Single Donor Platelet Pheresis (1 pheresis UNIT = approx 6 indiv units) the patient received during the procedure (including CPB PRIME). Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0

High Value: 100

Parent Long Name: Transfusion of Blood Products During Procedure

Parent Short Name: Transfusion

Parent Value(s): = "Yes"

July 2016: The definition says "Indicate the number of units of Single Donor Platelet Pheresis (1 pheresis UNIT = approx 6 indiv units) the patient received during the procedure (including CPB PRIME). Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation." We use pooled platelet pheresis and not single donor pheresis. We have no way of telling how many donors were used in each unit, so how do I indicate these? **Check with your blood bank; they should be able to give you a better answer than I can. How are your platelet concentrates ordered? By volume? When I last used**

**platelet pooled products (it has been some time) we ordered a specific number of pooled platelets (ie 4 units pooled). The pooling is mostly for convenience of administration (rather than having to draw the platelets from 4 separate bags they are combined into one by the blood bank).**

SeqNo: 2763

Long Name: Blood Products Transfused - Individual Platelets - Units Transfused During Procedure

Short Name: BldProdIndPlatDur

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Individual Platelets, including concentrated, the patient received during the procedure (including CPB PRIME). Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0 High Value: 100

Parent Long Name: Transfusion of Blood Products During Procedure

Parent Short Name: Transfusion

Parent Value(s): = "Yes"

SeqNo: 2766

Long Name: Blood Products Transfused - Cryoprecipitate - Units Transfused During Procedure

Short Name: BldProdCryoDur

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Cryoprecipitate the patient received during the procedure (including CPB PRIME). Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0 High Value: 100

Parent Long Name: Transfusion of Blood Products During Procedure  
Parent Short Name: Transfusion  
Parent Value(s): = "Yes"

---

SeqNo: 2769

Long Name: Blood Products Transfused - Fresh Whole Blood - Units Transfused During Procedure

Short Name: BldProdFreshWBDur

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Fresh Whole Blood (< 72 Hours post-collection) the patient received during the procedure (including CPB PRIME). Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0 High Value: 100

Parent Long Name: Transfusion of Blood Products During Procedure  
Parent Short Name: Transfusion  
Parent Value(s): = "Yes"

---

SeqNo: 2772

Long Name: Blood Products Transfused - Whole Blood - Units Transfused During Procedure

Short Name: BldProdWBDur

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Whole Blood (> 72 hours post-collection) the patient received during the procedure (including CPB PRIME). Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single

unit.

Low Value: 0      High Value: 100

Parent Long Name: Transfusion of Blood Products During Procedure  
 Parent Short Name: Transfusion  
 Parent Value(s): = "Yes"

---

SeqNo: 2779

Long Name: Transfusion of Blood Products Within 24 Hours Post-Procedure

Short Name: TransfusBldProdLT24

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether the patient received blood products within 24 hours post-procedure.

Parent Long Name: Transfusion of Non-Autologous Blood Products During or After  
 Parent Short Name: TransfusBldProdAny  
 Parent Value(s): = "Yes"

---

SeqNo: 2780

Long Name: Blood Products Transfused - Packed Red Blood Cells (PRBC) - Units Transfused Within 24 Hours Post-Procedure

Short Name: BldProdPRBCLT24

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Packed Red Blood Cells (PRBC) the patient received within 24 hours post-procedure. Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0      High Value: 100

Parent Long Name: Transfusion of Blood Products Within 24 Hours Post-Procedure  
 Parent Short Name: TransfusBldProdLT24  
 Parent Value(s): = "Yes"

---

SeqNo: 2781

Long Name: Blood Products Transfused - Fresh Frozen Plasma (FFP) - Units Transfused Within 24 Hours Post-Procedure

Short Name: BldProdFFPLT24

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Fresh Frozen Plasma (FFP) the patient received within 24 hours post-procedure. Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0 High Value: 100

Parent Long Name: Transfusion of Blood Products Within 24 Hours Post-Procedure

Parent Short Name: TransfusBldProdLT24

Parent Value(s): = "Yes"

SeqNo: 2782

Long Name: Blood Products Transfused - Fresh Plasma - Units Transfused Within 24 Hours Post-Procedure

Short Name: BldProdFreshPLT24

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Fresh Plasma (<72 Hours Post-collection, never frozen) the patient received within 24 hours post-procedure. Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0 High Value: 100

Parent Long Name: Transfusion of Blood Products Within 24 Hours Post-Procedure  
Parent Short Name: TransfusBldProdLT24  
Parent Value(s): = "Yes"

---

SeqNo: 2783

Long Name: Blood Products Transfused - Single Donor Platelet Pheresis - Units Transfused Within 24 Hours Post-Procedure

Short Name: BldProdSnglPlatLT24

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Single Donor Platelet Pheresis (1 pheresis UNIT = approx 6 indiv units) the patient received within 24 hours post-procedure. Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0 High Value: 100

Parent Long Name: Transfusion of Blood Products Within 24 Hours Post-Procedure  
Parent Short Name: TransfusBldProdLT24  
Parent Value(s): = "Yes"

---

SeqNo: 2784

Long Name: Blood Products Transfused - Individual Platelets - Units Transfused Within 24 Hours Post-Procedure

Short Name: BldProdIndPlatLT24

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Individual Platelets, including concentrated, the patient received within 24 hours post-procedure. Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

---

Low Value: 0      High Value: 100

Parent Long Name: Transfusion of Blood Products Within 24 Hours Post-Procedure  
 Parent Short Name: TransfusBldProdLT24  
 Parent Value(s): = "Yes"

SeqNo: 2785

Long Name: Blood Products Transfused - Cryoprecipitate - Units Transfused Within 24 Hours Post-Procedure

Short Name: BldProdCryoLT24

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Cryoprecipitate the patient received within 24 hours post-procedure. Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0      High Value: 100

Parent Long Name: Transfusion of Blood Products Within 24 Hours Post-Procedure  
 Parent Short Name: TransfusBldProdLT24  
 Parent Value(s): = "Yes"

SeqNo: 2786

Long Name: Blood Products Transfused - Fresh Whole Blood - Units Transfused Within 24 Hours Post-Procedure

Short Name: BldProdFreshWBLT24

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Fresh Whole Blood (< 72 Hours post-collection) the patient received within 24 hours post-procedure. Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and



finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0      High Value: 100

Parent Long Name: Transfusion of Blood Products Within 24 Hours Post-Procedure  
 Parent Short Name: TransfusBldProdLT24  
 Parent Value(s): = "Yes"

SeqNo: 2787

Long Name: Blood Products Transfused - Whole Blood - Units Transfused Within 24 Hours Post-Procedure

Short Name: BldProdWBLT24

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Whole Blood (> 72 hours post-collection) the patient received within 24 hours post-procedure. Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0      High Value: 100

Parent Long Name: Transfusion of Blood Products Within 24 Hours Post-Procedure  
 Parent Short Name: TransfusBldProdLT24  
 Parent Value(s): = "Yes"

SeqNo: 2788

Long Name: Transfusion of Blood Products After 24 Hours Post-Procedure

Short Name: TransfusBldProdGT24

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether the patient received blood products after 24 hours post-procedure.

Parent Long Name: Transfusion of Non-Autologous Blood Products During or After  
 Parent Short Name: TransfusBldProdAny

Parent Value(s): = "Yes"

---

SeqNo: 2789

Long Name: Blood Products Transfused - Packed Red Blood Cells (PRBC) - Units Transfused After 24 Hours Post-Procedure

Short Name: BldProdPRBCGT24

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Packed Red Blood Cells (PRBC) the patient received after 24 hours post-procedure. Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0

High Value: 100

Parent Long Name: Transfusion of Blood Products After 24 Hours Post-Procedure

Parent Short Name: TransfusBldProdGT24

Parent Value(s): = "Yes"

---

SeqNo: 2790

Long Name: Blood Products Transfused - Fresh Frozen Plasma (FFP) - Units Transfused After 24 Hours Post-Procedure

Short Name: BldProdFFPGT24

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Fresh Frozen Plasma (FFP) the patient received after 24 hours post-procedure. Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0

High Value: 100

---

Parent Long Name: Transfusion of Blood Products After 24 Hours Post-Procedure  
Parent Short Name: TransfusBldProdGT24  
Parent Value(s): = "Yes"

---

SeqNo: 2791

Long Name: Blood Products Transfused - Fresh Plasma - Units Transfused After 24 Hours Post-Procedure

Short Name: BldProdFreshPGT24

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Fresh Plasma (<72 Hours Post-collection, never frozen) the patient received after 24 hours post-procedure. Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0 High Value: 100

Parent Long Name: Transfusion of Blood Products After 24 Hours Post-Procedure  
Parent Short Name: TransfusBldProdGT24  
Parent Value(s): = "Yes"

---

SeqNo: 2792

Long Name: Blood Products Transfused - Single Donor Platelet Pheresis - Units Transfused After 24 Hours Post-Procedure

Short Name: BldProdSnglPlatGT24

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Single Donor Platelet Pheresis (1 pheresis UNIT = approx 6 indiv units) the patient received after 24 hours post-procedure. Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

---

Low Value: 0 High Value: 100

Parent Long Name: Transfusion of Blood Products After 24 Hours Post-Procedure  
Parent Short Name: TransfusBldProdGT24  
Parent Value(s): = "Yes"

---

SeqNo: 2793

Long Name: Blood Products Transfused - Individual Platelets - Units Transfused After 24 Hours Post-Procedure

Short Name: BldProdIndPlatGT24

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Individual Platelets, including concentrated, the patient received after 24 hours post-procedure. Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0 High Value: 100

Parent Long Name: Transfusion of Blood Products After 24 Hours Post-Procedure  
Parent Short Name: TransfusBldProdGT24  
Parent Value(s): = "Yes"

---

SeqNo: 2794

Long Name: Blood Products Transfused - Cryoprecipitate - Units Transfused After 24 Hours Post-Procedure

Short Name: BldProdCryoGT24

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Cryoprecipitate the patient received after 24 hours post-procedure. Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and

finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0      High Value: 100  
Parent Long Name: Transfusion of Blood Products After 24 Hours Post-Procedure  
Parent Short Name: TransfusBldProdGT24  
Parent Value(s): = "Yes"

---

SeqNo: 2795

Long Name: Blood Products Transfused - Fresh Whole Blood - Units Transfused After 24 Hours Post-Procedure

Short Name: BldProdFreshWBGT24

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Fresh Whole Blood (< 72 Hours post-collection) the patient received after 24 hours post-procedure. Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit. Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0      High Value: 100

Parent Long Name: Transfusion of Blood Products After 24 Hours Post-Procedure  
Parent Short Name: TransfusBldProdGT24  
Parent Value(s): = "Yes"

---

SeqNo: 2796

Long Name: Blood Products Transfused - Whole Blood - Units Transfused After 24 Hours Post-Procedure

Short Name: BldProdWBGT24

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the number of units of Whole Blood (> 72 hours post-collection) the patient received after 24 hours post-procedure. Unit(s) are defined as aliquots of blood originating from a single donor, regardless of volume transfused from that donation. For example, two 50 ml aliquots of PRBCs from the same donor count as ONE (1) unit.

Two aliquots from DIFFERENT donors count as TWO (2) units.

Units that are split over multiple time periods, such as starting a unit in the OR and finishing it in the ICU should be assigned to the period where the transfusion was begun or where the majority of the unit is given. The intent is not to double count a single unit.

Low Value: 0 High Value: 100

Parent Long Name: Transfusion of Blood Products After 24 Hours Post-Procedure  
Parent Short Name: TransfusBldProdGT24  
Parent Value(s): = "Yes"

---

SeqNo: 2797  
Long Name: Directed Donor Units  
Short Name: DirDonorUnits  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient received any directed donor transfusions during this procedure.

Parent Long Name: Transfusion of Non-Autologous Blood Products During or After  
Parent Short Name: TransfusBldProdAny  
Parent Value(s): = "Yes"

---

SeqNo: 2798  
Long Name: Antifibrinolytic Used Intraoperatively  
Short Name: AntifibUsage  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether antifibrinolytics were used intraoperatively.

---

SeqNo: 2799  
Long Name: Epsilon Amino-Caproic Acid (Amicar,EACA) Used  
Short Name: AntifibEpUse  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether EACA was used.

---

Parent Long Name: Antifibrinolytic Used Intraoperatively  
 Parent Short Name: AntifibUsage  
 Parent Value(s): = "Yes"

---

SeqNo: 2800

Long Name: Epsilon Amino-Caproic Acid (Amicar,EACA) Load mg/kg

Short Name: AntifibEpLoad

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the loading dose in mg/kg of epsilon aminocaproic acid (Amicar) given during this procedure. Enter zero if no loading dose given.

Low Value: 0      High Value: 300

Parent Long Name: Epsilon Amino-Caproic Acid (Amicar,EACA) Used

Parent Short Name: AntifibEpUse

Parent Value(s): = "Yes"

---

July 2016: I'm confused with the Antifibrinolytic section. I have a new surgeon who has Anesthesia give 100mg/kg bolus as they go on pump and as they come off. The perfusionist also gives one bolus dose of 100mg/kg. They do not give an infusion. So do I count the 1st bolus by Anesthesia as the answer to sequence AntifibEpLoad(2800), then the Perfusion dose in sequence AntifibEpPrime (2801) And then the infusion would 0 for sequence AntifibEpInfRate (2803). And my second bolus from Anesthesia as they go off pump doesn't go anywhere? Is the goal of this section to capture the amount of Amicar used or just know the mg/kg dosed. Also, please give more guidance on sequence AntifibEpPrimeDose (2802). If there is a value populated in seq 2802, wouldn't seq 2801 always be yes since it's in the prime for the pump? For example, my perfusionists add 100mg/kg to the pump prime but I would assume to do that they would have to figure out mg/ml in order to add it to the bag. **This is sort of an unusual dosing schema and doesn't neatly fit into our algorithm honestly. Most institutions I am familiar with utilize a combination of bolus and then infusion with an additional bolus in the pump prime.**

---

SeqNo: 2801

Long Name: Epsilon Amino-Caproic Acid (Amicar,EACA) Pump Prime mg/kg

Short Name: AntifibEpPrime

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the pump priming dose in mg/kg of epsilon aminocaproic acid (Amicar) given during this procedure. Enter zero if no pump priming dose given.

Low Value: 0      High Value: 300

Parent Long Name: Epsilon Amino-Caproic Acid (Amicar,EACA) Used

---

Parent Short Name: AntifibEpUse  
Parent Value(s): = "Yes"

---

SeqNo: 2802  
Long Name: EACA Dosed As mg per ml of Pump Prime  
Short Name: AntifibEpPrimeDose  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the Epsilon Amino-Caproic Acid was dosed as mg per ml of Pump Prime.

Parent Long Name: Epsilon Amino-Caproic Acid (Amicar,EACA) Pump Prime mg/kg  
Parent Short Name: AntifibEpPrime  
Parent Value(s): >0

---

SeqNo: 2803  
Long Name: Epsilon Amino-Caproic Acid (Amicar,EACA) Infusion Rate mg/kg/hr  
Short Name: AntifibEpInfRate  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the infusion rate in mg/kg/hour of epsilon aminocaproic acid (Amicar) given during this procedure. Enter zero if no infusion initiated.

Low Value: 0      High Value: 200

Parent Long Name: Epsilon Amino-Caproic Acid (Amicar,EACA) Used  
Parent Short Name: AntifibEpUse  
Parent Value(s): = "Yes"

---

SeqNo: 2804  
Long Name: Tranexamic Acid Used  
Short Name: AntifibTranexUse  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether tranexamic acid was used during this procedure.

Parent Long Name: Antifibrinolytic Used Intraoperatively  
Parent Short Name: AntifibUsage  
Parent Value(s): = "Yes"

September 2016: Our Tranexamic Acid documentation is done as "1000mg" IV or "500mg" IV. I don't see a



loading dose or pump prime or it given as mg/kg. How do I document the doses given? **Document one dose in the loading dose field and document one dose in the pump prime field. Leave the infusion rate field blank or document "0".**

---

October 2016: We document Tranexamic use in two ways: pump prime (covered by Sequence #2806) and then as bolus doses for every 6 hours of surgery. How do I code the bolus doses? And is Antifibrinolytic use a required field affecting star rating and M&M? **Document the initial bolus as the loading dose and then the rate would be "0". Antifibrinolytic use does not affect the star rating.**

---

SeqNo: 2805

Long Name: Tranexamic Acid Load mg/kg

Short Name: AntifibTranexLoad

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the loading dose in mg/kg of tranexamic acid given during this procedure. Enter zero if no loading dose given.

Low Value: 0      High Value: 150

Parent Long Name: Tranexamic Acid Used

Parent Short Name: AntifibTranexUse

Parent Value(s): = "Yes"

---

SeqNo: 2806

Long Name: Tranexamic Acid Pump Prime mg/kg

Short Name: AntifibTranexPrime

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the pump priming dose in mg/kg of tranexamic acid given during this procedure. Enter zero if no pump priming dose given.

Low Value: 0      High Value: 150

Parent Long Name: Tranexamic Acid Used

Parent Short Name: AntifibTranexUse

Parent Value(s): = "Yes"

---

SeqNo: 2807

Long Name: Tranexamic Dosed As mg per ml of Pump Prime

Short Name: AntifibTranexPrimeDose

Database Table Name: Operations

---

Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether the Tranexamic was dosed as mg per ml of Pump Prime.

Parent Long Name: Tranexamic Acid Pump Prime mg/kg  
 Parent Short Name: AntifibTranexPrime  
 Parent Value(s): >0

SeqNo: 2809

Long Name: Tranexamic Acid Infusion Rate mg/kg/hr  
 Short Name: AntifibTranexInfRate  
 Database Table Name: Operations  
 Data Source: User  
 Format: Integer  
 Definition: Indicate the infusion rate in mg/kg/hour of tranexamic acid given during this procedure. Enter zero if no infusion initiated.

Low Value: 0      High Value: 50

Parent Long Name: Tranexamic Acid Used  
 Parent Short Name: AntifibTranexUse  
 Parent Value(s): = "Yes"

May 2016: The question is asking for rate of TXA infusion in mg/kg/hr. It will not let me enter a decimal point, it is only accepting whole numbers. For example, the anesthesia record documents an infusion rate of 3.1mg/kg/hr for a total of 477 minutes. The field will not accept 3.1 it will only accept 3. Rounding on medication dosages to the nearest whole number is extremely inaccurate. Now, I cannot tell if this issue lies within my software, Axis Pats, or with the new version. **I confirmed with DCRI and that field is an integer field, so the software is working as it should. This field was an integer even in the previous data version 3.22.**

SeqNo: 2810

Long Name: Trasylol (Aprotinin) Used  
 Short Name: AntifibTrasylUse  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether trasylol (aprotinin) was given to the patient during this procedure.

Parent Long Name: Antifibrinolytic Used Intraoperatively  
 Parent Short Name: AntifibUsage  
 Parent Value(s): = "Yes"

SeqNo: 2811

Long Name: Trasylo1 (Aprotinin) Load cc/kg  
Short Name: AntifibTrasy1Load  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the loading dose of trasylo1 (aprotinin) in cc/kg used during this procedure.  
Enter zero if no loading dose was used.

Low Value: 0 High Value: 10

Parent Long Name: Trasylo1 (Aprotinin) Used  
Parent Short Name: AntifibTrasy1Use  
Parent Value(s): = "Yes"

---

SeqNo: 2812

Long Name: Trasylo1 (Aprotinin) Pump Prime cc/kg  
Short Name: AntifibTrasy1Prime  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the pump priming dose of trasylo1 (aprotinin) in cc/kg used during this procedure. Enter zero if no pump priming dose was used.

Low Value: 0 High Value: 10

Parent Long Name: Trasylo1 (Aprotinin) Used  
Parent Short Name: AntifibTrasy1Use  
Parent Value(s): = "Yes"

---

SeqNo: 2813

Long Name: Trasylo1 (Aprotinin) Infusion Rate cc/kg/hr  
Short Name: AntifibTrasy1InfRate  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the infusion rate of trasylo1 (aprotinin) in cc/kg/hour used during this procedure. Enter zero if no infusion initiated.

Low Value: 0 High Value: 10

Parent Long Name: Trasylo1 (Aprotinin) Used  
Parent Short Name: AntifibTrasy1Use  
Parent Value(s): = "Yes"

---

---

SeqNo: 2815  
 Long Name: Procoagulant Used Intraoperatively  
 Short Name: ProcoagUsage  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether procoagulents were used intraoperatively

---

SeqNo: 2816  
 Long Name: Factor VIIa (Novoseven) Usage  
 Short Name: ProcoagFactorVIIa  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether Factor VIIa (Novoseven) was administered intraoperatively.

Parent Long Name: Procoagulant Used Intraoperatively  
 Parent Short Name: ProcoagUsage  
 Parent Value(s): = "Yes"

---

SeqNo: 2817  
 Long Name: Factor VIIa (Novoseven) mcg/kg - Dose 1  
 Short Name: ProcoagFactorVIIa1  
 Database Table Name: Operations  
 Data Source: User  
 Format: Integer  
 Definition: Indicate the first dose in micrograms per kilogram of Factor VIIa given during this procedure.

Low Value: 1      High Value: 200

Parent Long Name: Factor VIIa (Novoseven) Usage  
 Parent Short Name: ProcoagFactorVIIa  
 Parent Value(s): = "Yes"

---

SeqNo: 2818  
 Long Name: Factor VIIa (Novoseven) mcg/kg - Dose 2  
 Short Name: ProcoagFactorVIIa2  
 Database Table Name: Operations  
 Data Source: User

---

Format: Integer  
Definition: Indicate the second dose in micrograms per kilogram of Factor VIIa given during this procedure. Enter zero if no second dose given.

Low Value: 0 High Value: 200

Parent Long Name: Factor VIIa (Novoseven) Usage  
Parent Short Name: ProcoagFactorVIIa  
Parent Value(s): = "Yes"

---

SeqNo: 2819

Long Name: Factor VIIa (Novoseven) mcg/kg - Dose 3

Short Name: ProcoagFactorVIIa3

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the third dose in micrograms per kilogram of Factor VIIa given during this procedure. Enter zero if no third dose given.

Low Value: 0 High Value: 200

Parent Long Name: Factor VIIa (Novoseven) mcg/kg - Dose 2  
Parent Short Name: ProcoagFactorVIIa2  
Parent Value(s): >0

---

SeqNo: 2823

Long Name: Prothrombin Complex Concentrate - 4 (PCC-4, KCentra) Usage

Short Name: ProCmplxCon4

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether Prothrombin Complex Concentrate - 4 (PCC-4, KCentra) was administered intraoperatively.

Parent Long Name: Procoagulant Used Intraoperatively  
Parent Short Name: ProcoagUsage  
Parent Value(s): = "Yes"

---

SeqNo: 2824

Long Name: Prothrombin Complex Concentrate - 4 (PCC-4, KCentra) units/kg - Dose 1

Short Name: ProCmplxCon4Ds1

Database Table Name: Operations

Data Source: User

---

Format: Integer  
Definition: Indicate the first dose in units per kilogram of Prothrombin Complex Concentrate - 4 (PCC-4, KCentra).

Low Value: 1 High Value: 100

Parent Long Name: Prothrombin Complex Concentrate - 4 (PCC-4, KCentra) Usage  
Parent Short Name: ProCmplxCon4  
Parent Value(s): = "Yes"

---

SeqNo: 2825

Long Name: Prothrombin Complex Concentrate - 4 (PCC-4, KCentra) units/kg - Dose 2  
Short Name: ProCmplxCon4Ds2  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the second dose in units per kilogram of Prothrombin Complex Concentrate - 4 (PCC-4, KCentra). Enter zero if no second dose given.

Low Value: 0 High Value: 100

Parent Long Name: Prothrombin Complex Concentrate - 4 (PCC-4, KCentra) Usage  
Parent Short Name: ProCmplxCon4  
Parent Value(s): = "Yes"

---

SeqNo: 2826

Long Name: Prothrombin Complex Concentrate - 4 (PCC-4, KCentra) units/kg - Dose 3  
Short Name: ProCmplxCon4Ds3  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the third dose in units per kilogram of Prothrombin Complex Concentrate - 4 (PCC-4, KCentra). Enter zero if no third dose given.

Low Value: 0 High Value: 100

Parent Long Name: Prothrombin Complex Concentrate - 4 (PCC-4, KCentra) units/kg - Dose 2  
Parent Short Name: ProCmplxCon4Ds2  
Parent Value(s): >0

---

SeqNo: 2827

Long Name: Prothrombin Complex Concentrate - 4 With Factor VIIa (FEIBA) Usage  
Short Name: ProCmplxCon4W7a  
Database Table Name: Operations

---

Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Prothrombin Complex Concentrate - 4 With Factor VIIa (FEIBA) was administered intraoperatively.

Parent Long Name: Procoagulent Used Intraoperatively  
Parent Short Name: ProcoagUsage  
Parent Value(s): = "Yes"

---

SeqNo: 2828  
Long Name: Prothrombin Complex Concentrate - 4 With Factor VIIa (FEIBA) units/kg - Dose 1  
Short Name: ProCmplxCon4W7a1  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the first dose in units per kilogram of Prothrombin Complex Concentrate - 4 With Factor VIIa (FEIBA).

Low Value: 1      High Value: 200

Parent Long Name: Prothrombin Complex Concentrate - 4 With Factor VIIa (FEIBA) Usage  
Parent Short Name: ProCmplxCon4W7a  
Parent Value(s): = "Yes"

---

SeqNo: 2829  
Long Name: Prothrombin Complex Concentrate - 4 With Factor VIIa (FEIBA) units/kg - Dose 2  
Short Name: ProCmplxCon4W7a2  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the second dose in units per kilogram of Prothrombin Complex Concentrate - 4 With Factor VIIa (FEIBA). Enter zero if no second dose given.

Low Value: 0      High Value: 200

Parent Long Name: Prothrombin Complex Concentrate - 4 With Factor VIIa (FEIBA) Usage  
Parent Short Name: ProCmplxCon4W7a  
Parent Value(s): = "Yes"

---

SeqNo: 2830  
Long Name: Prothrombin Complex Concentrate - 4 With Factor VIIa (FEIBA) units/kg - Dose 3  
Short Name: ProCmplxCon4W7a3  
Database Table Name: Operations

---

Data Source: User  
Format: Integer  
Definition: Indicate the third dose in units per kilogram of Prothrombin Complex Concentrate - 4 With Factor VIIa (FEIBA). Enter zero if no third dose given.

Low Value: 0 High Value: 200

Parent Long Name: Prothrombin Complex Concentrate - 4 With Factor VIIa (FEIBA) units/kg  
- Dose 2

Parent Short Name: ProCmplxCon4W7a2

Parent Value(s): >0

---

SeqNo: 2831

Long Name: Prothrombin Complex Concentrate - 3 (PCC-3, ProfilNine-SD) Usage

Short Name: ProCmplxCon3

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether Prothrombin Complex Concentrate - 3 (PCC-3, ProfilNine-SD) was administered intraoperatively.

Parent Long Name: Procoagulant Used Intraoperatively

Parent Short Name: ProcoagUsage

Parent Value(s): = "Yes"

---

SeqNo: 2832

Long Name: Prothrombin Complex Concentrate - 3 (PCC-3, ProfilNine-SD) units/kg - Dose 1

Short Name: ProCmplxCon3Ds1

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the first dose in units per kilogram of Prothrombin Complex Concentrate - 3 (PCC-3, ProfilNine-SD).

Low Value: 1 High Value: 5

Parent Long Name: Prothrombin Complex Concentrate - 3 (PCC-3, ProfilNine-SD) Usage

Parent Short Name: ProCmplxCon3

Parent Value(s): = "Yes"

---

SeqNo: 2833

Long Name: Prothrombin Complex Concentrate - 3 (PCC-3, ProfilNine-SD) units/kg - Dose 2

Short Name: ProCmplxCon3Ds2

Database Table Name: Operations

Data Source: User

---



Format: Integer  
Definition: Indicate the second dose in units per kilogram of Prothrombin Complex Concentrate - 3 (PCC-3, ProfilNine-SD). Enter zero if no second dose given.

Low Value: 0 High Value: 5

Parent Long Name: Prothrombin Complex Concentrate - 3 (PCC-3, ProfilNine-SD) Usage  
Parent Short Name: ProCmplxCon3  
Parent Value(s): = "Yes"

---

SeqNo: 2834

Long Name: Prothrombin Complex Concentrate - 3 (PCC-3, ProfilNine-SD) units/kg - Dose 3

Short Name: ProCmplxCon3Ds3

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the third dose in units per kilogram of Prothrombin Complex Concentrate - 3 (PCC-3, ProfilNine-SD). Enter zero if no third dose given.

Low Value: 0 High Value: 5

Parent Long Name: Prothrombin Complex Concentrate - 3 (PCC-3, ProfilNine-SD) units/kg -  
Dose 2

Parent Short Name: ProCmplxCon3Ds2

Parent Value(s): >0

---

SeqNo: 2835

Long Name: Octaplex Prothrombin Concentrate Usage

Short Name: Octaplex

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether Octaplex Prothrombin Concentrate was administered intraoperatively.

Parent Long Name: Procoagulant Used Intraoperatively

Parent Short Name: ProcoagUsage

Parent Value(s): = "Yes"

---

SeqNo: 2836

Long Name: Octaplex Prothrombin Concentrate Units - Dose 1

Short Name: OctaplexDs1

Database Table Name: Operations

Data Source: User

---

Format: Integer  
Definition: Indicate the first dose in international units (IU) of Octaplex Prothrombin Concentrate.

Low Value: 1 High Value: 6000

Parent Long Name: Octaplex Prothrombin Concentrate Usage  
Parent Short Name: Octaplex  
Parent Value(s): = "Yes"

---

SeqNo: 2837

Long Name: Octaplex Prothrombin Concentrate Units - Dose 2  
Short Name: OctaplexDs2  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the second dose in international units (IU) of Octaplex Prothrombin Concentrate.

Low Value: 0 High Value: 6000

Parent Long Name: Octaplex Prothrombin Concentrate Usage  
Parent Short Name: Octaplex  
Parent Value(s): = "Yes"

---

SeqNo: 2838

Long Name: Octaplex Prothrombin Concentrate Units - Dose 3  
Short Name: OctaplexDs3  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the third dose in international units (IU) of Octaplex Prothrombin Concentrate.

Low Value: 0 High Value: 6000

Parent Long Name: Octaplex Prothrombin Concentrate Units - Dose 2  
Parent Short Name: OctaplexDs2  
Parent Value(s): >0

---

SeqNo: 2840

Long Name: Fibrinogen Concentrate Usage  
Short Name: ProcoagFibrin  
Database Table Name: Operations  
Data Source: User

---

Format: Text (categorical values specified by STS)  
Definition: Indicate whether Fibrinogen Concentrate was administered intraoperatively.

Parent Long Name: Procoagulent Used Intraoperatively  
Parent Short Name: ProcoagUsage  
Parent Value(s): = "Yes"

---

SeqNo: 2841  
Long Name: Fibrinogen Concentrate mg/kg - Dose 1  
Short Name: ProcoagFibrin1  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the first dose in mg per kilogram of fibrinogen concentrate given during this procedure.

Low Value: 1      High Value: 100

Parent Long Name: Fibrinogen Concentrate Usage  
Parent Short Name: ProcoagFibrin  
Parent Value(s): = "Yes"

---

SeqNo: 2842  
Long Name: Fibrinogen Concentrate mg/kg - Dose 2  
Short Name: ProcoagFibrin2  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the second dose in mg per kilogram of fibrinogen concentrate given during this procedure. Enter zero if no second dose given.

Low Value: 0      High Value: 100

Parent Long Name: Fibrinogen Concentrate Usage  
Parent Short Name: ProcoagFibrin  
Parent Value(s): = "Yes"

---

SeqNo: 2843  
Long Name: Fibrinogen Concentrate mg/kg - Dose 3  
Short Name: ProcoagFibrin3  
Database Table Name: Operations  
Data Source: User

---

Format: Integer  
Definition: Indicate the third dose in mg per kilogram of fibrinogen concentrate given during this procedure. Enter zero if no third dose given.

Low Value: 0 High Value: 100

Parent Long Name: Fibrinogen Concentrate mg/kg - Dose 2  
Parent Short Name: ProcoagFibrin2  
Parent Value(s): >0

---

SeqNo: 2844  
Long Name: Antithrombin 3 (AT3) Concentrate Usage  
Short Name: ProcoagAntithrom  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether Antithrombin 3 (AT3) Concentrate was administered intraoperatively.  
  
Parent Long Name: Procoagulent Used Intraoperatively  
Parent Short Name: ProcoagUsage  
Parent Value(s): = "Yes"

---

SeqNo: 2845  
Long Name: Antithrombin 3 Concentrate units - Dose 1  
Short Name: ProcoagAntithrom1  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the first dose in units of antithrombin 3 concentrate given during this procedure.  
Low Value: 1 High Value: 5000  
  
Parent Long Name: Antithrombin 3 (AT3) Concentrate Usage  
Parent Short Name: ProcoagAntithrom  
Parent Value(s): = "Yes"

---

SeqNo: 2846  
Long Name: Antithrombin 3 Concentrate units - Dose 2  
Short Name: ProcoagAntithrom2  
Database Table Name: Operations  
Data Source: User  
Format: Integer

---

**Definition:** Indicate the second dose in units of antithrombin 3 concentrate given during this procedure. Enter zero if no second dose given.

**Low Value:** 0      **High Value:** 5000

**Parent Long Name:** Antithrombin 3 (AT3) Concentrate Usage

**Parent Short Name:** ProcoagAntithrom

**Parent Value(s):** = "Yes"

---

**SeqNo:** 2847

**Long Name:** Antithrombin 3 Concentrate units - Dose 3

**Short Name:** ProcoagAntithrom3

**Database Table Name:** Operations

**Data Source:** User

**Format:** Integer

**Definition:** Indicate the third dose in units of antithrombin 3 concentrate given during this procedure. Enter zero if no third dose given.

**Low Value:** 0      **High Value:** 5000

**Parent Long Name:** Antithrombin 3 Concentrate units - Dose 2

**Parent Short Name:** ProcoagAntithrom2

**Parent Value(s):** >0

---

**SeqNo:** 2848

**Long Name:** Desmopressin (DDAVP) Usage

**Short Name:** ProcoagDesmo

**Database Table Name:** Operations

**Data Source:** User

**Format:** Text (categorical values specified by STS)

**Definition:** Indicate whether Desmopressin (DDAVP) was administered intraoperatively.

**Parent Long Name:** Procoagulant Used Intraoperatively

**Parent Short Name:** ProcoagUsage

**Parent Value(s):** = "Yes"

---

**SeqNo:** 2849

**Long Name:** Desmopressin (DDAVP) mcg/kg - Dose 1

**Short Name:** ProcoagDesmo1

**Database Table Name:** Operations

**Data Source:** User

**Format:** Real, 2 decimal places

**Definition:** Indicate the first dose in micrograms per kilogram of desmopressin (DDAVP) given

during this procedure.

Low Value: 0.01 High Value: 6.00

Parent Long Name: Desmopressin (DDAVP) Usage  
 Parent Short Name: ProcoagDesmo  
 Parent Value(s): = "Yes"

SeqNo: 2850

Long Name: Desmopressin (DDAVP) mcg/kg - Dose 2

Short Name: ProcoagDesmo2

Database Table Name: Operations

Data Source: User

Format: Real, 2 decimal places

Definition: Indicate the second dose in micrograms per kilogram of desmopressin (DDAVP) given during this procedure. Enter zero if no second dose given.

Low Value: 0.00 High Value: 6.00

Parent Long Name: Desmopressin (DDAVP) Usage  
 Parent Short Name: ProcoagDesmo  
 Parent Value(s): = "Yes"

SeqNo: 2851

Long Name: Desmopressin (DDAVP) mcg/kg - Dose 3

Short Name: ProcoagDesmo3

Database Table Name: Operations

Data Source: User

Format: Real, 2 decimal places

Definition: Indicate the third dose in micrograms per kilogram of desmopressin (DDAVP) given during this procedure. Enter zero if no third dose given.

Low Value: 0.00 High Value: 6.00

Parent Long Name: Desmopressin (DDAVP) mcg/kg - Dose 2  
 Parent Short Name: ProcoagDesmo2  
 Parent Value(s): >0

SeqNo: 2852

Long Name: Point Of Care Coagulation Testing Utilized Intraoperatively

Short Name: POCCoagTstUtil

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)  
Definition: Indicate whether point of care coagulation testing was utilized intraoperatively.

---

SeqNo: 2853  
Long Name: Point Of Care Coagulation Testing - Thromboelastography (TEG)  
Short Name: POCCoagTstTEG  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether point of care coagulation testing included Thromboelastography (TEG).  
  
Parent Long Name: Point Of Care Coagulation Testing Utilized Intraoperatively  
Parent Short Name: POCCoagTstUtil  
Parent Value(s): = "Yes"

---

SeqNo: 2854  
Long Name: Point Of Care Coagulation Testing - ROTEM  
Short Name: POCCoagTstROTEM  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether point of care coagulation testing included ROTEM.  
  
Parent Long Name: Point Of Care Coagulation Testing Utilized Intraoperatively  
Parent Short Name: POCCoagTstUtil  
Parent Value(s): = "Yes"

---

SeqNo: 2855  
Long Name: Point Of Care Coagulation Testing - Sonoclot  
Short Name: POCCoagTstSon  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether point of care coagulation testing included Sonoclot.  
  
Parent Long Name: Point Of Care Coagulation Testing Utilized Intraoperatively  
Parent Short Name: POCCoagTstUtil  
Parent Value(s): = "Yes"

---

SeqNo: 2856  
Long Name: Point Of Care Coagulation Testing - Heparin Concentration (Hepcon, HMS)  
Short Name: POCCoagTstHep

---

Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether point of care coagulation testing included Heparin Concentration (Hepcon, HMS).

Parent Long Name: Point Of Care Coagulation Testing Utilized Intraoperatively  
Parent Short Name: POCCoagTstUtil  
Parent Value(s): = "Yes"

---

SeqNo: 2857  
Long Name: Point Of Care Coagulation Testing - INR/PT/aPTT (iStat or equivalent)  
Short Name: POCCoagTstINR  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether point of care coagulation testing included INR/PT/aPTT (iStat or equivalent).

Parent Long Name: Point Of Care Coagulation Testing Utilized Intraoperatively  
Parent Short Name: POCCoagTstUtil  
Parent Value(s): = "Yes"

---

## 12. CABG Procedures

SeqNo: 2900  
Long Name: CAB  
Short Name: OpCAB  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether coronary artery bypass grafting was done.

Parent Long Name: Operation Type  
Parent Short Name: OpType  
Parent Value(s): = "CPB" or "No CPB Cardiovascular"

---

SeqNo: 2910  
Long Name: Dist Anast - Art #  
Short Name: DistArt  
Database Table Name: Operations

---



Data Source: User  
Format: Integer  
Definition: Indicate the total number of distal anastomoses with arterial conduits, whether IMA, radial artery, etc.

Low Value: 0 High Value: 9

Parent Long Name: CAB  
Parent Short Name: OpCAB  
Parent Value(s): = "Yes"

---

SeqNo: 2920  
Long Name: Dist Anast - Vein #  
Short Name: DistVein  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the total number of distal anastomoses with venous conduits.

Low Value: 0 High Value: 9

Parent Long Name: CAB  
Parent Short Name: OpCAB  
Parent Value(s): = "Yes"

SeqNo: 2930  
Long Name: IMA Artery Used  
Short Name: IMAArtUs  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate which, if any, Internal Mammary Artery(ies) (IMA) were used for grafts.

Parent Long Name: CAB  
Parent Short Name: OpCAB  
Parent Value(s): = "Yes"

---

## 13. Valve Procedures

SeqNo: 2940  
Long Name: Valve  
Short Name: OpValve

---

Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a surgical procedure was done on the Aortic, Mitral, Tricuspid, Pulmonic, common AV valve or truncal valve.

Parent Long Name: Operation Type  
Parent Short Name: OpType  
Parent Value(s): = "CPB" or "No CPB Cardiovascular"

---

SeqNo: 3140  
Long Name: Valve Device Explanted And/Or Implanted  
Short Name: ValExImp  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a valve device of any type was explanted and/or implanted during this procedure.

Parent Long Name: Valve  
Parent Short Name: OpValve  
Parent Value(s): = "Yes"

---

September 2017: History of severe pulmonary insufficiency status post pulmonary valvotomy for congenital pulmonary stenosis. Now having RVOT reconstruction with 24 mm expanded polytetrafluoroethylene (ePTFE) valve conduit. Do I mark that a valve was explanted and implanted or just implanted? **Valve implanted only.**

---

SeqNo: 3150  
Long Name: Valve Explant Type #1  
Short Name: ValExType1  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the type of the first valve or device explanted

Parent Long Name: Valve Device Explanted And/Or Implanted  
Parent Short Name: ValExImp  
Parent Value(s): = "Yes, Explanted" or "Yes, Explanted and Implanted"

---

SeqNo: 3151  
Long Name: Valve Explant Unique Device Identifier (UDI) - 1  
Short Name: ValExpUDI1  
Database Table Name: Operations

---

Data Source: User  
Format: Text Data Length: 75  
Definition: Indicate the Unique Device Identifier (UDI) of the first explanted valve device if available, otherwise leave blank.

Parent Long Name: Valve Device Explanted And/Or Implanted  
Parent Short Name: ValExImp  
Parent Value(s): = "Yes, Explanted" or "Yes, Explanted and Implanted"

---

SeqNo: 3160  
Long Name: Second Valve Explanted or Device Removed  
Short Name: ValEx2  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a second valve or device was explanted

Parent Long Name: Valve Device Explanted And/Or Implanted  
Parent Short Name: ValExImp  
Parent Value(s): = "Yes, Explanted" or "Yes, Explanted and Implanted"

---

SeqNo: 3170  
Long Name: Valve Explant Type #2  
Short Name: ValExType2  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the type of the second valve or device explanted

Parent Long Name: Second Valve Explanted or Device Removed  
Parent Short Name: ValEx2  
Parent Value(s): = "Yes"

---

SeqNo: 3171  
Long Name: Valve Explant Unique Device Identifier (UDI) - 2  
Short Name: ValExpUDI2  
Database Table Name: Operations  
Data Source: User  
Format: Text Data Length: 75  
Definition: Indicate the Unique Device Identifier (UDI) of the second explanted valve device if available, otherwise leave blank.

Parent Long Name: Second Valve Explanted or Device Removed

---

Parent Short Name: ValEx2  
Parent Value(s): = "Yes"

---

SeqNo: 3180  
Long Name: Third Valve Explanted or Device Removed  
Short Name: ValEx3  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a third valve or device was explanted

Parent Long Name: Second Valve Explanted or Device Removed  
Parent Short Name: ValEx2  
Parent Value(s): = "Yes"

---

SeqNo: 3190  
Long Name: Valve Explant Type #3  
Short Name: ValExType3  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the type of the third valve or device explanted

Parent Long Name: Third Valve Explanted or Device Removed  
Parent Short Name: ValEx3  
Parent Value(s): = "Yes"

---

SeqNo: 3191  
Long Name: Valve Explant Unique Device Identifier (UDI) - 3  
Short Name: ValExpUDI3  
Database Table Name: Operations  
Data Source: User  
Format: Text Data Length: 75  
Definition: Indicate the Unique Device Identifier (UDI) of the third explanted valve device if available, otherwise leave blank.

Parent Long Name: Third Valve Explanted or Device Removed  
Parent Short Name: ValEx3  
Parent Value(s): = "Yes"

---

SeqNo: 3200  
Long Name: Fourth Valve Explanted or Device Removed  
Short Name: ValEx4  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a fourth valve or device was explanted

Parent Long Name: Third Valve Explanted or Device Removed  
Parent Short Name: ValEx3  
Parent Value(s): = "Yes"

SeqNo: 3210  
Long Name: Valve Explant Type #4  
Short Name: ValExType4  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the type of the fourth valve or device explanted

Parent Long Name: Fourth Valve Explanted or Device Removed  
Parent Short Name: ValEx4  
Parent Value(s): = "Yes"

---

SeqNo: 3211  
Long Name: Valve Explant Unique Device Identifier (UDI) - 4  
Short Name: ValExpUDI4  
Database Table Name: Operations  
Data Source: User  
Format: Text Data Length: 75  
Definition: Indicate the Unique Device Identifier (UDI) of the fourth explanted valve device if available, otherwise leave blank.

Parent Long Name: Fourth Valve Explanted or Device Removed  
Parent Short Name: ValEx4  
Parent Value(s): = "Yes"

---

SeqNo: 3220  
Long Name: Valve Implant Location #1  
Short Name: ValImpLoc1  
Database Table Name: Operations  
Data Source: User

---

Format: Text (categorical values specified by STS)  
Definition: Indicate the location of the first valve or device implanted

Parent Long Name: Valve Device Explanted And/Or Implanted  
Parent Short Name: ValExImp  
Parent Value(s): = "Yes, Implanted" or "Yes, Explanted and Implanted"

---

SeqNo: 3230  
Long Name: Valve Implant Type #1  
Short Name: VallmpType1  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the type of the first valve or device implanted

Parent Long Name: Valve Device Explanted And/Or Implanted  
Parent Short Name: ValExImp  
Parent Value(s): = "Yes, Implanted" or "Yes, Explanted and Implanted"

---

SeqNo: 3240  
Long Name: Valve Implant Surgeon Fashioned Material #1  
Short Name: VallmpSFMat1  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the material used to fashion the first valve or device

Parent Long Name: Valve Implant Type #1  
Parent Short Name: VallmpType1  
Parent Value(s): = "Surgeon fashioned"

SeqNo: 3250  
Long Name: Valve Implant Commercial Device Model Number #1  
Short Name: VallmpComMod1  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the name of the prosthesis implanted. The names provided include the manufacturer's model number with "xx" substituting for the device size. Note that the model number is different from the serial number.

Parent Long Name: Valve Implant Type #1  
Parent Short Name: VallmpType1

---

Parent Value(s): = "Commercially supplied device"

---

SeqNo: 3260

Long Name: Valve Implant Commercial Device Size #1

Short Name: VallmpComSz1

Database Table Name: Operations

Data Source: User

Format: Integer

Definition: Indicate the size of the first implanted valve or device

Low Value: 5 High Value: 50

Parent Long Name: Valve Implant Type #1

Parent Short Name: VallmpType1

Parent Value(s): = "Commercially supplied device"

---

SeqNo: 3261

Long Name: Valve Implant Unique Device Identifier (UDI) - 1

Short Name: VallmpUDI1

Database Table Name: Operations

Data Source: User

Format: Text Data Length: 75

Definition: Indicate the Unique Device Identifier (UDI) of the first implanted valve device if available, otherwise leave blank.

Parent Long Name: Valve Device Explanted And/Or Implanted

Parent Short Name: ValExImp

Parent Value(s): = "Yes, Implanted" or "Yes, Explanted and Implanted"

---

SeqNo: 3270

Long Name: Second Valve Implant

Short Name: Vallmp2

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether a second valve or device was implanted

Parent Long Name: Valve Device Explanted And/Or Implanted

Parent Short Name: ValExImp

Parent Value(s): = "Yes, Implanted" or "Yes, Explanted and Implanted"

---

SeqNo: 3280

Long Name: Valve Implant Location #2

---

Short Name: VallImpLoc2  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the location of the second valve or device implanted

Parent Long Name: Second Valve Implant  
Parent Short Name: VallImp2  
Parent Value(s): = "Yes"

---

SeqNo: 3290  
Long Name: Valve Implant Type #2  
Short Name: VallImpType2  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the type of the second valve or device implanted

Parent Long Name: Second Valve Implant  
Parent Short Name: VallImp2  
Parent Value(s): = "Yes"

---

SeqNo: 3300  
Long Name: Valve Implant Surgeon Fashioned Material #2  
Short Name: VallImpSFMat2  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the material used to fashion the second valve or device

Parent Long Name: Valve Implant Type #2  
Parent Short Name: VallImpType2  
Parent Value(s): = "Surgeon fashioned"

---

SeqNo: 3310  
Long Name: Valve Implant Commercial Device Model Number #2  
Short Name: VallImpComMod2  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the name of the prosthesis implanted. The names provided include the manufacturer's model number with "xx" substituting for the device size.

---



Parent Long Name: Valve Implant Type #2  
Parent Short Name: VallmpType2  
Parent Value(s): = "Commercially supplied device"

---

SeqNo: 3320  
Long Name: Valve Implant Commercial Device Size #2  
Short Name: VallmpComSz2  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the size of the second implanted valve or device

Low Value: 5 High Value: 50  
Parent Long Name: Valve Implant Type #2  
Parent Short Name: VallmpType2  
Parent Value(s): = "Commercially supplied device"

---

SeqNo: 3321  
Long Name: Valve Implant Unique Device Identifier (UDI) - 2  
Short Name: VallmpUDI2  
Database Table Name: Operations  
Data Source: User  
Format: Text Data Length: 75  
Definition: Indicate the Unique Device Identifier (UDI) of the second implanted valve device if available, otherwise leave blank.

Parent Long Name: Second Valve Implant  
Parent Short Name: Vallmp2  
Parent Value(s): = "Yes"

---

SeqNo: 3330  
Long Name: Third Valve Implant  
Short Name: Vallmp3  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a third valve or device was implanted

Parent Long Name: Second Valve Implant  
Parent Short Name: Vallmp2  
Parent Value(s): = "Yes"

---

---

SeqNo: 3340  
Long Name: Valve Implant Location #3  
Short Name: VallImpLoc3  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the location of the third valve or device implanted

Parent Long Name: Third Valve Implant  
Parent Short Name: VallImp3  
Parent Value(s): = "Yes"

---

SeqNo: 3350  
Long Name: Valve Implant Type #3  
Short Name: VallImpType3  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the type of the third valve or device implanted

Parent Long Name: Third Valve Implant  
Parent Short Name: VallImp3  
Parent Value(s): = "Yes"

---

SeqNo: 3360  
Long Name: Valve Implant Surgeon Fashioned Material #3  
Short Name: VallImpSFMat3  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the material used to fashion the third valve or device

Parent Long Name: Valve Implant Type #3  
Parent Short Name: VallImpType3  
Parent Value(s): = "Surgeon fashioned"

---

SeqNo: 3370  
Long Name: Valve Implant Commercial Device Model Number #3  
Short Name: VallImpComMod3  
Database Table Name: Operations

---

Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the name of the prosthesis implanted. The names provided include the manufacturer's model number with "xx" substituting for the device size.

Parent Long Name: Valve Implant Type #3  
Parent Short Name: VallmpType3  
Parent Value(s): = "Commercially supplied device"

---

SeqNo: 3380  
Long Name: Valve Implant Commercial Device Size #3  
Short Name: VallmpComSz3  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the size of the third implanted valve or device

Low Value: 5 High Value: 50

Parent Long Name: Valve Implant Type #3  
Parent Short Name: VallmpType3  
Parent Value(s): = "Commercially supplied device"

---

SeqNo: 3381  
Long Name: Valve Implant Unique Device Identifier (UDI) - 3  
Short Name: VallmpUDI3  
Database Table Name: Operations  
Data Source: User  
Format: Text Data Length: 75  
Definition: Indicate the Unique Device Identifier (UDI) of the third implanted valve device if available, otherwise leave blank.

Parent Long Name: Third Valve Implant  
Parent Short Name: Vallmp3  
Parent Value(s): = "Yes"

---

SeqNo: 3390  
Long Name: Fourth Valve Implant  
Short Name: Vallmp4  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)

---

Definition: Indicate whether a fourth valve or device was implanted

Parent Long Name: Third Valve Implant

Parent Short Name: Vallmp3

Parent Value(s): = "Yes"

---

SeqNo: 3400

Long Name: Valve Implant Location #4

Short Name: VallmpLoc4

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate the location of the fourth valve or device implanted

Parent Long Name: Fourth Valve Implant

Parent Short Name: Vallmp4

Parent Value(s): = "Yes"

---

SeqNo: 3410

Long Name: Valve Implant Type #4

Short Name: VallmpType4

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate the type of the fourth valve or device implanted

Parent Long Name: Fourth Valve Implant

Parent Short Name: Vallmp4

Parent Value(s): = "Yes"

---

SeqNo: 3420

Long Name: Valve Implant Surgeon Fashioned Material #4

Short Name: VallmpSFMat4

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate the material used to fashion the fourth valve or device

Parent Long Name: Valve Implant Type #4

Parent Short Name: VallmpType4

Parent Value(s): = "Surgeon fashioned"

---

SeqNo: 3430  
Long Name: Valve Implant Commercial Device Model Number #4  
Short Name: VallImpComMod4  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the name of the prosthesis implanted. The names provided include the manufacturer's model number with "xx" substituting for the device size.

Parent Long Name: Valve Implant Type #4  
Parent Short Name: VallImpType4  
Parent Value(s): = "Commercially supplied device"

---

SeqNo: 3440  
Long Name: Valve Implant Commercial Device Size #4  
Short Name: VallImpComSz4  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the size of the fourth implanted valve or device

Low Value: 5 High Value: 50  
Parent Long Name: Valve Implant Type #4  
Parent Short Name: VallImpType4  
Parent Value(s): = "Commercially supplied device"

---

SeqNo: 3441  
Long Name: Valve Implant Unique Device Identifier (UDI) - 4  
Short Name: VallImpUDI4  
Database Table Name: Operations  
Data Source: User  
Format: Text Data Length: 75  
Definition: Indicate the Unique Device Identifier (UDI) of the fourth implanted valve device if available, otherwise leave blank.

Parent Long Name: Fourth Valve Implant  
Parent Short Name: VallImp4  
Parent Value(s): = "Yes"

---

## 14. VAD Procedures

---

SeqNo: 3460

Long Name: VAD Explanted And/Or Implanted

Short Name: VADExImp

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether a ventricular assist device (VAD) was explanted and/or implanted during this procedure.

Parent Long Name: Operation Type

Parent Short Name: OpType

Parent Value(s): = "CPB Cardiovascular", "No CPB Cardiovascular", "CPB Non-Cardiovascular", "ECMO", "Thoracic", "VAD Operation Done With CPB", "VAD Operation Done Without CPB." or "Other"

---

SeqNo: 3500

Long Name: VAD-Indication for VAD

Short Name: VADInd

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate the reason the patient is receiving the ventricular assist device (VAD).

Parent Long Name: VAD Explanted And/Or Implanted

Parent Short Name: VADExImp

Parent Value(s): = "Yes, implanted" or "Yes, explanted and implanted"

---

SeqNo: 3560

Long Name: VAD-Implant Type

Short Name: VImpTy

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate the initial type of VAD implanted.

Parent Long Name: VAD Explanted And/Or Implanted

Parent Short Name: VADExImp

Parent Value(s): = "Yes, implanted" or "Yes, explanted and implanted"

---

September 2017: Question: Both a Berlin Heart LVAD and a Rotaflow RVAD were implanted in the same operation. What should I choose for Implant Type? **There is no current way to code both implanted VADS. This will be addressed with the upcoming spec upgrade. Perhaps code the VAD of the surgeons' choice.**

November 2017: A VAD was placed in a single ventricle patient, supporting his systemic circulation via his morphologic RV. Would you consider this an LVAD? **LVADs and RVADs short list to VAD but some software vendors force a decision to be made. Be consistent within your facility and what is going on with the patient specifically.**

SeqNo: 3570

Long Name: VAD-Product Type

Short Name: VProdTy

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate the specific product implanted. Implant defined as physical placement of the VAD.

Parent Long Name: VAD Explanted And/Or Implanted

Parent Short Name: VADExImp

Parent Value(s): = "Yes, implanted" or "Yes, explanted and implanted"

SeqNo: 3571

Long Name: VAD Implant Unique Device Identifier (UDI)

Short Name: VADImpUDI

Database Table Name: Operations

Data Source: User

Format: Text Data Length: 75

Definition: Indicate the Unique Device Identifier (UDI) of the implanted VAD if available, otherwise leave blank.

Parent Long Name: VAD Explanted And/Or Implanted

Parent Short Name: VADExImp

Parent Value(s): = "Yes, implanted" or "Yes, explanted and implanted"

SeqNo: 3610

Long Name: VAD-Explant Reason

Short Name: VExpRsn

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate the reason the VAD was explanted.

Parent Long Name: VAD Explanted And/Or Implanted  
Parent Short Name: VADExImp  
Parent Value(s): = "Yes, explanted" or "Yes, explanted and implanted"

---

SeqNo: 3611  
Long Name: VAD Explant Unique Device Identifier (UDI)  
Short Name: VADExUDI  
Database Table Name: Operations  
Data Source: User  
Format: Text  
Definition: Indicate the Unique Device Identifier (UDI) of the explanted VAD if available, otherwise leave blank.  
Data Length: 75

Parent Long Name: VAD Explanted And/Or Implanted  
Parent Short Name: VADExImp  
Parent Value(s): = "Yes, explanted" or "Yes, explanted and implanted"

---

SeqNo: 3850  
Long Name: VAD-Primary VAD Comp-Intracranial Bleed  
Short Name: PVCmpBld  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate if the patient had an intracranial bleed, confirmed by CT scan or other diagnostic studies.

Parent Long Name: VAD Explanted And/Or Implanted  
Parent Short Name: VADExImp  
Parent Value(s): = "Yes, explanted", "Yes, implanted" or "Yes, explanted and implanted"

---

SeqNo: 3860  
Long Name: VAD-Primary VAD Comp-Embolic Stroke  
Short Name: PVCmpES  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate if the patient had embolic stroke caused by a blood clot, air embolus, or tissue, confirmed by CT scan or other diagnostic studies.

Parent Long Name: VAD Explanted And/Or Implanted  
Parent Short Name: VADExImp

---



Parent Value(s): = "Yes, explanted", "Yes, implanted" or "Yes, explanted and implanted"

---

SeqNo: 3870

Long Name: VAD-Primary VAD Comp-Driveline and/or cannula Infection

Short Name: PVCmpDCI

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate if the patient had a driveline and/or cannula infection. Driveline and/or cannula infection is defined as the presence of erythema, drainage, or purulence at the VAD connection site whether entering or exiting the body in association with leukocytosis and in the presence of positive culture.

Parent Long Name: VAD Explanted And/Or Implanted

Parent Short Name: VAExImp

Parent Value(s): = "Yes, explanted", "Yes, implanted" or "Yes, explanted and implanted"

---

SeqNo: 3880

Long Name: VAD-Primary VAD Comp-Pump Pocket Infection

Short Name: PVCmpPPI

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate if the patient had a pump pocket infection. A pump pocket infection is defined as a persistent drainage in the physical location of the pump, located preperitoneally or intra-abdominally with positive cultures from the pocket site.

Parent Long Name: VAD Explanted And/Or Implanted

Parent Short Name: VAExImp

Parent Value(s): = "Yes, explanted", "Yes, implanted" or "Yes, explanted and implanted"

---

SeqNo: 3890

Long Name: VAD-Primary VAD Comp-VAD Endocarditis

Short Name: PVCmpEnd

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate if the patient had VAD endocarditis. VAD endocarditis is defined as an infection of the blood contacting surface of the VAD device itself. This may include:

- internal surfaces;
- graft material;
- inflow/outflow valves of the VAD.

---

Parent Long Name: VAD Explanted And/Or Implanted  
Parent Short Name: VADExImp  
Parent Value(s): = "Yes, explanted", "Yes, implanted" or "Yes, explanted and implanted"

---

SeqNo: 3900  
Long Name: VAD-Primary VAD Comp-Device Malfunction  
Short Name: PVCmpMal  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate if the pump itself is not functioning properly causing hemodynamic compromise, and/or requiring immediate intervention or VAD replacement.

Parent Long Name: VAD Explanted And/Or Implanted  
Parent Short Name: VADExImp  
Parent Value(s): = "Yes, explanted", "Yes, implanted" or "Yes, explanted and implanted"

---

SeqNo: 3910  
Long Name: VAD-Primary VAD Comp-Bowel Obstruction  
Short Name: PVCmpBO  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate if the patient was diagnosed with a bowel obstruction post VAD insertion by documentation in the medical record.

Parent Long Name: VAD Explanted And/Or Implanted  
Parent Short Name: VADExImp  
Parent Value(s): = "Yes, explanted", "Yes, implanted" or "Yes, explanted and implanted"

---

SeqNo: 3920  
Long Name: VAD-Primary VAD Comp-Hemolysis  
Short Name: PVCmpHemo  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate if the patient was diagnosed with hemolysis post VAD insertion by documentation in the medical record.

Parent Long Name: VAD Explanted And/Or Implanted  
Parent Short Name: VADExImp

---

Parent Value(s): = "Yes, explanted", "Yes, implanted" or "Yes, explanted and implanted"

---

## 15. Complications

SeqNo: 4180

Long Name: Complications Table Unique Record Identifier  
Short Name: CompUniqueID  
Database Table Name: Complications  
Data Source: Automatic  
Format: Text  
Definition: Unique identifier for the record in the Complications table.

---

SeqNo: 4190

Long Name: Complications Link to Operations Table  
Short Name: OperationID  
Database Table Name: Complications  
Data Source: Automatic  
Format: Text  
Definition: An arbitrary, unique value generated by the software that permanently identifies each operation record in the participant's database. This field is the foreign key that links the Complications record with the associated record in the Operations table.

---

SeqNo: 4200

Long Name: Complication  
Short Name: Complication  
Database Table Name: Complications  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Assign complication to the operation that is most closely associated with the complication. A complication is an event or occurrence that is associated with a disease or a healthcare intervention, is a departure from the desired course of events, and may cause, or be associated with, suboptimal outcome. A complication does not necessarily represent a breach in the standard of care that constitutes medical negligence or medical malpractice. An operative or procedural complication is any complication, regardless of cause, occurring (1) within 30 days after surgery or intervention in or out of the hospital, or (2) after 30 days during the same hospitalization subsequent to the operation or intervention. Operative and procedural complications include both intraoperative/intraprocedural complications and postoperative/postprocedural complications in this time interval.

---

An adverse event is a complication that is associated with a healthcare intervention and is associated with suboptimal outcome. Adverse events represent a subset of complications. Not all medical errors result in an adverse event; the administration of an incorrect dose of a medication is a medical error, but it does not always result in an adverse event. Similarly, not all adverse events are the result of medical error. A child may develop pneumonia after an atrial septal defect repair despite intra- and peri-operative management that is free of error. Complications of the underlying disease state, which are not related to a medical intervention, are not adverse events. For example, a patient who presents for medical care with metastatic lung cancer has already developed a complication (Metastatic spread) of the primary lung cancer without any healthcare intervention. Furthermore, complications not associated with suboptimal outcome or harm are not adverse events and are known as no harm events. The patient who receives an incorrect dose of a medication without harm has experienced a no harm event, but not an adverse event.

---

12/2015: If a patient has Pulmonary Hypertension preoperatively, should this still be coded as a complication if it is present post op as well? **Call it a complication if series of events happen, if there is a clinical or significant change in patient's condition.**

February 2016: If a post-op patient is extubated and re-intubated multiple times due to respiratory failure, but never for 7 consecutive days, do you code complication 150? Is it 7 days consecutively or 7 days cumulatively? **Consecutively**

February 2016: if a patient has surgery is extubated after several days, and then 3 weeks later the patient requires reintubation and eventual trach, would that be considered a post-op complication for the cardiac surgery? **Yes**

February 2016: If a patient requires a tracheostomy and it is discussed and planned 1-2 days prior to the trach being performed, would that still be considered an "unplanned non-cardiac reoperation during the postoperative time period" or would it be considered planned? **code an unplanned non-cardiac reoperation during the post-operative time period. While there is pre-surgical planning for the trach but it is an unplanned non-cardiac reoperation during this admission.**

April 2016: Please clarify the difference between "Thrombus, Central Vein" and "Systemic Vein Obstruction" - we can for see many situations with overlap - with the rare exception of a systemic vein that is occluded by something else, those will mostly be due to thrombus. Doesn't seem like a femoral vein occlusion, for example, should get 2 complications. **Complications aren't "added up"; a number of complications do overlap, pick what best fits the situation.**

**Systemic Vein Obstruction [120] is an option for Complication, as defined: "Clinically significant stenosis or obstruction of any major systemic vein (e.g., superior vena cava, inferior vena cava, femoral veins, internal jugular veins, etc.). A clinically significant event or condition is an event or condition that necessitates a change in treatment."**

**Complications related to thrombus/thrombosis/thromboembolism:**

- 1) intracardiac thrombus – a mass of platelets, fibrin, other blood elements (and potentially additional matter) located at least partially within one or more of the four chambers of the heart (may extend to or from a great vein or artery)
- 2) central vein thrombus – a mass of platelets, fibrin, other blood elements (and potentially additional matter) located in any of the major veins of the body within the space shared with the thoracic and abdominal organs

- 3) **peripheral deep vein thrombus** - a mass of platelets, fibrin, other blood elements (and potentially additional matter) located in any of the deep veins of the extremities (e.g. popliteal, femoral, cephalic, brachial, axillary, etc.) or the extra-thoracic portion of the internal jugular vein
- 4) **systemic-to-pulmonary shunt thrombosis** - a mass of platelets, fibrin, other blood elements (and potentially additional matter) occupying the lumen of a systemic-to-pulmonary artery shunt – may obstructive, occlusive, or neither.
- 5) **pulmonary artery thrombosis/thromboembolism** - a mass of platelets, fibrin, other blood elements (and potentially additional matter) located at least partially within the main pulmonary trunk, right or left pulmonary artery, or their respective branches. The thrombus may have developed in this location (in situ) or may have embolized from another point of origin and lodged within the pulmonary arteries.

April 2016: Is an unplanned sternal wound revision and closure considered an unplanned cardiac reoperation, or unplanned non-cardiac reoperation? **Cardiac Reoperation**

April 2016: How would we code a surgery for abdominal pacemaker pocket hematoma evacuation? In OR, pocket reopened, cleaned, etc. Cardiac or non-cardiac reoperation? Bleeding requiring reoperation? It was bleeding issue, he had to open pocket and clean up, stop oozing, etc. **Bleeding, Cardiac Reoperation**

Part two to this question: If hematoma develops and patient needs reop for bleeding, is that coded as bleeding requiring reop only? Or should there also be a second code for the hematoma (“other”)? **Can use ‘other’, if desired.**

April 2016: I have a question about listing complications for a patient who has had multiple procedures during one hospitalization. An example (not a real patient):

1/1/2016 Norwood procedure, 1/3/2016 Delayed sternal closure, 1/7/2016 Drainage of pericardial effusion, 1/20/2016 Extubation, 1/22/2016 Reintubation. So the complications would be ventilation longer than 7 days, reintubation, pericardial effusion and unexpected cardiac operation. I have been listing the long ventilation and reintubation for all operations, but the pericardial effusion and unexpected operation for the index operation. Do you know if I should be doing this differently? **No. Don’t have to code the re-intubation.**

May 2016: Patient had several STS complications while on ECMO, prior to the 3rd surgery of the admission, which was CPB. My understanding is that all complications will then be attributed to the CPB surgery. Is this correct? If so how do we handle this at our institution? 1st surgery of admission - ECMO cannulation (no other procedures done within the case) 2nd surgery of admission - ECMO repositioning (no other procedures done within the case) 3rd surgery of admission - Tricuspid valvuloplasty (CPB) **All complications regardless of their timing of occurrence will be attributed to the index procedure. Complications from operations that occur prior to the index operation can be included as preoperative factors, where applicable, but should not be recorded in the complications section of those procedures.**

May 2016: For complication #200 "pleural effusion, requiring drainage", does this only get indicated when a patient is given a new chest tube specifically for pleural drainage? What if a patient comes out of the OR with a chest tube and develops a pleural effusion that drains for 8 days, does that need to be indicated if it's the original chest tube the patient came out of the OR with? **This is not a complication. Only include complication “Pleural effusion, requiring drainage” if a new chest tube/drainage procedure was performed. Do not include this complication for an indwelling chest tube or persistent chest tube drainage.**

July 2016: When a patient’s chest is left open post operatively, do further surgical interventions such as mediastinal exploration, removal of clot or re-do of the repair count as an unplanned return to surgery? Our

surgeon is telling us that if the chest is open, further surgical procedures do not count as unplanned return to surgery. **If the sternum is left open at the time of surgery (either sternum left open planned, or sternum left open unplanned), then the first instance of delayed sternal closure is not considered an unplanned surgical intervention. All of the other procedures listed as examples above would be considered unplanned re-interventions.**

August 2016: vocal cord dysfunction: patient has been diagnosed with Papilloma virus. Would this still be considered a complication? **If vocal cord dysfunction is present it is a complication, regardless of the cause. You can code as 'other'.**

October 2016: If a patient is put on ECMO as their only procedure, should "postoperative mechanical circulatory support" be coded as a complication? **If ECMO is the primary procedure then don't code it as a complication.**

November 2016: During a redo sternotomy the surgeon was taking down adhesions on/near the heart using a bovie device. The patient sustained a VF arrest and was quickly defibrillated. The surgeon feels that this episode of VF was due to the electrocautery device disrupting the electrical circuit of the heart and was not an entirely unexpected event. Rather than a complication of surgery, he sees this more as one of those things that can happen and is then quickly rectified. Should this be included as a complication under "Unexpected cardiac arrest during or following procedure" since the surgeon feels it was something that was known to happen?

**Yes, you should code this.**

December 2016: If a child comes in in complete heart block from a prior surgery for a pacemaker generator change, I do not code arrhythmia requiring permanent pacemaker for this surgery correct? It was already coded at his initial repair which resulted in CHB. **This is not a complication.**

March 2017: Do I code these complications related to underlying disease process that occur after index procedure as complications? Or do we only code complications related to the procedure? i.e. one patient had PDA closure without complications from that, but also has Cri-du-Chat syndrome and is still in the hospital r/t Cri-Du-Chat. Has been on a vent. Another patient was severely premature and has complications related to the prematurity, not the PDA closure. **Record all postop events that are listed in the complications list regardless of any other concurrent disease processes. If the patient remained ventilated for greater than 7 days postoperatively for any reason, this complication should be selected.**

April 2017: Would a mediastinal exploration for tamponade be coded as a reop for bleeding, or just an unplanned cardiac reop? **Re-op for bleeding.**

May 2017: A patient goes to OR and postoperatively has an ultrasound revealing bilateral diaphragmatic movement. Unfortunately, the child had a prolonged stay related to co-morbidities and developed critical illness myopathy resulting in diaphragm paresis. Should this be included in the complications/major complication category? **Yes as the complications are encompassing of all post-operative events. It does not matter what caused the diaphragm paresis but the paresis did occur and should be captured.**

June 2017: Should infections other than sepsis or wound infections be captured under "Other complication". Patient had a nasal swab right before surgery on 3/23 which grew enterobacter & h. flu. He got his usual pre-op and post-op antibiotics x 48 hours. On 5/25, a respiratory culture was done and grew enterobacter and 7 days of IV antibiotics were given. Would this count as post-op complication? Or would it not count at all because of the nasal swab pre-op growing enterobacter? **Currently there is no way to capture an infection outside of wound infection or pneumonia. There is no requirement to include 'other' complications but you are able to for local use if desired.**

June 2017: I would like a clarification on the time period of complications. On the monthly calls, I hear people say if there is more than one surgery in a hospitalization, they put all the complications on the very first index procedure and then put "none" for any other surgeries. But looking at the definition "(1) within 30 days after surgery or intervention in or out of the hospital, or (2) after 30 days during the same hospitalization subsequent to the operation or intervention". For example, in one hospitalization, if we had a PDA ligation & atrial

septostomy on CPB in October then a B-T shunt in December followed by a trach two weeks later, would the trach go on the PDA case or the B-T shunt? **You can attribute the complications to the surgery it is most likely associated with, however upon analysis, DCRI will attribute all of the complications to the index operation.**

June 2017: For complications that occur outside of the hospital but within 30 days of the patient's index operation, do we code all that occur within the 30 days? **Yes** Or, if the patient is readmitted, do we capture all complications that occur within the subsequent admission regardless of timing? **All complications are captured for the entire hospital length of stay or if discharged, through post-operative day 30.**

July 2017: If a patient has vocal cord paralysis, but it is due to a transesophageal fistula repair done by general surgery, should this be recorded as a post op complication, as it was not due to the cardiac surgery? **Yes, all complications should be captured regardless of the cause.**

October 2017: Patient had repair of unbalanced left dominant atrioventricular septal defect and ligation of patent arterial duct, and subsequently developed complete heart block requiring insertion of a permanent pacemaker. We would code a complication of arrhythmia necessitating permanent pacemaker and unplanned cardiac reoperation, correct? **Yes, capture all complications/events that occurred in the intraoperative/postoperative time period even if there is overlap. If the pacemaker was inserted in the operating room, select both complications. If the pacemaker was inserted in the cath lab, select only permanent pacemaker.**

November 2017: If a patient is placed on ECMO after the index surgery, does the ECMO count as an unplanned cardiac re-op? **No, include the complication of post-operative mechanical circulatory support only as ECMO**

November 2017: If a patient is having a Hybrid procedure "Stage 1", application of RPA and LPA bands in the cath lab along with a cath procedure. During the procedure the patient arrested and received CPR. Would this complication be coded as 'unexpected cardiac arrest' or 'complication of cardiac cath' procedure? **Unexpected cardiac arrest should be coded as this was a cardiac surgery procedure.**

## 16. Discharge/Readmission

SeqNo: 4220

Long Name: Date of Hospital Discharge

Short Name: HospDischDt

Database Table Name: Operations

Data Source: User

Format: Date - mm/dd/yyyy

Definition: Indicate the date that the patient is discharged from the hospital where the surgery took place. In rare instances, the "Date of Hospital Discharge" differs from the "Date of Database Discharge". In situations where the patient is discharged to another acute care facility or to a chronic care facility, the "Date of Hospital Discharge" is the date the patient is transferred from the hospital where the surgery took place to another facility. This field is intended to capture the total length of stay in your hospital regardless of the



medical service managing the patient.

SeqNo: 4230

Long Name: Mortality Status At Hospital Discharge

Short Name: MtHospDisStat

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether the patient was Alive or Dead at date and time of "Date of Hospital Discharge" for this operation.

May 2017: If a newborn is delivered at hospital A and has cardiac surgery then transferred to hospital B and has cardiac surgery and then transferred back to hospital A months later and subsequently dies at hospital A. Both Hospital A and Hospital B would code that patient as a discharge mortality. The patient was never home from the beginning. This issue is not addressed in the Rules to determine the EACTS-STS discharge date. And thus both hospitals would count the case as a mortality? **Yes, both programs would have a discharge mortality and operative mortality.**

July 2017: Baby who went to the cath lab for pulmonary valve dilation and had a right ventricle perforation and who needed to go to the OR emergently for decompensation. Perforation repaired on bypass due to its proximity to coronary artery due to severity of right ventricle hypertrophy, the patient had a pulmonary valvotomy. If this patient does not make it, how do we code this procedure – we feel it should be a cath lab death and not necessarily a cardiac surgery death, due to the circumstances. **All cases are coded as they occur and this would count as an operative death.**

SeqNo: 4240

Long Name: Discharge Location

Short Name: DisLoctn

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate the location to where the patient was discharged at the Date of Hospital Discharge.

Parent Long Name: Mortality Status At Hospital Discharge

Parent Short Name: MtHospDisStat

Parent Value(s): = "Alive"

SeqNo: 4245

Long Name: VAD-Discharge Status

Short Name: VADDiscS

Database Table Name: Operations

Data Source: User



Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient had a VAD in place at discharge from the hospital.

---

SeqNo: 4250

Long Name: Date of Database Discharge

Short Name: DBDischDt

Database Table Name: Operations

Data Source: User

Format: Date - mm/dd/yyyy

Definition: Indicate the "Date of Database Discharge". The "Date of Database Discharge" is defined as a date that is determined by three rules (presented below as Rule A, Rule B, and Rule C), which specify how to complete the field "Date of Database Discharge".

[Rule A]: If a patient was admitted from their home, they must be either dead or discharged to home prior to completing the field "Date of Database Discharge". Their "Date of Database Discharge" is the date they are discharged to home or their date of mortality. If a patient was admitted from their home, the field "Date of Database Discharge" cannot be completed if the patient is transferred to another acute care facility or chronic care facility until they are either dead or discharged to home. However, if this patient survives in a chronic care facility for 6 postoperative months (i.e., 183 postoperative days in the chronic care facility), the patient can then be assigned a "Date of Database Discharge" that is the date when the patient is in the chronic care facility for 183 days. (Some institutions may not have a mechanism that allows transfer to a chronic care facility and instead utilizes their own institution as the chronic care facility. If an institution does not utilize a chronic care facility and instead keeps these chronic patients in-house, this institution can apply to this Rule [Rule A] whenever one of their patients survives for 6 postoperative months (i.e., 183 postoperative days) on "chronic care status" within their institution.)

[Rule B]: If a patient was admitted from (i.e., transferred from) a chronic care facility where they chronically reside, they must be either dead or discharged either to home or to a chronic care facility prior to completing the field "Date of Database Discharge". Their "Date of Database Discharge" is the date they are discharged either to home or to a chronic care facility, or their date of mortality.

[Rule C]: If a patient was admitted from (i.e., transferred from) another acute care facility, Rule A as previously stated applies if they lived at home prior to their admission to the transferring acute care facility. If a patient was transferred from another acute care facility, Rule B as previously stated applies if they lived in a chronic care facility prior to their admission to the transferring acute care facility.

These three rules are consistent with previously published rules defining Operative Mortality [1] and Operative Morbidity [2] in the following published manuscripts [1, 2]. [1]. Jacobs JP, Mavroudis C, Jacobs ML, Maruszewski B, Tchervenkov CI, Lacour-Gayet FG, Clarke DR, Yeh T, Walters HL 3rd, Kurosawa H, Stellin G, Ebels T, Elliott MJ.

What is Operative Mortality? Defining Death in a Surgical Registry Database: A Report from the STS Congenital Database Task Force and the Joint EACTS-STS Congenital Database Committee. The Annals of Thoracic Surgery, 81(5):1937-41, May 2006. [2]. Jacobs JP, Jacobs ML, Mavroudis C, Maruszewski B, Tchervenkov CI, Lacour-Gayet FG, Clarke DR, Yeh T, Walters HL 3rd, Kurosawa H, Stellin G, Ebels T, Elliott MJ, Vener DF, Barach P, Benavidez OJ, Bacha EA.. What is Operative Morbidity? Defining Complications in a Surgical Registry Database: A Report from the STS Congenital Database Task Force and the Joint EACTS-STS Congenital Database

---

April 2016: Patient has been a continuous patient since their first OR since 06Oct2014 and they have had 7 surgeries total with the last one being in April 2015, would I still enter the 07Apr2015 as the discharge date which would be 183 days even though they had a surgery on the 21Apr2015? How long can patients remain "open" without being submitted for harvest when they don't have a discharge date? **The 183 days is only for extended care.**

---

July 2016: We had a patient that was admitted on 07/21/15 and had a Ross-Konno done. She had a long complicated hospital stay that ended in trach/ventilator dependent status and transfer to a long term care facility on 01/06/16. She was then readmitted on 01/10/16 for 1 day and sent back to LTC 01/11/16; readmitted 01/29/16 for 1 day and sent back 01/30/16 to LTC; and finally readmitted 02/06/16 to us and she expired yesterday in our facility. Throughout all of the readmissions there were no more cardiac procedures done. I believe that her database D/C date should be 07/11/16, but my surgeons feel that her database D/C date should be at 183 days post-op and that this should not be an operative mortality. **Database discharge date is 7\_11\_2016 & it is an operative mortality**

June 2017: Patient has CPB cardiovascular procedure on 12/20/16 is discharged home on 4/28/2017, but readmitted to the hospital stepdown unit w/in 12 hours of hospital discharge.

Question 1 - Would hospital discharge date and database discharge date be 4/28? **Yes**

Question 2 - If patient has complications during readmission would these be attributed to his 12/20/16 CPB cardiovascular procedure? **No, as the complications would only be collected to the 04/28/2017 date. There would be a readmission to the hospital within 30 days.**

---

SeqNo: 4260

Long Name: Mortality Status At Database Discharge

Short Name: MtDBDisStat

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether the patient was Alive or Dead at the date and time of "Date of Database Discharge" for this operation.

---

SeqNo: 4270

Long Name: Readmission Within 30 Days

Short Name: Readmit30

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

---

**Definition:** Indicate whether the patient was readmitted within thirty days of discharge.

**Parent Long Name:** Mortality Status At Hospital Discharge

**Parent Short Name:** MtHospDisStat

**Parent Value(s):** = "Alive"

---

12/2015: If a patient is re-admitted within the 30-day period for less than 24 hours (e.g., ER, feeding device mal-positioned, cath lab procedure, etc.) and discharged home, is this coded as a re-admittance? **If the patient is actually admitted, yes; if only observation status then no. If did interventional cath lab procedure – add this clarification.**

---

**SeqNo:** 4280

**Long Name:** Readmission Date

**Short Name:** ReadmitDt

**Database Table Name:** Operations

**Data Source:** User

**Format:** Date - mm/dd/yyyy

**Definition:** Indicate the date on which the patient was readmitted.

**Parent Long Name:** Readmission Within 30 Days

**Parent Short Name:** Readmit30

**Parent Value(s):** = "Yes"

---

**SeqNo:** 4290

**Long Name:** Primary Readmission Reason

**Short Name:** ReadmitRsn

**Database Table Name:** Operations

**Data Source:** User

**Format:** Text (categorical values specified by STS)

**Definition:** Indicate the primary reason for readmission.

Whenever possible, use the most appropriate specific organ system and/or lesion based choice from the list to document the reason for admission.

Please only use one of the three choices beginning with the word "Other" when no other choice is appropriate.

If the readmission is for the patient to undergo a procedure related to the index operation (the first operation of the given hospitalization that has an Operation Type of "CPB" or "No CPB Cardiovascular"), please document the cause of this readmission to be assigned to the specific organ system and/or lesion based choice if possible. If no specific organ system and/or lesion based choice is appropriate and the readmission is for the patient to undergo a procedure related to the index operation, please choose "Other Cardiovascular Complication" if the planned procedure is cardiac, and "Other - Readmission related to this index operation" if the planned procedure is noncardiac.

---

Parent Long Name: Readmission Within 30 Days  
Parent Short Name: Readmit30  
Parent Value(s): = "Yes"

SeqNo: 4300  
Long Name: Mortality - 30-Day Status  
Short Name: Mt30Stat  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient was alive or dead on the 30th day post-surgical procedure whether in hospital or not.

---

SeqNo: 4310  
Long Name: Mortality - 30-Day Status - Method Of Verification  
Short Name: Mt30StatMeth  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the primary method used to verify the patient's 30-day mortality status.

---

SeqNo: 4330  
Long Name: Mortality - Operative Death  
Short Name: MtOpD  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Operative Mortality includes: (1) all deaths, regardless of cause, occurring during the hospitalization in which the operation was performed, even if after 30 days (including patients transferred to other acute care facilities); and (2) all deaths, regardless of cause, occurring after discharge from the hospital, but before the end of the thirtieth postoperative day.

---

12/2015: : If a patient does not have a CPB or a No CPB case, and only has an operation op type "thoracic" and the patient dies, should the Mortality Case field be marked yes or no? Based on the STS definition, it sounds like this should only be marked yes for cases that are CPB or no CPB but I am not completely clear.

**Should be marked yes for local use, but it is not analyzed. (harvested, not analyzed)**

---

February 2016: A premature neonate with a birth weight of 630gms has a PDA closure done on day of life 110 and has a weight of 3.3kg at the time of surgery. Will the child be excluded from our center's mortality calculation based on the birth weight or included in the mortality calculation based on weight at time of surgery? **Patient would be included in the mortality analysis based on the weight at the time of surgery and not birth weight.**

February 2017: We had a patient with HLHS that we performed surgery for bi-lateral PA bands. The patient was transferred to another hospital for the Stg I Norwood procedure. The patient had numerous issues that resulted in the recommendation of withdrawal of support. The patient was transferred back to our hospital for w/d of support and the patient expired. Would this mortality be coded to our hospital? **Yes it would.**

September 2017: Patient was born at our facility and had initial cardiac surgery with CPB in neonatal period. Had revision of initial repair one month later with ECMO support. Successfully weaned from ECMO. Patient was born with multiple other anomalies and required GT and trach two months after the index operation. Seven months after the index operation, patient was transferred to another acute care facility for a complex airway procedure that we do not perform in our facility. While at the other facility, patient underwent cardiac catheterization and further cardiac surgery with CPB. Patient was then transferred back to our facility two months later and then was transferred to rehab facility one month later (care of chronic trach/vent since family was not equipped to care for patient at home). Patient was readmitted to our facility four months later for airway bleeding event leading to cardiac arrest and the need for ECMO. Head CT showed findings consistent with hypoxic brain injury and support was withdrawn. Question about where/if operative mortality should be assigned. Index case was done at our facility, but patient received additional cardiac surgery outside of our facility. Death was related to the airway anomaly and occurred 15 months after the index case. Please help us code this correctly. **Patient did not D/C to home following cardiac repair from either center. Both centers have admissions with index operations thus both centers would have an operative mortality.**

September 2017: For a 4 mo infant status post Norwood who is not discharged from the hospital, and dies s/p BDG how should the Norwood mortality be coded? **The Norwood is the index operation of the admission and would be the case the mortality is attributed to.**

October 2017: A child less than 24 hours old received a pacemaker. The same patient later had an ECMO run and PA banding. The Pacemaker was the first surgery, this child ended up dying before 30 days of life. Will this mortality be excluded from our Risk Adjusted Mortality because the pacemaker was the index procedure and was inserted at less than 24 hours of life and died before 30 days of life? **The index operation is the pacemaker insertion. Because the patient is a neonate at the time of the index operation, the case would not be analyzed in the mortality analysis. The timing of the death (before or after 30 days) during of the admission does not impact the analysis or assignment of death.**

SeqNo: 4331

Long Name: Eligibility For CHSS Study

Short Name: CHSSElig

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate patient's eligibility for the Congenital Heart Surgeon Society (CHSS) study.

## 17. Patient Process Measures

SeqNo: 4340

Long Name: Patient's care discussed at preoperative multidisciplinary planning conference

Short Name: CareDiscussed

Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether this patient's care was discussed at a preoperative multidisciplinary planning conference to plan pediatric and congenital heart surgery cases.

A preoperative multidisciplinary planning conference involves attendance by multiple members of the healthcare team, with recommended participation including but not limited to: cardiology, cardiac surgery, anesthesia, and critical care.

Parent Long Name: Operation Type  
Parent Short Name: OpType  
Parent Value(s): = "CPB Cardiovascular", "No CPB Cardiovascular" or "CPB Non-Cardiovascular"

---

SeqNo: 4350  
Long Name: Reason why patient's care was not discussed  
Short Name: CareDiscussedRsn  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the reason why the patient's case was not discussed at a preoperative multidisciplinary planning conference.

Parent Long Name: Patient's care discussed at preoperative multidisciplinary planning conference  
Parent Short Name: CareDiscussed  
Parent Value(s): = "No"

---

SeqNo: 4370  
Long Name: Transesophageal Echocardiography (TEE) available for case  
Short Name: TEEAvail  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether intraoperative transesophageal echocardiography (TEE) was available for this case (or epicardial echocardiography if TEE contraindicated or not informative).

Availability is defined as the presence and availability of equipment and staff to perform the study. Reporting of compliance will be as the fraction of all Cardiac Operations with availability (as opposed to use) of TEE and/or epicardial echocardiography.

Parent Long Name: Operation Type  
Parent Short Name: OpType  
Parent Value(s): = "CPB Cardiovascular", "No CPB Cardiovascular" or "CPB Non-Cardiovascular"

---

SeqNo: 4380

Long Name: Intraoperative transesophageal echocardiography (TEE) performance  
 Short Name: TEEpicEchoPerf  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether TEE / epicardial echocardiography was performed for this case.

If available, TEE may not be performed due to surgeon preference, size of patient, not indicated, etc.

Parent Long Name: Transesophageal Echocardiography (TEE) available for case  
 Parent Short Name: TEEAvail  
 Parent Value(s): = "Yes"

SeqNo: 4400

Long Name: Preoperative antibiotic prophylaxis given  
 Short Name: PreopAntiProph  
 Database Table Name: Operations  
 Data Source: User  
 Format: Text (categorical values specified by STS)  
 Definition: Indicate whether a preoperative antibiotic prophylaxis was given to this patient.

Measure is satisfied for each Cardiac Operation, when there is documentation that the patient has received prophylactic antibiotic(s) within the hour immediately preceding surgical incision (two hours if receiving vancomycin). To satisfy this measure, the field named "Skin Incision Start Time" must be completed.

Parent Long Name: Operation Type  
 Parent Short Name: OpType  
 Parent Value(s): = "CPB Cardiovascular", "No CPB Cardiovascular" or "CPB Non-Cardiovascular"

March 2017: We only have "Yes" or "No" as options for prophylactic antibiotics within one hour (two if Vanc), but when we perform a re-operation or the patient is already on a scheduled antibiotic, is it counted against us in any way if we answer "No"? If they are on a scheduled antibiotic, should I go ahead and put the time of the last dose in seq 4700 PreopAntiProphTime even though it's more than an hour before skin incisions? **Answer 'no' and do not include the time of the antibiotic if it is outside of the 1 hour time before skin incision.**

SeqNo: 4410

Long Name: Preoperative antibiotic prophylaxis - Cephalosporin  
 Short Name: PreopAntiProphCeph  
 Database Table Name: Operations

Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the preoperative antibiotic prophylaxis included Cephalosporin.

Parent Long Name: Preoperative antibiotic prophylaxis given  
Parent Short Name: PreopAntiProph  
Parent Value(s): = "Yes"

---

SeqNo: 4420

Long Name: Preoperative antibiotic prophylaxis - Penicillin or related medication  
Short Name: PreopAntiProphPen  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the preoperative antibiotic prophylaxis included penicillin or related medications (i.e., Oxacillin, Nafcillin, Ampicillin, etc.)

Parent Long Name: Preoperative antibiotic prophylaxis given  
Parent Short Name: PreopAntiProph  
Parent Value(s): = "Yes"

---

SeqNo: 4430

Long Name: Preoperative antibiotic prophylaxis - Aminoglycoside  
Short Name: PreopAntiProphAmino  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the preoperative antibiotic prophylaxis included Aminoglycoside.

Parent Long Name: Preoperative antibiotic prophylaxis given  
Parent Short Name: PreopAntiProph  
Parent Value(s): = "Yes"

---

SeqNo: 4440

Long Name: Preoperative antibiotic prophylaxis - Vancomycin  
Short Name: PreopAntiProphVan  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the preoperative antibiotic prophylaxis included Vancomycin.

Parent Long Name: Preoperative antibiotic prophylaxis given

---



Parent Short Name: PreopAntiProph  
Parent Value(s): = "Yes"

---

SeqNo: 4450  
Long Name: Preoperative antibiotic prophylaxis - Other  
Short Name: PreopAntiProphOth  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the preoperative antibiotic prophylaxis included any other class of antibiotic.

Parent Long Name: Preoperative antibiotic prophylaxis given  
Parent Short Name: PreopAntiProph  
Parent Value(s): = "Yes"

---

SeqNo: 4470  
Long Name: Preoperative antibiotic prophylaxis - Time started  
Short Name: PreopAntiProphTime  
Database Table Name: Operations  
Data Source: User  
Format: Time - hh:mm (24-hour clock)  
Definition: Indicate the time when the antibiotic infusion started.

Parent Long Name: Preoperative antibiotic prophylaxis given  
Parent Short Name: PreopAntiProph  
Parent Value(s): = "Yes"

---

SeqNo: 4480  
Long Name: Conventional preprocedure time-out.  
Short Name: ConvTimeOut  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a conventional preprocedural "time-out", which includes identification of patient, operative site, procedure, and history of any allergies, was performed.

Parent Long Name: Operation Type  
Parent Short Name: OpType  
Parent Value(s): = "CPB Cardiovascular", "No CPB Cardiovascular" or "CPB Non-Cardiovascular"

---

SeqNo: 4490  
Long Name: Surgeon shares essential elements of operative plan  
Short Name: PreProcBrief  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a pre-procedural briefing was performed wherein the surgeon shares with all members of the operating room team the essential elements of the operative plan; including diagnosis, planned procedure, outline of essentials of anesthesia and bypass strategies, antibiotic prophylaxis, availability of blood products, anticipated or planned implants or device applications, and anticipated challenges.

Parent Long Name: Operation Type  
Parent Short Name: OpType  
Parent Value(s): = "CPB Cardiovascular", "No CPB Cardiovascular" or "CPB Non-Cardiovascular"

---

SeqNo: 4500  
Long Name: Postprocedure debriefing  
Short Name: PostProcDebrief  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a postprocedural debriefing was performed wherein the surgeon succinctly reviews with all members of the operating room team the essential elements of the operative plan, identifying both the successful components and the opportunities for improvement. This debriefing should take place prior to the patient leaving the operating room or its equivalent, and may be followed by a more in-depth dialogue involving team members at a later time. (The actual debriefing in the operating room is intentionally and importantly brief, in recognition of the fact that periods of transition may be times of instability or vulnerability for the patient.)

Parent Long Name: Operation Type  
Parent Short Name: OpType  
Parent Value(s): = "CPB Cardiovascular", "No CPB Cardiovascular" or "CPB Non-Cardiovascular"

---

SeqNo: 4510  
Long Name: Hand-off protocol at the time of transfer to the Intensive Care Unit  
Short Name: HandoffProtocol  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a briefing and execution of a hand-off protocol (checklist) was performed at the time of transfer (arrival) to the Intensive Care Unit at the end of the

operation, involving ALL of the following: the anesthesiologist, surgeon, physician staff of the Intensive Care Unit (including critical care and cardiology) and nursing.

Parent Long Name: Operation Type  
Parent Short Name: OpType  
Parent Value(s): = "CPB Cardiovascular", "No CPB Cardiovascular" or "CPB Non-Cardiovascular"

---

SeqNo: 4520  
Long Name: Hand-off protocol - Anesthesiologist  
Short Name: HandoffAnesth  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the anesthesiologist or designee attended the hand-off protocol at the time of transfer to the Intensive Care Unit at the end of the operation.

Parent Long Name: Hand-off protocol at the time of transfer to the Intensive Care Unit  
Parent Short Name: HandoffProtocol  
Parent Value(s): = "Yes - Not all required team members present"

---

SeqNo: 4530  
Long Name: Hand-off protocol - Surgeon  
Short Name: HandoffSurg  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the surgeon or designee attended the hand-off protocol at the time of transfer to the Intensive Care Unit at the end of the operation.

Parent Long Name: Hand-off protocol at the time of transfer to the Intensive Care Unit  
Parent Short Name: HandoffProtocol  
Parent Value(s): = "Yes - Not all required team members present"

---

SeqNo: 4540  
Long Name: Hand-off protocol - Physician staff of the Intensive Care Unit  
Short Name: HandoffPhysStaff  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the physician staff of the Intensive Care Unit or designee attended the

hand-off protocol at the time of transfer to the Intensive Care Unit at the end of the operation.

Parent Long Name: Hand-off protocol at the time of transfer to the Intensive Care Unit  
Parent Short Name: HandoffProtocol  
Parent Value(s): = "Yes - Not all required team members present"

---

SeqNo: 4550

Long Name: Hand-off protocol - Nursing  
Short Name: HandoffNursing  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a nurse or designee attended the hand-off protocol at the time of transfer to the Intensive Care Unit at the end of the operation.

Parent Long Name: Hand-off protocol at the time of transfer to the Intensive Care Unit  
Parent Short Name: HandoffProtocol  
Parent Value(s): = "Yes - Not all required team members present"

---

SeqNo: 4560

Long Name: Patient died or had major postoperative complication(s)  
Short Name: PostOpComp  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient died before hospital discharge and/or had any of these major postoperative complication(s):  
a. New postoperative renal failure requiring dialysis  
b. New postoperative neurological deficit persisting at discharge  
c. Arrhythmia necessitating permanent pacemaker insertion  
d. Paralyzed diaphragm  
e. Need for postoperative mechanical circulatory support  
f. Unplanned reoperation and/or interventional cardiovascular catheterization procedure

The detailed definitions for the six postoperative complications are the definitions used in the current version of the STS Congenital Heart Surgery Database.

These detailed definitions for these six postoperative complications may be found in the following manuscript: Jacobs JP et al. Quality measures for congenital and pediatric cardiac surgery. World Journal for Pediatric and Congenital Heart Surgery 2012;3:32-47

---

SeqNo: 4570

Long Name: Patient management and outcomes reviewed  
Short Name: PostOpReview  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient's management and outcomes were reviewed as a part of a regularly scheduled Quality Assurance and Quality Improvement Cardiac Care Conference (i.e., Morbidity and Mortality conference).

Parent Long Name: Patient died or had major postoperative complication(s)  
Parent Short Name: PostOpComp  
Parent Value(s): = "Yes"

---

SeqNo: 4580

Long Name: Patient management and outcomes reviewed - date  
Short Name: PostOpReviewDate  
Database Table Name: Operations  
Data Source: User  
Format: Date - mm/dd/yyyy  
Definition: Indicate the date this patient's management and outcome was reviewed as a part of a regularly scheduled Quality Assurance and Quality Improvement Cardiac Care Conference (i.e., Morbidity and Mortality conference).

Parent Long Name: Patient management and outcomes reviewed  
Parent Short Name: PostOpReview  
Parent Value(s): = "Reviewed at conference"

---

## 18. Anesthesia Administrative

SeqNo: 4581

Long Name: Anesthesiologist Present  
Short Name: AnesPresent  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether an anesthesiologist was present for the procedure.

---

SeqNo: 4590

Long Name: Primary Anesthesiologist Attending Name  
Short Name: PrimAnesName

---

Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by user)  
Definition: Indicate the name of the primary anesthesiologist (attending physician present at induction of anesthesia). The name, NPI and signature of all anesthesiologists contributing data to the database must be on file with the STS for data files to be accepted.

Parent Long Name: Anesthesiologist Present  
Parent Short Name: AnesPresent  
Parent Value(s): = "Yes"

---

SeqNo: 4600  
Long Name: Primary Anesthesiologist National Provider Identifier  
Short Name: PrimAnesNPI  
Database Table Name: Operations  
Data Source: Lookup  
Format: Text  
Definition: Indicate the individual-level National Provider Identifier (NPI) of the anesthesiologist performing the procedure.

Parent Long Name: Anesthesiologist Present  
Parent Short Name: AnesPresent  
Parent Value(s): = "Yes"

---

SeqNo: 4610  
Long Name: Secondary Anesthesiologist Attending  
Short Name: SecAnes  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a relieving anesthesiologist and/or second anesthesiology attending was present during this procedure.

Parent Long Name: Anesthesiologist Present  
Parent Short Name: AnesPresent  
Parent Value(s): = "Yes"

---

SeqNo: 4630  
Long Name: Fellow or Resident Present  
Short Name: FelRes  
Database Table Name: Operations

---

Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a Fellow or Resident was present during this procedure.

---

SeqNo: 4640  
Long Name: Mid-Level Provider (CRNA, AA) Present  
Short Name: CRNA  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a Certified Registered Nurse Anesthetist (CRNA) or Anesthesia Assistant (AA) participated in the patient care during all or part of this procedure.

---

## 19. Anesthesia Preoperative

SeqNo: 4670  
Long Name: Preoperative Medications Table Unique Record Identifier  
Short Name: PMUniqueID  
Database Table Name: PreopMeds  
Data Source: Automatic  
Format: Text  
Definition: Unique identifier for the record in the Preoperative Medications table.

---

SeqNo: 4680  
Long Name: Preoperative Medication Link to Operations Table  
Short Name: OperationID  
Database Table Name: PreopMeds  
Data Source: Automatic  
Format: Text  
Definition: An arbitrary, unique value generated by the software that permanently identifies each operation record in the participant's database. This field is the foreign key that links the Preoperative Medications record with the associated record in the Operations table.

SeqNo: 4700  
Long Name: Preoperative Medication Category  
Short Name: PreopMedCat  
Database Table Name: PreopMeds  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the categories of preoperative medication(s) given to the patient within **24** hours (unless noted otherwise) prior to the period of anesthetic care.

---

April 2015: clarification of time frame for preoperative medications. Time frame updated to 24 hours except for the exceptions listed.

---

SeqNo: 4710

Long Name: Preoperative Sedation

Short Name: PreopSed

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether the patient received preoperative sedation.

---

SeqNo: 4720

Long Name: Preoperative Sedation Route

Short Name: PreopSedRte

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate the route used for preoperative sedation.

Parent Long Name: Preoperative Sedation

Parent Short Name: PreopSed

Parent Value(s): = "Yes"

---

SeqNo: 4730

Long Name: Preoperative Sedation Drug - Atropine

Short Name: PreopSedDrugAtro

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether the patient received Atropine for preoperative sedation.

Parent Long Name: Preoperative Sedation

Parent Short Name: PreopSed

Parent Value(s): = "Yes"

---

SeqNo: 4740

Long Name: Preoperative Sedation Drug - Demerol

Short Name: PreopSedDrugDem

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

---



**Definition:** Indicate whether the patient received Demerol for preoperative sedation.

**Parent Long Name:** Preoperative Sedation

**Parent Short Name:** PreopSed

**Parent Value(s):** = "Yes"

---

**SeqNo:** 4741

**Long Name:** Preoperative Sedation Drug - Dexmedetomidine

**Short Name:** PreopSedDrugDex

**Database Table Name:** Operations

**Data Source:** User

**Format:** Text (categorical values specified by STS)

**Definition:** Indicate whether the patient received Dexmedetomidine for preoperative sedation.

**Parent Long Name:** Preoperative Sedation

**Parent Short Name:** PreopSed

**Parent Value(s):** = "Yes"

---

**SeqNo:** 4750

**Long Name:** Preoperative Sedation Drug - Diazepam

**Short Name:** PreopSedDrugDiaz

**Database Table Name:** Operations

**Data Source:** User

**Format:** Text (categorical values specified by STS)

**Definition:** Indicate whether the patient received Diazepam for preoperative sedation.

**Parent Long Name:** Preoperative Sedation

**Parent Short Name:** PreopSed

**Parent Value(s):** = "Yes"

---

**SeqNo:** 4751

**Long Name:** Preoperative Sedation Drug - Fentanyl

**Short Name:** PreopSedDrugFent

**Database Table Name:** Operations

**Data Source:** User

**Format:** Text (categorical values specified by STS)

**Definition:** Indicate whether the patient received Fentanyl for preoperative sedation.

**Parent Long Name:** Preoperative Sedation

**Parent Short Name:** PreopSed

**Parent Value(s):** = "Yes"

---

---

SeqNo: 4760  
Long Name: Preoperative Sedation Drug - Glycopyrrolate  
Short Name: PreopSedDrugGlyco  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient received Glycopyrrolate for preoperative sedation.

Parent Long Name: Preoperative Sedation  
Parent Short Name: PreopSed  
Parent Value(s): = "Yes"

---

SeqNo: 4770  
Long Name: Preoperative Sedation Drug - Ketamine  
Short Name: PreopSedDrugKet  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient received Ketamine for preoperative sedation.

Parent Long Name: Preoperative Sedation  
Parent Short Name: PreopSed  
Parent Value(s): = "Yes"

---

SeqNo: 4780  
Long Name: Preoperative Sedation Drug - Lorazepam  
Short Name: PreopSedDrugLoraz  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient received Lorazepam for preoperative sedation.

Parent Long Name: Preoperative Sedation  
Parent Short Name: PreopSed  
Parent Value(s): = "Yes"

SeqNo: 4790  
Long Name: Preoperative Sedation Drug - Midazolam  
Short Name: PreopSedDrugMidaz  
Database Table Name: Operations  
Data Source: User

---

Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient received Midazolam for preoperative sedation.

Parent Long Name: Preoperative Sedation  
Parent Short Name: PreopSed  
Parent Value(s): = "Yes"

---

SeqNo: 4800  
Long Name: Preoperative Sedation Drug - Morphine  
Short Name: PreopSedDrugMorph  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient received Morphine for preoperative sedation.

Parent Long Name: Preoperative Sedation  
Parent Short Name: PreopSed  
Parent Value(s): = "Yes"

---

SeqNo: 4810  
Long Name: Preoperative Sedation Drug - Pentobarbital  
Short Name: PreopSedDrugPent  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient received Pentobarbital for preoperative sedation.

Parent Long Name: Preoperative Sedation  
Parent Short Name: PreopSed  
Parent Value(s): = "Yes"

---

SeqNo: 4820  
Long Name: Preoperative Baseline Oxygen Saturation  
Short Name: PreopO2Sat  
Database Table Name: Operations  
Data Source: User  
Format: Real  
Definition: Indicate the preoperative resting pulse oximeter saturation (%) recorded either in the clinic or immediately prior to the procedure.

---

Low Value: 1.0 High Value: 100.0

---

SeqNo: 4830

Long Name: Preoperative Oxygen Supplementation

Short Name: PreopOxygen

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether the patient received preoperative oxygen supplementation.

---

SeqNo: 4840

Long Name: Transport to Procedure Location Date and Time

Short Name: PLocTransDT

Database Table Name: Operations

Data Source: User

Format: Date/Time - mm/dd/yyyy hh:mm

Definition: Indicate the date (mm/dd/yyyy) and time (hh:mm 24-hour clock) of day when the patient was transferred to the procedure location or when anesthesia started.

---

## 20. Anesthesia Monitoring

SeqNo: 4850

Long Name: Arterial Line

Short Name: ArtLine

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether an arterial line was used during this procedure.

---

SeqNo: 4860

Long Name: Arterial Line Type - Radial

Short Name: ArtLineTypeRad

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether a radial arterial line type was used during this procedure.

Parent Long Name: Arterial Line

Parent Short Name: ArtLine

---

Parent Value(s): = "Yes"

---

SeqNo: 4870

Long Name: Arterial Line Type - Brachial

Short Name: ArtLineTypeBrach

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether a brachial arterial line type was used during this procedure.

Parent Long Name: Arterial Line

Parent Short Name: ArtLine

Parent Value(s): = "Yes"

---

SeqNo: 4880

Long Name: Arterial Line Type - Axillary

Short Name: ArtLineTypeAx

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether an axillary arterial line type was used during this procedure.

Parent Long Name: Arterial Line

Parent Short Name: ArtLine

Parent Value(s): = "Yes"

---

SeqNo: 4890

Long Name: Arterial Line Type - Femoral

Short Name: ArtLineTypeFem

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether a femoral arterial line type was used during this procedure.

Parent Long Name: Arterial Line

Parent Short Name: ArtLine

Parent Value(s): = "Yes"

---

SeqNo: 4900

Long Name: Arterial Line Type - Ulnar

---

Short Name: ArtLineTypeUlnar  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether an ulnar arterial line type was used during this procedure.

Parent Long Name: Arterial Line  
Parent Short Name: ArtLine  
Parent Value(s): = "Yes"

---

SeqNo: 4910  
Long Name: Arterial Line Type - Dorsalis Pedis  
Short Name: ArtLineTypeDors  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a dorsalis pedis arterial line type was used during this procedure.

Parent Long Name: Arterial Line  
Parent Short Name: ArtLine  
Parent Value(s): = "Yes"

---

SeqNo: 4920  
Long Name: Arterial Line Type - Posterior Tibial  
Short Name: ArtLineTypePost  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a posterior tibial arterial line type was used during this procedure.

Parent Long Name: Arterial Line  
Parent Short Name: ArtLine  
Parent Value(s): = "Yes"

---

SeqNo: 4930  
Long Name: Arterial Line Type - Umbilical  
Short Name: ArtLineTypeCent  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether an umbilical arterial line type was used during this procedure.

---

Parent Long Name: Arterial Line  
Parent Short Name: ArtLine  
Parent Value(s): = "Yes"

---

SeqNo: 4931  
Long Name: Arterial Line In-Situ Pre-Procedure  
Short Name: ArtLinePreProc  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the arterial line was in-situ pre-procedure.

Parent Long Name: Arterial Line  
Parent Short Name: ArtLine  
Parent Value(s): = "Yes"

---

SeqNo: 4940  
Long Name: Cutdown  
Short Name: Cutdown  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a cutdown was used during this procedure.

---

SeqNo: 4950  
Long Name: Cutdown Type - Radial  
Short Name: CutdownRad  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a radial cutdown was used.

Parent Long Name: Cutdown  
Parent Short Name: Cutdown  
Parent Value(s): = "Yes"

SeqNo: 4960  
Long Name: Cutdown Type - Femoral  
Short Name: CutdownFem  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)

---

Definition: Indicate whether a femoral cutdown was used.

Parent Long Name: Cutdown

Parent Short Name: Cutdown

Parent Value(s): = "Yes"

---

SeqNo: 4970

Long Name: Cutdown Type - Ulnar

Short Name: CutdownUln

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether an ulnar cutdown was used.

Parent Long Name: Cutdown

Parent Short Name: Cutdown

Parent Value(s): = "Yes"

---

SeqNo: 4980

Long Name: Cutdown Type - Other

Short Name: CutdownOth

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether any other type of cutdown was used.

Parent Long Name: Cutdown

Parent Short Name: Cutdown

Parent Value(s): = "Yes"

---

SeqNo: 4990

Long Name: Percutaneous Central Pressure

Short Name: PercCentPress

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether percutaneous central pressure was used during this procedure.

---

SeqNo: 5000

Long Name: Percutaneous Central Pressure Location - Right Internal Jugular

Short Name: PCPLocRJug

Database Table Name: Operations

---



Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the percutaneous central pressure was located in the right internal jugular.

Parent Long Name: Percutaneous Central Pressure  
Parent Short Name: PercCentPress  
Parent Value(s): = "Yes"

---

SeqNo: 5010  
Long Name: Percutaneous Central Pressure Location - Left Internal Jugular  
Short Name: PCPLocLJug  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the percutaneous central pressure was used in the left internal jugular.

Parent Long Name: Percutaneous Central Pressure  
Parent Short Name: PercCentPress  
Parent Value(s): = "Yes"

---

SeqNo: 5020  
Long Name: Percutaneous Central Pressure Location - Right Subclavian  
Short Name: PCPLocRSub  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the percutaneous central pressure was used in the right subclavian.

Parent Long Name: Percutaneous Central Pressure  
Parent Short Name: PercCentPress  
Parent Value(s): = "Yes"

---

SeqNo: 5030  
Long Name: Percutaneous Central Pressure Location - Left Subclavian  
Short Name: PCPLocLSub  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the percutaneous central pressure was used in the left subclavian.

Parent Long Name: Percutaneous Central Pressure  
Parent Short Name: PercCentPress

---

Parent Value(s): = "Yes"

---

SeqNo: 5040

Long Name: Percutaneous Central Pressure Location - Right Femoral Vein

Short Name: PCPLocRFem

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether the percutaneous central pressure was used in the right femoral vein.

Parent Long Name: Percutaneous Central Pressure

Parent Short Name: PercCentPress

Parent Value(s): = "Yes"

---

SeqNo: 5050

Long Name: Percutaneous Central Pressure Location - Left Femoral Vein

Short Name: PCPLocLFem

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether the percutaneous central pressure was used in the left femoral vein.

Parent Long Name: Percutaneous Central Pressure

Parent Short Name: PercCentPress

Parent Value(s): = "Yes"

---

SeqNo: 5051

Long Name: Percutaneous Central Pressure Location - PICC

Short Name: PCPLocPICC

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether the percutaneous central pressure was used in the PICC.

Parent Long Name: Percutaneous Central Pressure

Parent Short Name: PercCentPress

Parent Value(s): = "Yes"

---

SeqNo: 5060

Long Name: Percutaneous Central Pressure Location - Other

Short Name: PCPLocOth

Database Table Name: Operations

---

Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the percutaneous central pressure was used in any other location.

Parent Long Name: Percutaneous Central Pressure  
Parent Short Name: PercCentPress  
Parent Value(s): = "Yes"

---

SeqNo: 5062  
Long Name: CVP, PICC, LA or RA Line(s) In-Situ Pre-Procedure  
Short Name: CVPPICCPreProc  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether CVP, PICC, LA or RA line(s) were in place prior to entering the OR.

Parent Long Name: Percutaneous Central Pressure  
Parent Short Name: PercCentPress  
Parent Value(s): = "Yes"

---

SeqNo: 5070  
Long Name: CVP Placed By Anesthesia  
Short Name: CVPPlaced  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a CVP was placed by anesthesia during this procedure.

---

SeqNo: 5071  
Long Name: Surgeon Placed Lines INSTEAD of Anesthesia Placed Central Lines  
Short Name: SurgMonLines  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the surgeon placed one or more central monitoring / medication lines directly in the Right, Left or Common Atria during the procedure INSTEAD of pre-incision placement of a central line by anesthesia or the use of existing percutaneous CVL or PICC. This does not include monitoring lines placed during the procedure in addition to the anesthesia or in-situ catheters.

---

SeqNo: 5080

---

Long Name: Swan-Ganz Catheter  
Short Name: SGCath  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a Swan-Ganz catheter was inserted or utilized by anesthesia during this procedure.

---

SeqNo: 5090  
Long Name: Oximetric Central Line  
Short Name: ScVO2  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether an oximetric central line was inserted or utilized by anesthesia during this procedure.

---

SeqNo: 5100  
Long Name: Ultrasound Guidance Used For Line Placement  
Short Name: UltraGuide  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether real-time ultrasound imaging was used for line placement (i.e., Sonosite or equivalent).

---

SeqNo: 5110  
Long Name: Neurologic Monitoring  
Short Name: NeuroMonitor  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient received neurologic monitoring during this procedure.

---

SeqNo: 5130  
Long Name: Neurologic Monitoring - Bispectral Index  
Short Name: NeuroMonBIS  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the neurologic monitoring performed during this procedure included

Bispectral Index (BIS).

Parent Long Name: Neurologic Monitoring  
Parent Short Name: NeuroMonitor  
Parent Value(s): = "Yes"

---

SeqNo: 5140  
Long Name: Neurologic Monitoring - Transcranial Doppler  
Short Name: NeuroMonTCD  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the neurologic monitoring performed during this procedure included Transcranial Doppler (TCD).

Parent Long Name: Neurologic Monitoring  
Parent Short Name: NeuroMonitor  
Parent Value(s): = "Yes"

---

SeqNo: 5141  
Long Name: Neurologic Monitoring - NIRS (Cerebral)  
Short Name: NeuroMonNIRS  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the neurologic (cerebral) monitoring performed during this procedure included Near Infrared Spectroscopy (NIRS).

Parent Long Name: Neurologic Monitoring  
Parent Short Name: NeuroMonitor  
Parent Value(s): = "Yes"

---

SeqNo: 5150  
Long Name: Neurologic Monitoring - Other  
Short Name: NeuroMonOth  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the neurologic monitoring performed during this procedure included some other method.

Parent Long Name: Neurologic Monitoring

---

Parent Short Name: NeuroMonitor  
Parent Value(s): = "Yes"

---

SeqNo: 5160  
Long Name: Lowest Recorded Intraoperative Temperature  
Short Name: LowIntraopTemp  
Database Table Name: Operations  
Data Source: User  
Format: Real  
Definition: Indicate the patient's lowest temperature (in degrees Centigrade) recorded during the intraoperative period.

Low Value: 0.1      High Value: 40.9

---

SeqNo: 5170  
Long Name: Lowest Intraoperative Temperature Monitoring Site  
Short Name: IntraopTempSite  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the site where the patient's lowest temperature was being recorded intraoperatively.

---

SeqNo: 5180  
Long Name: Transesophageal Echocardiography  
Short Name: TEE  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a transesophageal echocardiography probe was placed or attempted during this procedure.

---

## **21. Anesthesia Anesthetic Technique**

SeqNo: 5190  
Long Name: Induction Date and Time  
Short Name: InductionDT  
Database Table Name: Operations

---

Data Source: User  
Format: Date/Time - mm/dd/yyyy hh:mm  
Definition: Indicate the date (mm/dd/yyyy) and time (hh:mm 24-hour clock) of day when the patient was first induced.

---

SeqNo: 5200  
Long Name: Induction Type - Inhalation  
Short Name: IndTypeInh  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether an inhalation drug was used as an induction agent.

---

SeqNo: 5220  
Long Name: Induction Agent - Inhalation - Sevoflurane  
Short Name: IndAgentInhalSevo  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether sevoflurane was used for induction of anesthesia.

Parent Long Name: Induction Type - Inhalation  
Parent Short Name: IndTypeInh  
Parent Value(s): = "Yes"

SeqNo: 5230  
Long Name: Induction Agent - Inhalation - Isoflurane  
Short Name: IndAgentInhalIso  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether isoflurane was used for induction of anesthesia.

Parent Long Name: Induction Type - Inhalation  
Parent Short Name: IndTypeInh  
Parent Value(s): = "Yes"

---

SeqNo: 5240  
Long Name: Induction Type - Intravenous  
Short Name: IndTypeIV  
Database Table Name: Operations  
Data Source: User

---

Format: Text (categorical values specified by STS)  
Definition: Indicate whether an intravenous drug was used as an induction agent.

---

SeqNo: 5260  
Long Name: Induction Agent - Intravenous - Sodium Thiopental  
Short Name: IndAgentIVSodT  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether sodium thiopental was used for induction of anesthesia.  
  
Parent Long Name: Induction Type - Intravenous  
Parent Short Name: IndTypeIV  
Parent Value(s): = "Yes"

---

SeqNo: 5270  
Long Name: Induction Agent - Intravenous - Ketamine  
Short Name: IndAgentIVKet  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether ketamine was used for induction of anesthesia.  
  
Parent Long Name: Induction Type - Intravenous  
Parent Short Name: IndTypeIV  
Parent Value(s): = "Yes"

---

SeqNo: 5280  
Long Name: Induction Agent - Intravenous - Etomidate  
Short Name: IndAgentIVEtom  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether etomidate was used for induction of anesthesia.  
  
Parent Long Name: Induction Type - Intravenous  
Parent Short Name: IndTypeIV  
Parent Value(s): = "Yes"

---

SeqNo: 5290  
Long Name: Induction Agent - Intravenous - Propofol

---



Short Name: IndAgentIVProp  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether propofol was used for induction of anesthesia.

Parent Long Name: Induction Type - Intravenous  
Parent Short Name: IndTypeIV  
Parent Value(s): = "Yes"

---

SeqNo: 5300  
Long Name: Induction Agent - Intravenous - Fentanyl  
Short Name: IndAgentIVFent  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether fentanyl was used for induction of anesthesia.

Parent Long Name: Induction Type - Intravenous  
Parent Short Name: IndTypeIV  
Parent Value(s): = "Yes"

---

SeqNo: 5310  
Long Name: Induction Agent - Intravenous - Midazolam  
Short Name: IndAgentIVMid  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether intravenous midazolam was used for induction of anesthesia.

Parent Long Name: Induction Type - Intravenous  
Parent Short Name: IndTypeIV  
Parent Value(s): = "Yes"

---

SeqNo: 5320  
Long Name: Induction Agent - Intravenous - Dexmedetomidine  
Short Name: IndAgentIVDex  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether dexmedetomidine was used for induction of anesthesia.

---

Parent Long Name: Induction Type - Intravenous  
Parent Short Name: IndTypeIV  
Parent Value(s): = "Yes"

---

SeqNo: 5330  
Long Name: Induction Agent - Intravenous - Sufentanil  
Short Name: IndAgentIVSuf  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether sufentanil was used for induction of anesthesia.

Parent Long Name: Induction Type - Intravenous  
Parent Short Name: IndTypeIV  
Parent Value(s): = "Yes"

---

SeqNo: 5340  
Long Name: Induction Agent - Intravenous - Remifentanil  
Short Name: IndAgentIVRem  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether remifentanil was used for induction of anesthesia.

Parent Long Name: Induction Type - Intravenous  
Parent Short Name: IndTypeIV  
Parent Value(s): = "Yes"

---

SeqNo: 5350  
Long Name: Induction Type - Intramuscular  
Short Name: IndTypeIM  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether an intramuscular drug was used as an induction agent.

---

SeqNo: 5370  
Long Name: Induction Agent - Intramuscular - Ketamine  
Short Name: IndAgentIMKet  
Database Table Name: Operations  
Data Source: User

---

Format: Text (categorical values specified by STS)  
Definition: Indicate whether ketamine was used for induction of anesthesia.

Parent Long Name: Induction Type - Intramuscular  
Parent Short Name: IndTypeIM  
Parent Value(s): = "Yes"

---

SeqNo: 5380  
Long Name: Induction Agent - Intramuscular - Midazolam  
Short Name: IndAgentIMMid  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether intramuscular midazolam was used for induction of anesthesia.

Parent Long Name: Induction Type - Intramuscular  
Parent Short Name: IndTypeIM  
Parent Value(s): = "Yes"

---

SeqNo: 5400  
Long Name: Regional Anesthetic  
Short Name: RegionalAnes  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a regional anesthetic was used during this operation.

---

SeqNo: 5410  
Long Name: Regional Anesthetic Site  
Short Name: RegAnesSite  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the technique used for the regional anesthetic.

Parent Long Name: Regional Anesthetic  
Parent Short Name: RegionalAnes  
Parent Value(s): = "Yes"

---

SeqNo: 5420  
Long Name: Regional Anesthetic Drug - Bupivacaine

---

Short Name: RegAnesDrugBup  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the regional anesthetic drug Bupivacaine was used during this procedure.

Parent Long Name: Regional Anesthetic  
Parent Short Name: RegionalAnes  
Parent Value(s): = "Yes"

---

SeqNo: 5430  
Long Name: Regional Anesthetic Drug - Bupivacaine/Fentanyl  
Short Name: RegAnesDrugBupFen  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the regional anesthetic drug Bupivacaine/Fentanyl was used during this procedure.  
Parent Long Name: Regional Anesthetic  
Parent Short Name: RegionalAnes  
Parent Value(s): = "Yes"

SeqNo: 5440  
Long Name: Regional Anesthetic Drug - Clonidine  
Short Name: RegAnesDrugClon  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the regional anesthetic drug Clonidine was used during this procedure.  
Parent Long Name: Regional Anesthetic  
Parent Short Name: RegionalAnes  
Parent Value(s): = "Yes"

---

SeqNo: 5450  
Long Name: Regional Anesthetic Drug - Fentanyl  
Short Name: RegAnesDrugFen  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the regional anesthetic drug Fentanyl was used during this procedure.

---

Parent Long Name: Regional Anesthetic  
Parent Short Name: RegionalAnes  
Parent Value(s): = "Yes"

---

SeqNo: 5460  
Long Name: Regional Anesthetic Drug - Hydromorphone  
Short Name: RegAnesDrugHydro  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the regional anesthetic drug Hydromorphone was used during this procedure.  
Parent Long Name: Regional Anesthetic  
Parent Short Name: RegionalAnes  
Parent Value(s): = "Yes"

---

SeqNo: 5470  
Long Name: Regional Anesthetic Drug - Lidocaine  
Short Name: RegAnesDrugLido  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the regional anesthetic drug Lidocaine was used during this procedure.  
Parent Long Name: Regional Anesthetic  
Parent Short Name: RegionalAnes  
Parent Value(s): = "Yes"

---

SeqNo: 5480  
Long Name: Regional Anesthetic Drug - Morphine  
Short Name: RegAnesDrugMorph  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the regional anesthetic drug Morphine was used during this procedure.  
Parent Long Name: Regional Anesthetic  
Parent Short Name: RegionalAnes  
Parent Value(s): = "Yes"

---

SeqNo: 5490  
Long Name: Regional Anesthetic Drug - Ropivacaine

---

Short Name: RegAnesDrugRop  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the regional anesthetic drug Ropivacaine was used during this procedure.

Parent Long Name: Regional Anesthetic  
Parent Short Name: RegionalAnes  
Parent Value(s): = "Yes"

---

SeqNo: 5500  
Long Name: Regional Anesthetic Drug - Ropivacaine/Fentanyl  
Short Name: RegAnesDrugRopFen  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the regional anesthetic drug Ropivacaine/Fentanyl was used during this procedure.

Parent Long Name: Regional Anesthetic  
Parent Short Name: RegionalAnes  
Parent Value(s): = "Yes"

---

SeqNo: 5510  
Long Name: Regional Anesthetic Drug - Tetracaine  
Short Name: RegAnesDrugTetra  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the regional anesthetic drug Tetracaine was used during this procedure.

Parent Long Name: Regional Anesthetic  
Parent Short Name: RegionalAnes  
Parent Value(s): = "Yes"

---

SeqNo: 5520  
Long Name: Regional Anesthetic Drug - Other  
Short Name: RegAnesDrugOth  
Database Table Name: Operations  
Data Source: User

---

Format: Text (categorical values specified by STS)  
Definition: Indicate whether any other regional anesthetic drug was used during this procedure.

Parent Long Name: Regional Anesthetic  
Parent Short Name: RegionalAnes  
Parent Value(s): = "Yes"

---

SeqNo: 5530  
Long Name: Intercostal Nerve Infiltration By Surgeon or Anesthesia  
Short Name: IntNerveInf  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether intercostal nerve infiltration was performed by the surgeon or anesthesiologist.

SeqNo: 5540  
Long Name: Regional Field Block by Surgeon or Anesthesia  
Short Name: RegFieldBlock  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether a regional field block was performed by the surgeon or anesthesiologist.

---

## **22. Anesthesia Airway**

SeqNo: 5550  
Long Name: Airway In-situ (ETT or Tracheostomy)  
Short Name: AirwayInsitu  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether an Endotracheal Tube (ETT) or tracheostomy was in place prior to arrival in the procedure area.

---

SeqNo: 5551  
Long Name: ETT or Tracheostomy Replaced For Procedure  
Short Name: AirwayReplaced  
Database Table Name: Operations  
Data Source: User

---

**Format:** Text (categorical values specified by STS)  
**Definition:** Indicate whether the Endotracheal Tube or tracheostomy was electively replaced prior to the procedure. For example, oral to nasal ETT, tracheostomy to ETT, uncuffed to cuffed ETT.

**Parent Long Name:** Airway In-situ (ETT or Tracheostomy)  
**Parent Short Name:** AirwayInsitu  
**Parent Value(s):** = "Yes"

---

**SeqNo:** 5560  
**Long Name:** Airway Type  
**Short Name:** AirwayType  
**Database Table Name:** Operations  
**Data Source:** User  
**Format:** Text (categorical values specified by STS)  
**Definition:** Indicate the type of airway support that was used during this procedure.

**SeqNo:** 5570  
**Long Name:** Airway Size - Laryngeal Mask Airway  
**Short Name:** AirwaySizeLMA  
**Database Table Name:** Operations  
**Data Source:** User  
**Format:** Text (categorical values specified by STS)  
**Definition:** Indicate the size of the laryngeal mask airway used during this operation

**Parent Long Name:** Airway Type  
**Parent Short Name:** AirwayType  
**Parent Value(s):** = "Laryngeal Mask Airway (LMA)"

---

**SeqNo:** 5580  
**Long Name:** Airway Size - Endotracheal Intubation  
**Short Name:** AirwaySizeIntub  
**Database Table Name:** Operations  
**Data Source:** User  
**Format:** Text (categorical values specified by STS)  
**Definition:** Indicate the size of the endotracheal intubation airway used during this procedure. Measurement should be the inner diameter (ID) size measured in millimeters (mm).

**Parent Long Name:** Airway Type  
**Parent Short Name:** AirwayType  
**Parent Value(s):** = "Endotracheal intubation"

---

**SeqNo:** 5590

---



Long Name: Cuffed  
Short Name: Cuffed  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the endotracheal tube was cuffed.

Parent Long Name: Airway Type  
Parent Short Name: AirwayType  
Parent Value(s): = "Endotracheal intubation"

---

SeqNo: 5600  
Long Name: Airway Site  
Short Name: AirwaySite  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the endotracheal intubation site.

Parent Long Name: Airway Type  
Parent Short Name: AirwayType  
Parent Value(s): = "Endotracheal intubation" or "Tracheostomy"

---

SeqNo: 5610  
Long Name: Endobronchial Isolation (DLETT, Bronchial Blocker)  
Short Name: Endobronclso  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether endobronchial isolation was employed using a double lumen ETT or bronchial blocker.

---

SeqNo: 5611  
Long Name: Endobronchial Isolation Method  
Short Name: EndobronclsoMeth  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the method used to isolate lung.

Parent Long Name: Endobronchial Isolation (DLETT, Bronchial Blocker)  
Parent Short Name: Endobronclso

---

Parent Value(s): = "Yes"

---

SeqNo: 5620

Long Name: ICU-Type Ventilator Used Intraop

Short Name: ICUTypeVent

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether an ICU-type ventilator was used during the procedure.

---

SeqNo: 5621

Long Name: Anesthesia Ready / End of Induction

Short Name: EndOfInductDT

Database Table Name: Operations

Data Source: User

Format: Date/Time - mm/dd/yyyy hh:mm

Definition: Indicate the date and time at which anesthesia preparations for surgery, such as placement of desired airway and vascular access, have been completed.

---

## 24. Anesthesia Intraoperative Pharmacology

SeqNo: 6120

Long Name: Intraoperative Pharmacology Table Unique Record Identifier

Short Name: IPUniqueID

Database Table Name: IntraopPharm

Data Source: Automatic

Format: Text

Definition: Unique identifier for the record in the Intraoperative Pharmacology table.

---

SeqNo: 6130

Long Name: Intraoperative Pharmacology Link to Operations Table

Short Name: OperationID

Database Table Name: IntraopPharm

Data Source: Automatic

Format: Text

Definition: An arbitrary, unique value generated by the software that permanently identifies each operation record in the participant's database. This field is the foreign key that links the Intraoperative Pharmacology record with the associated record in the Operations

table.

---

SeqNo: 6140  
Long Name: IntraOperative Pharmacology (Including CPB)  
Short Name: IntraopPharm  
Database Table Name: IntraopPharm  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the medications that were given during the intraoperative time period.

---

## 25. Anesthesia Pharmacology On Arrival To ICU/PACU

SeqNo: 6150  
Long Name: ICU Pharmacology Table Unique Record Identifier  
Short Name: ICUPUniqueID  
Database Table Name: ICUPharm  
Data Source: Automatic  
Format: Text  
Definition: Unique identifier for the record in the ICU Pharmacology table.

---

SeqNo: 6160  
Long Name: ICU Pharmacology Link to Operations Table  
Short Name: OperationID  
Database Table Name: ICUPharm  
Data Source: Automatic  
Format: Text  
Definition: An arbitrary, unique value generated by the software that permanently identifies each operation record in the participant's database. This field is the foreign key that links the ICU Pharmacology record with the associated record in the Operations table.

---

SeqNo: 6170  
Long Name: ICU/PACU Arrival Pharmacology  
Short Name: ICUPharm  
Database Table Name: ICUPharm  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the medications that were given to the patient on arrival to ICU (Intensive Care Unit) / PACU (Post Anesthesia Care Unit).

---

---

## 26. Anesthesia ICU/PACU Care

SeqNo: 6180

Long Name: ICU/PACU Arrival Date and Time

Short Name: ICUArrDT

Database Table Name: Operations

Data Source: User

Format: Date/Time - mm/dd/yyyy hh:mm

Definition: Indicate the date (mm/dd/yyyy) and time (hh:mm 24-hour clock) the patient arrived to the ICU / PACU.

---

SeqNo: 6190

Long Name: Initial FiO2

Short Name: InitialFiO2

Database Table Name: Operations

Data Source: User

Format: Real

Definition: Indicate the initial FiO2 (closest to the patient's arrival).

Low Value: 0.17    High Value: 1.0

---

SeqNo: 6200

Long Name: Mechanical Circulatory Support (ECMO/VAD)

Short Name: MechCircSup

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether the patient was on extracorporeal membrane oxygenation (ECMO) or on Ventricular Assist Device (VAD) on arrival.

---

SeqNo: 6211

Long Name: ICU/PACU Arrival Labs

Short Name: ICUPACULabs

Database Table Name: Operations

Data Source: User

Format: Text (categorical values specified by STS)

Definition: Indicate whether lab tests were drawn upon arrival to PACU or ICU.

---

SeqNo: 6220  
Long Name: pH  
Short Name: pH  
Database Table Name: Operations  
Data Source: User  
Format: Real  
Definition: Indicate the pH level from the first ABG obtained.

Low Value: 6.00      High Value: 8.00  
Parent Long Name: ICU/PACU Arrival Labs  
Parent Short Name: ICUPACULabs  
Parent Value(s): = "Yes"

---

SeqNo: 6230  
Long Name: pCO2  
Short Name: pCO2  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the pCO2 level from the first ABG obtained.

Low Value: 20      High Value: 150  
Parent Long Name: ICU/PACU Arrival Labs  
Parent Short Name: ICUPACULabs  
Parent Value(s): = "Yes"

---

SeqNo: 6240  
Long Name: pO2  
Short Name: pO2  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the pO2 level from the first ABG obtained.

Low Value: 15      High Value: 650  
Parent Long Name: ICU/PACU Arrival Labs  
Parent Short Name: ICUPACULabs  
Parent Value(s): = "Yes"

---

SeqNo: 6250  
Long Name: Base Excess

---

Short Name: BaseExcess  
Database Table Name: Operations  
Data Source: User  
Format: Integer  
Definition: Indicate the Base Excess level from the first ABG obtained.

Low Value: -30      High Value: 30  
Parent Long Name: ICU/PACU Arrival Labs  
Parent Short Name: ICUPACULabs  
Parent Value(s): = "Yes"

---

SeqNo: 6260  
Long Name: Lactate  
Short Name: Lactate  
Database Table Name: Operations  
Data Source: User  
Format: Real  
Definition: Indicate the Lactate level from the first ABG obtained.

Low Value: 0.1      High Value: 30.0  
Parent Long Name: ICU/PACU Arrival Labs  
Parent Short Name: ICUPACULabs  
Parent Value(s): = "Yes"

---

SeqNo: 6270  
Long Name: Hematocrit  
Short Name: Hematocrit  
Database Table Name: Operations  
Data Source: User  
Format: Real  
Definition: Indicate the hematocrit level from the first ABG obtained.

Low Value: 5.0      High Value: 70.0  
Parent Long Name: ICU/PACU Arrival Labs  
Parent Short Name: ICUPACULabs  
Parent Value(s): = "Yes"

---

SeqNo: 6280  
Long Name: Initial Pulse Oximeter  
Short Name: InitPulseOx  
Database Table Name: Operations  
Data Source: User

---

Format: Real  
Definition: Indicate the first pulse oximeter measurement after arrival to ICU / PACU.

Low Value: 50.0 High Value: 100.0

---

SeqNo: 6290  
Long Name: Temperature ICU/PACU Arrival  
Short Name: TempICUArr  
Database Table Name: Operations  
Data Source: User  
Format: Real  
Definition: Indicate the patient's temperature in degrees centigrade on arrival to the ICU/PACU.  
  
Low Value: 30.0 High Value: 43.0

---

SeqNo: 6300  
Long Name: Temperature Measurement Site  
Short Name: TempSite  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the location where the patient's temperature was measured.  
  
Parent Long Name: Temperature ICU/PACU Arrival  
Parent Short Name: TempICUArr  
Parent Value(s): Is Not Missing

---

SeqNo: 6310  
Long Name: Temporary Pacemaker on Arrival In ICU/PACU  
Short Name: TempPace  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the need for a temporary pacemaker on arrival to the ICU/PACU.

---

SeqNo: 6320  
Long Name: Temporary Pacemaker Site  
Short Name: TempPaceSite  
Database Table Name: Operations  
Data Source: User

---

Format: Text (categorical values specified by STS)  
Definition: Indicate the site of the temporary pacemaker.  
  
Parent Long Name: Temporary Pacemaker on Arrival In ICU/PACU  
Parent Short Name: TempPace  
Parent Value(s): = "Yes"

---

SeqNo: 6330  
Long Name: Type of Temporary Pacing  
Short Name: TempPaceType  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the type of temporary pacing.  
  
Parent Long Name: Temporary Pacemaker on Arrival In ICU/PACU  
Parent Short Name: TempPace  
Parent Value(s): = "Yes"

---

SeqNo: 6340  
Long Name: Disposition Under Anesthesia  
Short Name: DispUnderAnes  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate patient disposition after completion of anesthetic management.

---

SeqNo: 6350  
Long Name: Peri-Anesthetic Demise (Within 24 Hours of Last Anesthesia End Time)  
Short Name: PeriAnesDemise  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether the patient died within 24 hours of end of anesthesia.

---

## 27. Anesthesia Adverse Events

SeqNo: 6360  
Long Name: Anesthesia Adverse Events Unique Record Identifier

---



Short Name: AAEUniqueID  
Database Table Name: AAdvEvents  
Data Source: Automatic  
Format: Text  
Definition: Unique identifier for the record in the Anesthesia Adverse Events table.

SeqNo: 6370  
Long Name: Anesthesia Adverse Events Link to Operation Table  
Short Name: OperationID  
Database Table Name: AAdvEvents  
Data Source: Automatic  
Format: Text  
Definition: An arbitrary, unique value generated by the software that permanently identifies each operation record in the participant's database. This field is the foreign key that links the Anesthesia Adverse Events record with the associated record in the Operations table.

---

SeqNo: 6380  
Long Name: Anesthesia Adverse Event  
Short Name: AnesAdvEvent  
Database Table Name: AAdvEvents  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate the anesthesia-related adverse events that occurred.

---

SeqNo: 6381  
Long Name: Anesthesia Adverse Event - Additional Intervention Required  
Short Name: AnesAdvEventInt  
Database Table Name: AAdvEvents  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: Indicate whether additional intervention was required as a result of this adverse event.

Parent Long Name: Anesthesia Adverse Event  
Parent Short Name: AnesAdvEvent  
Parent Value(s): Is Not "None" And Is Not Missing

---

## 28. STS Temporary Fields

SeqNo: 6721  
Long Name: Temporary Yes/No Field #1  
Short Name: TempYN1

---

Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: This is a temporary field that should not be used for data collection until expressly instructed to by the STS.

SeqNo: 6722

Long Name: Temporary Yes/No Field #2  
Short Name: TempYN2  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: This is a temporary field that should not be used for data collection until expressly instructed to by the STS.

SeqNo: 6723

Long Name: Temporary Date Field  
Short Name: TempDt  
Database Table Name: Operations  
Data Source: User  
Format: Date - mm/dd/yyyy  
Definition: This is a temporary field that should not be used for data collection until expressly instructed to by the STS.

---

SeqNo: 6724

Long Name: Temporary Coded Field  
Short Name: TempCode  
Database Table Name: Operations  
Data Source: User  
Format: Text (categorical values specified by STS)  
Definition: This is a temporary field that should not be used for data collection until expressly instructed to by the STS.

---

SeqNo: 6725

Long Name: Temporary Text Field  
Short Name: TempText  
Database Table Name: Operations  
Data Source: User  
Format: Text  
Definition: This is a temporary field that should not be used for data collection until expressly instructed to by the STS.

---