



**The Society of Thoracic Surgeons**  
633 N. St. Clair, Suite 2320  
Chicago, IL 60611  
[www.sts.org](http://www.sts.org)

# News

## STS Press Release

### Media Contact:

Jennifer Bagley

312-202-5865

[jbagley@sts.org](mailto:jbagley@sts.org)

## Obese Heart Surgery Patients Require Significantly More ICU Resources

*Growing obesity problem prompts closer look at ways to maximize critical care resource management*

CHICAGO (August 10, 2017) — After heart surgery, obese patients tend to require additional intensive care unit (ICU) services and longer recovery times when compared to non-obese patients. This results in more expensive, more labor-intensive care, according to a study published online today in *The Annals of Thoracic Surgery*.

“Obesity is a growing problem for society that has reached epidemic proportions,” said lead author Brandon R. Rosvall, BSc, of Dalhousie Medicine New Brunswick in Canada. “In our study, we saw that as patients became more obese, the hospital resources required to care for them after heart surgery also increased.”

Rosvall, along with senior author Ansar Hassan, MD and other colleagues, used data from the New Brunswick Heart Centre Cardiac Surgery Database to identify patients who underwent cardiac surgery at the New Brunswick Heart Centre between January 2006 and December 2013. The authors also examined data specific to the patients’ ICU stays, which were obtained from logbooks and individual charts.

Of the 5,365 patients included in the final analysis, 1,948 (36%) were classified as obese. The patients were grouped into the following weight categories, as defined by the World Health Organization: Obese Class I (BMI 30.00-34.99), n=1,363 (25%); Obese Class II (BMI 35.00-39.99), n=441 (8%); and Obese Class III (BMI greater than 40.00), n=144 (3%). Body mass index (BMI) is a measure of body fat based on weight in relation to height. In general, the higher the BMI number, the more body fat a person has.

The study showed that following surgery, patients with higher levels of obesity were four times more likely to require extra time in the ICU, three times more likely to need additional time on mechanical ventilation, and three times more likely to be readmitted to the ICU. Researchers also learned that these patients experienced longer overall hospital lengths of stay and discharges with home care.

“The consistent relationship we have shown between increased BMI and these primary adverse outcomes confirms the robust nature of our findings,” said Rosvall.

### Key Points

- Obesity may negatively impact recovery from cardiac surgery.
- The larger the patient, the greater the hospital resources required to care for him/her after heart surgery.
- Patients with greater obesity have significantly increased risk of spending more time in the ICU and requiring more time on mechanical ventilation.

According to the Centers for Disease Control and Prevention (CDC), more than one-third (36.5%) of American adults are obese. The estimated yearly medical cost of obesity in the US reached \$147 billion in 2008 (the latest data available), which translates to \$1,429 more for each obese patient than a patient of normal weight.

“The ICU provides a number of highly specialized services to care for patients who are seriously ill,” said Rosvall. “Expensive resources including staff, medical equipment, and medication are needed to provide these services. Health care is costly, so by more efficiently using ICU resources, we can save the health care system money, while also improving overall patient care.”

Researchers said that strategies should be developed that improve ICU resource utilization among patients with increased BMI. For example, knowing that obese patients spend more time in the ICU after their operations enables health care providers to better predict ICU bed vacancies and be more thoughtful when scheduling staff and surgeries. Rosvall explained that efficient booking allows more patients to be treated while preventing overworked staff and canceled procedures. In addition, there are preemptive actions that doctors should put into practice, such as closer monitoring of obese patients and preoperative discussions about the unique surgical risks that they may face.

“Patients should be aware that obesity may negatively impact recovery from heart surgery,” said Rosvall. “More in-depth conversations regarding surgical risks and alternatives to surgery should take place with obese patients so that they are aware of the true impact of increased obesity. While obesity has become a social issue, it is one that can be addressed on a personal level. By working with their health care providers, patients can learn how to achieve and maintain a healthy body weight.”

According to Rosvall, this research group is currently conducting research on biomarkers (specific molecules in the blood) that will help doctors predict how obese patients will recover from cardiac surgery.

###

Rosvall BR, Forgie K, MacLeod JB, Yip AM, Aguiar C, Lutchmedial S, Brown C, Forgie R, Légaré JF, and Hassan A. Impact of Obesity on Intensive Care Unit Resource Utilization Following Cardiac Surgery. DOI: 10.1016/j.athoracsur.2017.05.047.

**Find comprehensive medical information presented for patients by leading experts in cardiothoracic surgery at the STS Patient Website ([ctsurgerypatients.org](http://ctsurgerypatients.org)). For a copy of *The Annals* article, contact Jennifer Bagley at 312-202-5865 or [jbagley@sts.org](mailto:jbagley@sts.org).**

Founded in 1964, The Society of Thoracic Surgeons is a not-for-profit organization representing more than 7,300 cardiothoracic surgeons, researchers, and allied health care professionals worldwide who are dedicated to ensuring the best possible outcomes for surgeries of the heart, lung, and esophagus, as well as other surgical procedures within the chest. The Society’s mission is to enhance the ability of cardiothoracic surgeons to provide the highest quality patient care through education, research, and advocacy.

*The Annals of Thoracic Surgery* is the official journal of STS and the Southern Thoracic Surgical Association. It has an impact factor of 3.700.